

Research Article

Effects of geraniol and citronellal compounds on biochemical and reproductive parameters of *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

Carolina A. Guedes^{1*}, Valéria Wanderley-Teixeira², Glaucilane Dos Santos Cruz¹, Kamilla De Andrade Dutra¹, Clóvis José Cavalcanti Lapa Neto², Álvaro A. C. Teixeira² and José V. De Oliveira¹

1. Federal Rural University of Pernambuco, Department of Agronomy- Entomology, Dom Manoel de Medeiros Ave., Dois Irmãos, 52171-900, Recife-PE, Brazil.

2. Federal Rural University of Pernambuco, Department of Animal Morphology and Physiology, Dom Manoel de Medeiros Av. Dois Irmãos, 52171-900, Recife-PE, Brazil.

Abstract: The essential oils of plants are composed of several volatile compounds, which may have different concentrations and may be determinant for their toxicity. Thus, in this work, sublethal and lethal effects of the compounds, geraniol and citronellal on the biochemical and reproductive parameters of *Spodoptera frugiperda* were investigated. For the geraniol compound the LD₃₀ of 9.42 mg/g and LD₅₀ of 13.65 mg/g was used, while for the citronellal LD₃₀ of 0.06 mg/g and LD₅₀ of 0.08 mg/g. Pure acetone was used in the control. Third instar caterpillars were treated topically in the prothoracic region by applying 1 µl of the respective compounds with a HamiltonTM 50 µl syringe. After 48 h the caterpillars were macerated in sodium phosphate buffer at a ratio of 4 caterpillars / 5 ml of the buffer, the levels of total protein, total sugar, lipid and glycogen were measured. The daily posture was evaluated in order to determine the periods of pre-oviposition, oviposition, post-oviposition and egg quantity. The results showed a reduction in the concentration of proteins and sugars for both compounds and concentrations studied. There was no change in lipid concentration. Citronellal increased the concentration of glycogen for both concentrations. The oviposition period and the number of eggs were reduced. However, there was no difference for the pre-oviposition and post-oviposition periods. Thus, it is inferred that geraniol and citronellal compounds cause alterations in the biochemical parameters that reflect in the reproduction of *S. frugiperda*.

Keywords: armyworm, isolated compounds, nutritional constituents, reproduction

Introduction

The high corn yield can be obtained by efficiently controlling its main pest, *Spodoptera frugiperda* (J.E. Smith). However, in Brazil the

predominance of synthetic insecticides is still recurrent, which are used without taking into account the principles of Integrated Pest Management. Thus, the adoption of insecticides with low environmental impact, such as insecticidal plants, is fundamental to the success of pest control (Diéz-Rodrigues and Omoto, 2001).

One of the classes of plant-derived compounds, which has been prominent in insect

Handling Editor: Saeid Moharramipour

*Corresponding author, e-mail: carolguedes84@yahoo.com.br

Received: 16 November 2017 Accepted: 8 May 2018

Published online: 20 July 2018

control, are the essential oils, which are already part of formulations of pesticides, capable of killing and repelling insects (Isman, 2000). Some essential oils have already been evaluated for *S. frugiperda*, such as citronella, *Cymbopogon winterianus* Jowitt causing food deterrence and mortality (Labinas and Crocomo 2002). Extracts from other species such as *Foeniculum vulgare* (Mill.), *Ocimum gratissimum* L. and *Eucalyptus staigeriana* F. Muell., in sublethal doses interfered in biological parameters and *O. gratissimum* also altered reproductive parameters (Cruz *et al.*, 2017).

The essential oils are composed of several volatile compounds, which may have different concentrations and may be determinant for their toxicity (Jemaa *et al.*, 2012). Gusmão *et al.* (2013) and Silva *et al.* (2016), through chromatographic analyzes coupled to mass spectrometry, verified that the oil of *C. winterianus* presents the main constituents of geraniol and citronellal. These chemical compounds were also found by Cruz *et al.*, (2016) in *O. gratissimum* oil, in addition citronellal was also present in the oils of *F. vulgare*, *E. staigeriana* and *E. citriodora*. Geraniol and citronellal are terpene compounds that have been shown to be of agricultural interest and can act as repellents, insecticides and growth inhibitors (Isman, 2000; Hummelbrunner and Isman, 2001; Isman, 2006; Chen and Viljoen, 2010).

Studies that address the interference of isolated chemical compounds on biochemical parameters are relevant, since researches have shown that botanical insecticides may interfere with the absorption and metabolism of nutrient constituents of insects (Sharma *et al.*, 2011). Changes that occur in this sense can have negative consequences on the biological cycle of insects. Silva *et al.* (2016) verified that the *C. winterianus* oil in sublethal dose affects the biochemical profile of *S. frugiperda* caterpillars, causes reduction of proteins, lipids and total sugars, and increase of glycogen.

In the case of isolated or associated chemical compounds, Cruz *et al.* (2017) found that the chemical compounds limonene, trans-anethole

and limonene + trans-anethole reduced the amount of lipids, protein, total sugars and glycogen, showing greater expression when the compounds were associated, except glycogen. Therefore, chemical compounds of important essential oils in the control of *S. frugiperda* can be explored for a better elucidation of their effects on the pest. Thus, the present study aimed to evaluate the sublethal and lethal effects of geraniol and citronellal on biochemical and reproductive parameters of *S. frugiperda*.

Materials and Methods

The present research was developed in the Laboratory of Histology of the Department of Morphology and Animal Physiology, Federal Rural University of Pernambuco (UFRPE).

Obtaining and breeding insects

S. frugiperda caterpillars were obtained from the stock of the Agricultural Entomology Laboratory of the Department of Agronomy of the Federal Rural University of Pernambuco (UFRPE), maintained at a temperature of 25.2 ± 1.4 °C, relative humidity of $67 \pm 0.7\%$ and photophase of 12 h, and fed with modified Greene artificial diet, recommended for the species (Busato *et al.*, 2006).

Obtaining Chemical Constituents

The chemical compounds were obtained from Sigma-Aldrich® with purity of 99%.

Extraction and Quantification of Total Soluble Proteins

For the experiments, third instar caterpillars (10 days old) were used, with an average weight of 78mg. The treatments consisted of: dilution of acetone compounds in LD₃₀ and LD₅₀ previously estimated by Cruz *et al.* (2016). For the geraniol compound the LD₃₀ of 9.42 mg/g (7.38-10.92) and LD₅₀ of 13.65 (11.34-15.97) mg/g were used, while for the citronellal the LD₃₀ 0.06 (0.05-0.07) mg/g and LD₅₀ of 0.08 (0.07-0.08) mg/g. Pure acetone was used in the control. In the prothoracic region 1.0 µl of the

respective compounds were applied with the aid of a HamiltonTM 50 µl syringe. The caterpillars were individualized in flat bottom glass tubes and sealed with plastic film. Artificial diet was used for feeding. Each sample was composed of four caterpillars, and for each treatment 10 samples were obtained, totaling 40 caterpillars/treatment. The method of Bradford (1976) was used to determine the soluble proteins. The processing occurred after 48h of the installation of the experiments. The caterpillars were immobilized at 40 °C and then macerated in sodium phosphate buffer (pH 7.4 and 0.1 M) in the ratio of four caterpillars / 5 ml of the buffer. 1.0 ml of the mixture (caterpillars + buffer) was pipetted and stored in a properly labeled microtube. The procedure was carried out at low temperature to avoid oxidation of the samples. These were centrifuged for 3 minutes at 3000rpm. After centrifugation, 100 µl of each sample was taken and placed in glass tubes and 5 ml of Bradford dye (0.01% Comassie Blue G-250, 8.5% phosphoric acid and 4.7% ethanol). The tubes were homogenized in a vortex type stirrer and were allowed to stand for 2 min. The spectrophotometer was then read at a wavelength of 595 nm. The unit used was µg/mL. The results were submitted to ANOVA and the means were compared by the tukey test at the 5% probability level using SAS Proc GLM (SAS Institute 2001).

Analyses of lipid, total sugar and glycogen

3rd instar larvae were macerated in sodium phosphate buffer (pH 7.4 and 0.1 M) at a ratio of 4 caterpillars / 5 ml of the buffer. Each sample consisted of four caterpillars and for each treatment 10 samples were obtained, totaling 40 caterpillars/treatment. The content of lipid, total sugar and glycogen were evaluated using the Van Handel method (Van Handel, 1985a,b), where 200 µl of the homogenate was added to 200 µl of sodium sulfate plus 800 µl of methanol and chloroform (1: 1), And centrifuged at 2000 rpm for 2 min. The precipitate was used for the glycogen analysis, and the supernatant was transferred to a test tube where the lipid and sugar were separated. The lipid was analyzed

spectrophotometrically, using the phosphoric-vanillin acid method, whereas for the total sugar and glycogen, the sulfuric-anthrone acid method was used. The absorbance was read at 625 nm. The results of the lipid, sugar and glycogen contents were submitted to ANOVA and the means were compared by the tukey test at the 5% probability level using SAS ProcGLM (SAS Institute 2001).

Effect of compounds on reproduction of *S. frugiperda*

Third instars were subjected to the LDs 30 and 50 of the geraniol and citronellal compounds through topical contact. Then they were individualized in flat bottom glass tubes, fed daily with specific artificial diet until pupae formation. These were sexed and after adult emergence, couples were formed, two per cage with five replicates per treatment. The couples were housed in a PVC cage with dimensions of 10 cm x 15 cm (diameter and height) coated internally with sulphite paper, as a substrate for oviposition. The moths were fed with 10% honey solution and kept in an air-conditioned room. All experiments were performed at 25.2 ± 1.4 ° C, $67 \pm 0.7\%$ RH and 12 h photophase. The eggs laid were collected daily, counted and conditioned in Petri dishes with dimensions of 10 cm in diameter and maintained in the same conditions as previously described. Finally, the period of pre-oviposition, oviposition, total number of eggs and post-oviposition were evaluated. The analyses were submitted to the tukey test at 5% probability by SAS (SAS Institute 2001). The data were transformed into $(x + 1)^{1/2}$. Through confirmed mortality data the average survival rate was determined.

Results

Quantification of total soluble proteins

Biochemical tests for the extraction and quantification of total soluble proteins showed that the third instar larvae of *S. frugiperda* treated with LD₃₀ and LD₅₀ of the geraniol and citronellal compounds showed a reduction in their content (Fig. 1).




Figure 1 Amount of total proteins from *Spodoptera frugiperda* 3rd instar caterpillars subjected to LD₃₀ and LD₅₀, by topical contact of the geraniol and citronellal compounds.

Analysis of lipid, total sugar and glycogen

Neither of the two compounds nor the doses used affected the amount of lipid compared to the control (Fig. 2). For the analysis of total sugar, there was a significant reduction in the caterpillars of both treatments and concentrations (Fig. 3). Geraniol at LD₃₀ did not affect the glycogen content, whereas at LD₅₀ it induced its reduction. However, the citronellal compound at both doses caused an increase of glycogen (Fig. 4).




Figure 2 Quantity of total lipids derived from *Spodoptera frugiperda* 3rd instar caterpillars subjected to LD₃₀ and LD₅₀, by topical contact of the geraniol and citronellal compounds.




Figure 3 Amount of total sugar from *Spodoptera frugiperda* 3rd instar caterpillars submitted to LD₃₀ and LD₅₀, by topical contact of the geraniol and citronellal compounds.




Figure 4 Quantity of total glycogen from third instar caterpillars of *Spodoptera frugiperda* submitted to LD₃₀ and LD₅₀, by topical contact of the geraniol and citronellal compounds.

Effect of compounds on reproduction of *S. frugiperda*

The geraniol and citronellal compounds at LD₃₀ and LD₅₀ caused a significant reduction in the oviposition period and in the amount of eggs of adults from 3rd instar caterpillars treated by topical contact. However, there was no change in the pre-oviposition and post-oviposition periods (Table 1).

Table 1 Pre-oviposition period, oviposition, post-oviposition period and total number of eggs of two *Spodoptera frugiperda* females treated as larvae with LD₃₀ and LD₅₀ of the geraniol and citronellal compounds. Temperature: 25.2 ± 1.4 °C. Humidity: 67 ± 0.7% and 12 h photophase.

Treatments	Pre-oviposition (days) ¹	Oviposition (days) ¹	Post-oviposition (days) ¹	Number of eggs Per cage ¹
Control	2.40 ± 0.24a	7.60 ± 0.24a	2.60 ± 0.24a	2170.80 ± 197.02a
Geraniol LD ₃₀	3.60 ± 0.24a	5.00 ± 0.55b	2.20 ± 0.20a	704.20 ± 171.02b
Geraniol LD ₅₀	3.00 ± 0.32a	3.80 ± 0.37b	2.80 ± 0.86a	729.00 ± 257.02b
Citronellal LD ₃₀	2.60 ± 0.40a	3.40 ± 0.60b	3.60 ± 0.98a	598.40 ± 86.51b
Citronellal LD ₅₀	3.00 ± 0.32a	4.20 ± 0.37b	2.20 ± 0.58a	527.00 ± 88.95b

¹ Original data for analysis were transformed into $(x + 1)^{1/2}$, Means followed by the same letter in the columns do not differ significantly by the Tukey's test ($P < 0.05$).

Discussion

Geraniol and citronellal caused reduction in some biochemical parameters, such as protein and sugars. According to Sharma *et al.* (2011) and Yazdani *et al.* (2013) nutritional parameters may suffer interference caused by botanical insecticides, a fact that is in line with the results of the present research. Proteins participate in innumerable processes related to insect survival, and alterations in protein levels can cause negative reproductive effects, as they directly affect vitellogenesis (Panizzi and Parra, 2009; Guizzo *et al.*, 2012; Rosas-Mejía *et al.*, 2015). As verified in the research in question according to Senthilkumar *et al.* (2009) protein reduction is probably due to interference caused by compounds present in botanical insecticides that act on the hormones that regulate protein synthesis. Many of the insecticides have deterrent properties and reduce feed efficiency in insects, which in turn reduces some vital components like proteins in the body. The compounds may have inhibited the feeding of the larvae and as a result the protein decrease could be due to hunger stress (Etebari *et al.*, 2006).

The reduction in carbohydrate content may compromise insect survival (Parra, 1999), because these constituents function as an energy source, besides being responsible for innumerable metabolic and structural functions, and in the formation of chitin; Can be converted into lipids, as well as participate in amino acid synthesis (Chapman, 2013; Arrese *et al.*, 2010).

In insects, lipids play important roles, are constituents of cellular structures, act as hormones and form important energy reserves

of great metabolic activity such as flight and egg production (Arrese *et al.*, 2001). Although the results obtained in this research did not present negative changes for the lipids in either of the compounds and concentrations studied; oviposition and fecundity showed a significant reduction. These results may suggest that lipids were not metabolized and/or absorbed adequately for egg production in *S. frugiperda*.

It can be inferred, therefore, that compounds of essential oils have an effect on the nutritional indexes of treated larvae. This suggests a reduction in the efficiency of using ingested foods when these products are used (Etebari *et al.*, 2006).

In general, studies related to biochemical alterations are performed with extracts or essential oils, such as extracts of *Eucalyptus globulus* Labill. *Cymbopogon citratus* (DC.) Stapf, *Artemisia annua* L., *Gendarussian justice* Burm F., *Myristica fragrans* Houtt, *Annona squamosa* L., and *Centella asiatica*, which reduced rates of protein, carbohydrate, lipid and some amino acids of larvae and adults of *Anopheles stephensi* (Senthilkumar *et al.*, 2009).

According to tests performed by Nasr *et al.* (2017) the essential oil of *Origanum vulgare* L. significantly reduced the total protein and triglycerides of third instar larvae of *Plutella xylostella* L., in sublethal and lethal doses, corroborating with several other studies that point out the toxic properties of insect oils - Prague. These authors suggest that in physiological studies, determination of the total protein and of many chemical macromolecules such as lipids and carbohydrates are very important. Etebari *et al.* (2006) reiterated that

many products have anti-feedant properties and reduce the efficiency of insect feeding, which in turn reduces some vital components, such as proteins.

As regards effect of components of essential oils on the aforementioned macromolecules, Cruz *et al.* (2017) reported decreases in proteins, lipids, sugars and glycogen in *S. frugiperda* caterpillars when treated with limonene and trans-anethole compounds, and these results were more significant when the compounds were used in combination. In addition, the experiments showed different relationships with glycogen levels, since the citronellal compound caused an increase of glycogen corroborating with data obtained by Silva *et al.* (2016), who also observed increase of glycogen caused by *C. winterianus* oil. These authors suggested that the increase was caused by the citronellal, the major compound present in this oil which is the object of our study as well.

Regarding the reproductive aspects, it is possible to state that the geraniol and citronellal compounds had a negative effect on the oviposition and fecundity of *S. frugiperda*, confirming the correlation that the reproductive success of the insects is related to the nutrients acquired in the immature phase (Sousa *et al.*, 2015), since nutritional resources are acquired to ensure growth and reproduction. The vitellogenesis and maturation of ovules to occur satisfactorily depends on the energy resources that have been acquired. A reduction in reproduction shows that the compounds may have influenced the non-metabolization of most nutritional constituents or inefficiently acquired in the larval period (Attardo *et al.*, 2005; Milano *et al.*, 2010; Etebari *et al.*, 2006).

Other studies deal with the effects of compounds and essential oils on reproductive aspects such as Sharab *et al.* (2009), who verified that the compound eugenol and the oil of *Mentha piperita* L., each at concentration of 0.01% caused significant reduction in the fecundity of *Phthorimaea operculella* Zell. The same authors also tested leaves, fruits or seeds of 14 plants for their essential oils and verified that the oils of *Allium cepa* L., *Curcuma longa*

L., *Clolocasia antiquorum* Schott, *Ocimum basilicum* L., *Dodonaea viscosa* (L.) Jacq. and *Thuja orientalis* L. caused a reduction in egg deposition. (+) - fenchone and (-) - β - pinene significantly reduced the oviposition potential of *Pseudaletia unipuncta* Haworth females by 46% and 33% (Sousa *et al.*, 2015).

The present study showed efficacy with respect to the isolated compounds, whereas Hummelbrunner and Isman (2001) found that isolated compounds were less effective than conventional insecticides. On the other hand, when mixtures of some compounds were tested, they proved to be more effective than botanical insecticides such as rotenone and neem. These same authors verified that a mixture of R-terpineol, eugenol and thyme oil was 300 times less toxic to fish than azadirachtin, rotenone and pyrethrum. This can be attributed to detoxification pathways and mode of action of monoterpenoids (Koul *et al.* 2013). Several compounds of essential oils have been reported as octopamine blockers, which is a unique neurotransmitter for insects, and which is also why they are less toxic to vertebrates (Hummelbrunner and Isman, 2001). Compounds of essential oils may be considered safer for the environment than other plant-derived chemicals, such as azadirachtin, rotenone, or pyrethrum (Stroh *et al.*, 1998). In general, it is believed that essential oils by having a large number of compounds and thereby increasing the spectrum of insecticidal action are more effective than pure compounds. However, there are controversies regarding the efficacy of essential oil compounds, since different species have different responses to individual compounds.

Conclusions

It can be concluded from the present study that the geraniol and citronellal compounds cause reduction in biochemical and reproductive parameters, and thus can be used in pest control in ecologically based integrated management programs. In addition, the identification of essential oil compounds may allow the development of more effective control agents.

Acknowledgements

Thanks are expressed to: CAPES for granting the scholarship to the first author, making this research possible, Dmfa-Ufrpe (Department of Morphology and Animal Physiology) for the realization of the bioassays, Sigma-Aldrich® for obtaining the chemical constituents.

References

- Arrese, E. L., Canavoso, L. E., Jouni, Z. E., J. E. Pennington, K. Tsuchida and M. A. Ells. 2001. Lipid storage and mobilization in insects: current status and future directions. *Insect Biochemistry and Molecular Biology*, 31 (1): 7-17.
- Arrese, E. L., Howard, A. D., Patel, R. T., Rimoldi, O. J. and Soulages, J. L. 2010. Mobilization of lipid stores in *Manduca sexta*: cDNA cloning and developmental expression of fat body triglyceride lipase, TGL. *Insect Biochemistry and Molecular Biology*, 40 (2): 91-99.
- Attardo, G. M., Hansen, I. A. and Raikhel, A. S. 2005. Nutricional regulation of vitellogenesis in mosquitoes: implications for anautogeny. *Insect Biochemistry and Molecular Biology*, 35 (7): 661-675.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72 (7): 248-254.
- Busato, G. R., Garcia, M. S., Loeck, A. E., Zart, M., Nunes, A. M., Bernardi, O. and Andersson, F. S. 2006. Adequação de uma dieta artificial para os biótipos “milho” e “arroz” de *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Bragantia*, 65 (2): 317-323.
- Chapman, R. F. 2013. *The Insects Structure and Function*. Cambridge, Cambridge University Press.
- Chen, W. and Viljoen. A. M. 2010. Geraniol-A review of commercially important fragrance material. *South African Journal of Botany*, 76 (4): 643-651.
- Cruz, G. S. Wanderley-Teixeira, V., Oliveira, J. V., D’assunção, C. G., Cunha, F. M., Teixeira, A. A. C., Guedes, C. A., Dutra, K. A., Barbosa, D. R. S. and Breda, M. O. 2017. Effect of trans-anethole, limonene and your combination in nutritional componentes and their reflection on reproductive parameters and testicular apoptosis in *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Chemico-Biological Interactions*, 263 (1): 74-80.
- Cruz, G. S. Wanderley-Teixeira, V., Oliveira, J. V., Lopes, F. S. C., Barbosa, D. R. S., Breda, M. O., Dutra, K. A., Guedes, C. A., Navarro, D. M. A. F. and Teixeira, A. A. C. 2016. Sublethal effects of essential oils from *Eucalyptus staigeriana* (Myrtales: Myrtaceae), *Ocimum gratissimum* (Lamiales: Lamiaceae), and *Foeniculum vulgare* (Apiaceae) on the biology of *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*, 109 (2): 660-666.
- Diéz-Rodrigues, G. I. and Omoto, C. 2001. Herança da resistência de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) a lambda-cialotrina. *Neotropical Entomology*, 30 (2): 311-316.
- Etebari, K., Bizhannia, A. R., Sorati, R. and Matindoost, L. 2006. Biochemical changes in haemolymph of silkworm larvae due to pyriproxyphen residue. *Pesticide Biochemistry and Physiology*, 88 (1): 14-19.
- Guizzo, M. G., Abreu, L., Masuda, A., Logullo, C. and Junior, I. S. V. 2012. Metabolism of Biomolecules in the Embryogenesis of the Tick *Rhipicephalus (Boophilus) microplus*. *Acta Scientiae Veterinariae*, 40 (1): 1010-1022.
- Gusmão, N. M. S., Oliveira, J. V., Navarro, D. M. A. F., Dutra, K. A., Silva, W. A. and Wanderley, M. J. A. 2013. Contact and fumigant toxicity and repellency of *Eucalyptus citriodora* Hook., *Eucalyptus staigeriana* F., *Cymbopogon winterianus* Jowitt and *Foeniculum vulgare* Mill. essential oils in the management of *Callosobruchus maculatus* (Fabr.) (Coleoptera: Chrysomelidae, Bruchinae). *Journal of Stored Products Research*, 54: 41-47.

- Hummelbrunner, L. A. and Isman, M. B. 2001. Acute, Sublethal, Antifeedant, and Synergistic Effects of Monoterpenoid Essential Oil Compounds on the Tobacco Cutworm, *Spodoptera litura* (Lep., Noctuidae). *Journal of Agricultural and Food Chemistry*, 49 (2): 715-720.
- Isman, M. B. 2000. Plant essential oils for pest and disease management. *Crop Protection*, 19 (8-10): 603-608.
- Isman, M. B. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology*, 51: 45-66.
- Jemâa, J. M. B., Haquel, S., Bouaziz, M. and Khouja, M. L. 2012. Seasonal variations in chemical composition and fumigant activity of five Eucalyptus essential oils against three moth pests of stored dates in Tunisia. *Journal of Stored Products Research*, 48: 61-67.
- Koul, O., Singh, R., Kaur, B. and Kanda, D. 2013. Comparative study on the behavioral response and acute toxicity of some essential oil compounds and their binary mixtures to larvae of *Helicoverpa armigera*, *Spodoptera litura* and *Chilo partellus*. *Industrial Crops and Products*, 49: 428-436.
- Labinas, M. A. and Crocomo, W. B. 2002. Effect of java grass (*Cymbopogon wintwrianus*) essential oil on fall armyworm *Spodoptera frugiperda* (J. E. Smith, 1979) (Lepidoptera: Noctuidae). *Acta Scientiae*, 24: 1401-1405.
- Milano, P., Berti Filho, E., Parra, J. R. P., Oda, M. L., and Cônsoli, F. L. 2010. Efeito da alimentação da fase adulta na reprodução e longevidade de espécies de Noctuidae, Crambidae, Tortricidae e Elachistidae. *Neotropical Entomology*. 39 (2): 172-180.
- Nasr, M., Sendi, J. J., Moharramipour, S. and Zibae, A. 2017. Evaluation of *Origanum vulgare* L. essential oil as a source of toxicant and an inhibitor of physiological parameters in diamondback moth, *Plutella xylostella* L. (Lepidoptera: Pyralidae). *Journal of the Saudi Society of Agricultural Sciences*, 16 (2): 184-190.
- Panizzi, A. R. and Parra, J. R. P. 2009. Bioecologia e Nutrição de Insetos: Base Para o Manejo Integrado de Pragas. Brasília, Embrapa.
- Parra, J. R. P. 1999. Técnicas de Criação de Insetos Para Programas de Controle Biológico. Piracicaba, FEALQ.
- Rosas-Mejía, M., Correa-Sandoval, A., Venegas-Barrera, C. S. and Horta-Veja, J. V. 2015. Preferências entre cinco carboidratos en *Pheidole bilimeki* (Hymenoptera: Formicidae). *Acta Zoologica Mexicana*, 31 (2): 291-297.
- SAS Institute. 1999-2001. SAS/STAT User's Guide, Version 8.02, TS level 2MO. SAS Institute Inc., Cary, NC.
- Senthilkumar, N., Varma, P. and Gurusubramanian, G. 2009. Larvidal and adulticidal activities of some medicinal plants against the Malarial Vector, *Anopheles stephensi* (Liston). *Parasitology Research*, 104 (2): 237-244.
- Sharaby, A., Abdel-Rahman, H. and Moawad, S. 2009. Biological effects of some natural and chemical compounds on the potato tuber moth, *Phthorimaea operculella* Zell. (Lepidoptera: Gelechiidae). *Saudi Journal of Biological Sciences*, 16 (1): 1-9.
- Sharma, P., Mohan, L., Dua, K. K. and Srivastava, C. N. 2011. Status of carbohydrate, protein and lipid profile in the mosquito larvae treated with certain phytoextracts. *Asian Pacific Journal of Tropical Medicine*, 4 (4): 301-304.
- Silva, C. T. S., Wanderley-Teixeira, V., Cunha, F. M., Oliveira, J. V., Dutra, K. A., Navarro, D. M. A. F. and Teixeira, A. A. C. 2016. Biochemical parameters of *Spodoptera frugiperda* (J. E. Smith, 1979) treated with citronella oil (*Cymbopogon wintwrianus* Jowitt ex Bor) and its influence on reproduction. *Acta Histochemica*, 118: 347-352.
- Sousa, R. M., Rosa, J. S., Oliveira, L., Cunha, A. and Fernandes-Ferreira, M. 2015. Activities of Apiaceae essential oils and volatile compounds on hatchability, development, reproduction and nutrition of *Pseudaletia unipuncta* (Lepidoptera: Noctuidae). *Industrial Crops and Products*, 63: 226-237.

- Stroh, J., Wan, M. T., Isman, M. B. and Moul, D. J. 1998. Evaluation of the acute toxicity to juvenile Pacific, *Coho salmon* and rainbow trout of some plant essential oils, a formulated product, and the carrier. *Bulletin of Environmental Contamination and Toxicology*, 60 (6): 923-930.
- Van Handel, E. 1985a. Rapid determination of total lipids in mosquitoes. *Journal of the American Mosquito Control Association*. 1 (3): 302-304.
- Van Handel, E. 1985b. Rapid determination of glycogen and sugars in mosquitoes. *Journal of the American Mosquito Control Association*. 1 (3): 299-301.
- Yazdani, E., Sendi, J. J. Aliakbar, A. and Senthil-Nathan, S. 2013. Effect of *Lavandula angustifolia* essential oil against lesser mulberry pyralid *Glyphodes pyloalis* Walker (Lep: Pyralidae) and identification of its major derivatives. *Pesticide Biochemistry and Physiology*, 107 (2): 250-257.

اثرات ترکیبات ژرانیول و سیترونلال روی پارامترهای بیوشیمیایی و تولید مثلی *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

کارولینا گوندس^{۱*}، والریا وندرلی تیکسرا^۲، گلوسیلا، دوس سانتوس کروزا^۱، کامیلا آندره دوترا^۱، کلویس خوزه کاوالکانتیلاپا ناتو^۲، آلوارو تیکسریا^۲ و جوزف اولیویرا^۱

۱- دانشگاه فدرال پرنامبوکو، دانشکده زراعت و حشره‌شناسی، رسیف، برزیل.

۲- دانشگاه فدرال پرنامبوکو، گروه مورفولوژی و فیزیولوژی دامی، رسیف، برزیل.

* پست الکترونیکی نویسنده مسئول مکاتبه: carolguedes84@yahoo.com.br

دریافت: ۲۵ آبان ۱۳۹۶؛ پذیرش: ۱۸ اردیبهشت ۱۳۹۷

چکیده: اسانس‌های گیاهی از ترکیبات فرآری تشکیل شده‌اند که غلظت‌های مختلف آن می‌تواند خاصیت سمی و کشنده روی حشرات آفت داشته باشد. لذا در این پژوهش، اثرات کشنده و زیرکشنده ترکیبات ژرانیول و سیترونلال روی پارامترهای بیوشیمیایی و تولیدمثلی *Spodoptera frugiperda* مورد بررسی قرار گرفت. برای ترکیب ژرانیول مقدار LD₃₀ ۹/۴۲ میلی گرم بر گرم و LD₅₀ ۱۳/۶۵ میلی گرم بر گرم استفاده شد، درحالی که برای سیترونلال از LD₃₀ ۰/۰۶ میلی گرم بر گرم و LD₅₀ ۰/۰۸ میلی گرم بر گرم استفاده شد. از استون خالص به عنوان شاهد استفاده شد. با استفاده از سرنگ ۵۰ میکرولیتری همیلتون مقدار ۱ میلی لیتر از هر غلظت روی پیش قفس سینه لارو سن سوم قرار داده شد. پس از ۴۸ ساعت، لاروها در محلول بافر سدیم فسفات با نسبت ۴ لارو در ۵ میلی لیتر بافر همگن سازی شدند. سپس مقدار پروتئین کل، قند کل، چربی و گلیکوژن تعیین شدند. همچنین به طور روزانه طول دوره پیش از تخم گذاری، دوره تخم گذاری، دوره پس از تخم گذاری و تعداد تخم‌های گذاشته شده تعیین گردید. نتایج نشان داد که غلظت پروتئین کل و قند کل برای هر دو ترکیب در غلظت‌های مورد بررسی کاهش یافته است. هیچ تغییری در غلظت چربی مشاهده نشد. اما سیترونلال مقدار گلیکوژن را در هر دو غلظت افزایش داد. دوره تخم گذاری و تعداد تخم‌های گذاشته شده کاهش یافت. با این حال، دوره‌های قبل از تخم گذاری و بعد از تخم گذاری اختلافی نداشت. بنابراین نتیجه گیری می‌شود که ترکیبات ژرانیول و سیترونلال باعث تغییراتی در پارامترهای بیوشیمیایی می‌شوند که این امر در تولیدمثل حشرات کامل *S. frugiperda* مؤثر است.

واژگان کلیدی: لارو، ترکیبات استخراج شده، ترکیبات غذایی، تولیدمثل