Research Article

Using ISSR and URP-PCR markers in detecting genetic diversity among *Macrophomina phaseolina* isolates of sesame in Iran

Reza Salahlou¹, Naser Safaie^{2*} and Masoud Shams-Bakhsh²

1. Department of Biotechnology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

2. Department of Plant Pathology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

Abstract: To assess the variability of *Macrophomina phaseolina* (Tassi) Goid, the causal agent of charcoal rot of Sesame, sixty isolates recovered from ten geographic regions, were analyzed using inter simple sequence repeat (ISSR) and universal rice primer (URP) markers. Isolates were grouped into eight clusters at 78% genetic similarity level. Our results showed that the five ISSR primers produced 105 bands of which 77.11% were polymorphic and eight URP-PCR primers generated 135 bands of which 66.84% were polymorphic. These methods showed a considerable genetic diversity among Iranian isolates, but no correlation was found between genetic diversity and geographical origins of the isolates. Analysis of molecular variance revealed that a large proportion of genetic variability resulted from the differences among isolates within regions. The findings of this study demonstrated that the low-genetic differentiation (G_{ST}) and high gene flow (N_m) among populations had a significant effect on the emergence and evolutionary development of *M. phaseolina*.

Keywords: charcoal rot, *Sesamum indicum*, DNA fingerprinting, population genetics

Introduction

Charcoal rot is one of the factors limiting the cultivation of Sesame *Sesamum indicum* L., an oilseed crop widely grown in different parts of the world. *Macrophomina phaseolina* (Tassi) Goid. a soil-and seed-borne fungus causes charcoal rot. The fungus can infect the root and lower stem of more than 500 plant families and has a wide geographic distribution (Dhingra and Sinclair, 1978). Charcoal rot is an important soil borne disease and is favored by hot and dry weather or when unfavorable environmental conditions impose stress on plant (Etebarian, 2006).

In order to design optimal breeding methods to produce sesame cultivars which can have resistance to charcoal rot, knowledge of population biology and its interaction with sesame cultivar would be quite useful. The use of resistant varieties is the most appropriate approach to control of M. phaseolina (Ahmad and Burney, 1990). The existence of high level variations in a fungal population increase its adaptation to various conditions and overcomes host resistance deployed by growers (Yeh et al., 1999). To determine the level of variability in all organisms, basic morphological traits and pathogenesis as the first character are used, but usually these methods are time-consuming and their quantification is difficult. Molecular methods are an appropriate analysis approach for a rapid evaluation of the variability within among species (Chakravarthi and and Naravaneni, 2006).



Handling Editor: Vahe Minassian

^{*} Corresponding author, e-mail: nsafaie@modares.ac.ir Received: 18 October 2018, Accepted: 15 June 2019 Published online: 21 July 2019

The host correlates development of disease resistance with the accumulation of new polypeptides syntheses (Broglie *et al.*, 1986). The new protein contents depend on sensitivity to infection, host genotype and the virulence genes of the pathogens (Hlinkova and Sykora, 1996; Radwan, 2000). Pecina *et al.* (2000) reported Double-stranded RNA (dsRNA) in *M. phaseolina* with sizes ranging from 0.4kbp to 10kbp and the number of dsRNA ranging from 1 to 10.

In recent years, different DNA markers have been employed to study genetic diversity within populations of *M. phaseolina* such as Restriction Fragment Length Polymorphism (RFLP) of rDNA-ITS regions (Aghakhani and Dubey, 2009; Almeida et al. 2003), Random Polymorphic Amplified DNA (RAPD) (Aboshosha et al. 2007; Aghakhani and Dubey, 2009; Almeida et al. 2003), Amplified Fragment Length Polymorphism (AFLP) (Linhai et al. 2011; Mayek-Perez et al. 2001; Reyes-Franco et al. 2006), Universal Rice Primer PCR (URP-PCR) (Jana et al., 2005b), ISSR (Jana et al., 2005a), Repetitive Sequence-Based Polymerase Chain Reaction (Rep-PCR) (Purkayastha et al., 2008) and Simple Sequence Repeats (SSR) (Baird et al., 2010).

Among DNA markers, ISSR is a dominant molecular marker system that includes the use of one primer complementary to a target Simple Sequence Repeat (SSR) region in PCR and allows DNA amplification between SSR regions. This technique was first reported as marker to assess genetic diversity and taxonomic studies in animals, plants and later used to obtain DNA markers in fungi (Bornet and Branchard, 2001; Menzies et al. 2003; Zietkiewicz et al. 1994). The advantages of ISSR over other markers include; (1) low development costs; (2) no prior information or lengthy mapping studies are required; (3) laboratory procedures can easily be applied to any plant species and fungi (Geo et al., 2006; Maleki et al., 2018).

URP-PCR markers are powerful tools which can be utilized to examine genetic diversity of various organisms including Micro-organisms, animals and plants at intraspecific and interspecific levels and were designed based on repetitive sequence of rice genome by Kang *et al.* (2002).

In Iran, dry rot occurs in many economically important crops, including common bean, sesame, soybean, chickpea, sunflower and sugarbeet, among others, and it might lead to yield reduction (Mahdizadeh *et al.*, 2011). Seed infection by this pathogen may be responsible for its spreading in the main agricultural regions of the country. Infection to this pathogen has been reported in different region of Iran (Ershad, 1995). Although *M. phaseolina* has been present for decades in Iran and a vast literature about its agronomic aspects is available, information about the population biology of this pathogen is still interesting subject.

The objective of this study was to assess genetic diversity and population structure of M. *phaseolina* pathogen of sesame in Iran, using different genetic markers, URP-PCR and ISSR and compare the discriminative power of these markers in elucidating the degree of heterogeneity in the population of M. *phaseolina*.

Materials and Methods

Fungal material

Sixty *M. phaseolina* isolates used in this study were obtained in a previous comprehensive sampling from different geographical regions of Iran including Khuzestan, Isfahan, Kerman, Golestan, Hormozgan, Mazandaran, Bushehr, Fars, Yazd and Khorasan-jonoobi (South Khorasan) provinces (Table 1). The isolates were confirmed using species-specific primers (MpKFI/MpKRI) previously described by Babu *et al.* (2007).

DNA extraction

Samples were grown in 250ml glass bottles containing 50ml of Potato Dextrose Broth (PDB) for 5 days at 30 °C in the dark. Mycelia from PDB were harvested by filtering through whatman NO.1 filter paper then dried and deep frozen. Fungal DNA was extracted according to Lee and Taylor, (1990) as modified by Safaie *et al.* (2005). The Quantity and quality were determined spectrophotometrically at 260nm and electrophoretically, respectively and were stored at -20 °C for further use.

| Table | 1 | Name | and | source | of | Macrophomina |
|--------|------|----------|----------|------------|--------|--------------|
| phaseo | lina | isolates | s utiliz | zed in thi | is stı | ıdy. |

| Isolates | Origin | Isolates | Origin |
|----------|-----------|----------|------------------|
| B1 | Bushehr | H10 | Hormozgan |
| B4 | Bushehr | H11 | Hormozgan |
| B5 | Bushehr | H12 | Hormozgan |
| B6 | Bushehr | K1 | Kerman |
| B7 | Bushehr | K2 | Kerman |
| B8 | Bushehr | K3 | Kerman |
| B9 | Bushehr | KH6 | Khuzestan |
| B10 | Bushehr | KH15 | Khuzestan |
| E1 | Isfahan | KH17 | Khuzestan |
| E2 | Isfahan | KH27 | Khuzestan |
| F5 | Fars | KH30 | Khuzestan |
| F7 | Fars | KH32 | Khuzestan |
| F12 | Fars | KH33 | Khuzestan |
| F15 | Fars | KH35 | Khuzestan |
| F23 | Fars | KH36 | Khuzestan |
| F27 | Fars | KH37 | Khuzestan |
| F31 | Fars | KJ1 | Khorasan-jonoobi |
| F33 | Fars | KJ4 | Khorasan-jonoobi |
| F38 | Fars | KJ7 | Khorasan-jonoobi |
| F41 | Fars | KJ8 | Khorasan-jonoobi |
| F43 | Fars | M2 | Mazandaran |
| F45 | Fars | M3 | Mazandaran |
| G1 | Golestan | M9 | Mazandaran |
| G2 | Golestan | M13 | Mazandaran |
| G3 | Golestan | M16 | Mazandaran |
| G4 | Golestan | M19 | Mazandaran |
| H3 | Hormozgan | M20 | Mazandaran |
| H7 | Hormozgan | Y1 | Yazd |
| H8 | Hormozgan | Y2 | Yazd |
| H9 | Hormozgan | Y3 | Yazd |

Primers and polymerase chain reaction

The sequences of five ISSR primers and 12 URP-PCR primers used for the molecular analysis are given in Table 2 and were synthesized by CinnaGen, Iran. The PCR were carried out in a thermocycler (Mastercycler® gradient, Eppendorf, Hamburg, Germany). For both methods final volume of 20µl contained 20 ng of template DNA, 1µM primer, 200mM each of four deoxyribonucleotide triphosphate (dNTP_s), 1.5 units of taq polymerase (CinnaGen, Iran), 2mM MgCl₂ and 2.5µl PCR Buffer (100mM Tris-HCl, 500mM KCl (pH = 8.4)). The PCR reactions were performed for both methods under following conditions: initial denaturation at 94 °C for 4min; 35 cycles of 1min denaturation at 94 °C; 1min annealing at a temperature specific to the primer and extension primer at 70 °C for 2min and 7min at 72 °C for the final product extension. The amplification product were separated by electrophoresis on agarose (1.4%) gel in $0.5 \times$ TAE Buffer run for 2h at 85v and stained with ethidium bromide. A 1Kb DNA ladder (Fermentase, Lithuania) was used as standard molecular weight marker. All PCR assays were repeated at least twice to confirm the reproducibility of the results.

Data analysis

Amplified fragments were considered as a binary character for the present (scored 1) and absent (Scored 0). Similarities based on their fingerprinting groups were estimated by Jaccard's (J) coefficient. Cluster analysis was carried out with NTSYS-pc software, UPGMA (Rholf, 2000). Polymorphism algorithm information content (PIC) was calculated as: PIC = 1 - [(p) 2 + (q)2], where p is the frequency of allele band present and q is frequency of allele band absent (Lynch and Walsh, 1998). Effective multiplex ratio (EMR) and marker index (MI) were calculated according to Powell et al. (1996) to determine the utility of each of the markers in phaseolina Percentage system. М. of Polymorphic Loci (PPL), Shannon's Information Index and Nei's gene diversity were calculated using 'POPGENE' software package ver. 1.31 (Yeh et al., 1999). The total genetic diversity (H_T) , genetic diversity within each population (H_s) coefficient of gene differentiation (G_{st}) were also determined to characterize the gene diversity and the distribution of the variation using 'POPGENE' program. The gene flow estimates (Nm) among these populations were computed as Nm ¹/₄ (1/Gst _ 1)/4 (Nei 1973; Slatkin and Barton 1989). An analysis of molecular variance (AMOVA) was performed to calculate the hierarchical apportionment of variation using GenAlEx version 6.501 (Peakall and Smouse, 2006).

| Table | 2 | Primer | names | and | sequences, | annealing | temperature, | number | of | generated | bands, | number | of : |
|---------|------|----------|----------|-------|-----------------------|-------------|--------------|----------|-----|-----------|--------|----------|------|
| polym | orpl | hic band | ls and d | egree | of polymor | phism of ar | nplified DNA | for each | pri | mer using | URP-PC | R and IS | SSR |
| analysi | is o | f Macro | phomina | a pha | s <i>eolina</i> isola | ites. | | | | | | | |

| Marker | Primer | Sequence | Annealing | No. bands | No. bands | Polymorphism |
|---------|---------|-----------------------------|------------------|-----------|-------------|--------------|
| | | - | temperature (°C) | generated | polymorphic | (%) |
| ISSR | ISSR10 | 5'-CACCACCACCACCAC-3' | 59 | 23 | 19 | 82.20 |
| | ISSR02 | 5'-ACTG ACTGACTGACTG-3' | 48 | 23 | 19 | 82.20 |
| | ISSR09 | 5'-CCACCACCACCA-3' | 55 | 24 | 22 | 83.30 |
| | PcMs | 5'-GTCGTCGTCGTCGTCGTCGTC-3' | 55 | 18 | 10 | 55.50 |
| | P4 | 5'-ATGATGATGATGATGATG-3' | 51 | 17 | 15 | 88.23 |
| URP-PCR | URP-2F | 5'-GTGTGCGATCAGTTGCTGGG-3' | 47 | 13 | 10 | 76.92 |
| | URP-30F | 5'-GGACAAGAAGAGGATGTGGA-3' | 45 | 18 | 11 | 61.11 |
| | URP-2R | 5'-CCCAGCAACTGATCGCACAC-3' | 45 | 17 | 12 | 70.58 |
| | URP-4R | 5'-AGGACTCGATAACAGGCTCC-3' | 46 | 17 | 11 | 64.70 |
| | URP-6R | 5'-GGCAAGCTGGTGGGAGGTAC-3' | 45 | 17 | 9 | 52.94 |
| | URP-13R | 5'-TACATCGCAAGTGACACAGG-3' | 46 | 18 | 14 | 77.77 |
| | URP-1F | 5'-ATCCAAGGTCCGAGACAACC-3' | 47 | 22 | 9 | 69.23 |
| | URP-17R | 5'-AATGTGGGCAAGCTGGTGGT-3' | 55 | 13 | 8 | 61.53 |
| | URP-25F | 5'-GATGTGTTCTTGGAGCCTGT-3' | - | - | - | - |
| | URP-9F | 5'-ATGTGTGCGATCAGTTGCTG-3' | - | - | - | - |
| | URP-32F | 5'-TACACGTCTCGATCTACAGG-3' | - | - | - | - |
| | URP-38F | 5'-AAGAGGCATTCTACCACCAC-3' | - | - | - | - |
| | URP-1F | 5'-ATCCAAGGTCCGAGACAACC-3' | - | - | - | - |

Results

Identification of M. phaseolina isolates using specific primers

In the present study, sixty isolates were identified at the molecular level using species specific primers (MpFI/MoKRI) as designed by Babu *et al.* (2007). A fragment 350-bp was amplified by the primer pair in tested isolates (Fig. 1). The result showed that the isolates belong to the species *M. phaseolina*. The DNA from *Sclerotinia sclerotiorum*, *Fusarium graminearum*, *Alternaria alternata*, *Trichoderma viride* and *Cytospora* sp. were used as negative controls. No fragment was detected in the negative control.

ISSR analysis

Five ISSR primers used to assess genetic diversity of 60 *M. phaseolina* isolates yielded reproducible and polymorphic amplification patterns. A total of 105 bands were produced of which 85 (80.95%) bands were polymorphic. The number of ISSR bands varied from 24 (for the ISSR10) to 17 (for the P4 primers), ranging

in size from 250 to 3000 bp. Genetic similarity coefficient varied from 0.51 to 0.91.

The dendrogram obtained by UPGMA analysis indicated that the isolates were grouped into five clusters at 70% similarity coefficient level (Fig. 2). Cluster I was comprised of two isolates from Bushehr and Yazd. Cluster II contained 54 isolates, consisting of seven isolates from Bushehr, 11 isolates from Fars, nine isolates from Khuzestan, two isolates from Golestan, two isolates Yazd, all from Khorasan-jonoobi, Kerman. isolates Mazandaran, Hormozgan and Isfahan provinces. Clusters III, IV and V contained one, two and one isolates from Khuzestan, Golestan and Fars provinces, respectively. Fifty four isolates grouped in one cluster indicating there is high similarity among isolates from different regions.

The first three most informative principal components of ISSR data explained 80.17% of the total genetic variation observed among isolates. The result of three-dimensional plot of principal component analysis confirmed the cluster analysis results and it indicated that F12 isolate from Fars is the most genetically distant from the other isolates (Fig. 3).



Figure 1 Template DNA was amplified using species-specific primers (MpKFI/MpKRI), M: 1 kb molecular ladder; B7, F45, G4, KH38 and M9 *M. phaseolina* isolates from Iran. R1: *Sclerotinia sclerotiorum*, R2: *Fusarium graminearum*, R3: *Alternaria alternata*, R4 *Trichoderma viride* and R5: *Cytospora ehrenb* are negative controls; NC: Negative control.



Figure 2 Dendrogram derived (constructed) from 60 isolates *Macrophomina phaseolina* by UPGMA clustering based on ISSR.



Figure 3 Genetic relationships of 60 Macrophomina phaseolina isolates based on principal component analysis.

UPR-PCR analysis

Out of 12 URP-PCR primers tested for amplification, 8 primers generated reproducible and well-defined fragments (Fig. 4). A total of 135 fragments were generated, 84 (62.22%) of them were polymorphic. Amplified fragments were from 200bp to 2500bp. The number of polymorphic fragments ranged from 8 to 15, with an average of 10.5 fragments. Genetic similarity coefficient was varied from 0.61 to 0.93. Based on URP-PCR analysis, the isolates were grouped into six clusters by UPGMA at the 77% similarity coefficient level (Fig. 5). Cluster I contained 53 isolates, consisting of 2 sub-clusters; the first sub-cluster contained six isolates from Bushehr, seven isolates from Fars, from Khorasan-jonoobi, three six from Hormozgan, two from Mazandaran, two from Khuzestan, one from Golestan and all isolates from both Isfahan and Yazd provinces. The second sub-cluster included six isolates from four from Fars, four Khuzestan, from Mazandaran, all isolates from Kerman and a single isolate from Bushehr, Khorasan-jonoobi, Hormozgan and Golestan provinces. Cluster III consisted of only two isolates from Fars and Mazandaran and cluster IV contained two isolates from Khuzestan. Clusters II, V and VI are single isolate, namely G1, F45 and M13 which were collected from Golestan, Fars and Mazandaran, respectively.

The first three principal components of URP-PCR data explained 83.34% of the total variation observed among isolates. The grouping of isolates by three first principal components was in agreement with results in clustering. It indicated that G4 isolate from eastern Golestan is the most genetically distant isolate from the other isolates.

Combination of ISSR and URP-PCR analysis The UPGMA dendrogram obtained from combination of URP-PCR and ISSR fingerprints indicated that *M. phaseolina* isolates could be differentiated into eight clusters at 78%

similarity coefficient level (Fig. 6).



Figure 4 URP-PCR profile generated using URP-13R; B = Bushehr; E = Isfahan; F = Fars; G = Golestan; H = Hormozgan; K = Kerman; KH = Khuzestan; KJ = Khorasan-jonoobi; M = Mazandaran; Y = Yazd; M = molecular weight marker GeneRulerTM 1 kb DNA ladder (Fermentase, Lithuania); NC = Negative control.



Figure 5 Dendrogram derived (constructed) from 60 isolates *Macrophomina phaseolina* by UPGMA clustering based on URP-PCR.

Cluster I consisted of two isolates from Bushehr, two from Hormozgan and one from Yazd. Cluster II had two sub-cluster, the first sub-cluster was comprised of three isolates from Bushehr, four from Fars, seven from Khuzestan, four from Mazandaran, all isolates from Kerman, two from Golestan and single isolates from Hormozgan and Khorasanjonoobi provinces and the second sub-cluster was composed of two isolates from Bushehr, three from Khorasan-jonoobi, two from Yazd, six from Fars, four from Hormozgan, two from Mazandaran, all isolates from Isfahan and single one each from Khuzestan and Golestan provinces. Cluster III contained two isolates from Fars and Mazandaran provinces. Clusters IV, V, VI, VII and VIII that each contained one isolate from Khuzestan, Khuzestan, Bushehr, Fars and Golestan, respectively. Genetic similarity coefficient was varied from 0.61 to 0.90.

The first four principal components from combination of ISSR and URP-PCR data explained 82.74% of the total variation observed among isolates. The result of threedimensional plot of principle component analysis confirmed the cluster analysis results and it indicated that G4 isolate from Golestan is the most genetically distant from the others.



Figure 6 Dendrogram derived (constructed) from 60 isolates *Macrophomina phaseolina* by UPGMA clustering based on URP-PCR and ISSR.

Comparison of ISSR and URP-PCR

EMR, MI and the product of the polymorphism information content (PIC) were used as criterion for the assessment the usefulness of each marker. For ISSR marker, MI, EMR and PIC were 3.28, 13.94 and 0.246, respectively, while for URP-PCR these indices were 1.56, 5.54 and 0.33, respectively.

The cophenetic correlation (r) value for similarity matrices of URP-PCR and ISSR were 0.76 and 0.86, respectively, demonstrated a good fit of the cluster analysis to the similarity data. The cophenetic correlation with combination of URP-PCR and ISSR fingerprints was r = 0.83, which according to Mantel, (1967) is significant.

The Mantel test showed that the correlation coefficient between the two similarity matrices

obtained by ISSR and URP-PCR was significant (r = 0.44, P = 0.002), suggesting existence of positive correlation between two molecular markers.

Genetic differentiation between populations

The higher values, Shannon's Information Index (0.258 \pm 0.269) and PPL (53.33%) were in Fars populations and Nei's gene diversity (0.17 \pm 0.2096) in Golestan populations. Percentage of Polymorphic Loci (PPL) (9.58%,), Shannon's Information Index (0.061 \pm 0.1878) and Nei's gene diversity (0.042 \pm 0.1311) had the lowest values in the populations of Kerman (Table 3).

The AMOVA conducted using URP-PCR + ISSR data showed that from the total genetic variability, 10% (p < 0.01) could be

attributed by the differences between populations while 90% (p < 0.01) could be attributed by the differences within populations. The within and among population variability or the URP-PCR and ISSR data were 88-12% and 91-9%, respectively (Table 4).

Table 3 Genetic analysis among MacrophominaphaseolinapopulationsobtainedfromURP-PCRandISSR data.

| N | Н | Ι | р |
|----|---|--|--|
| 8 | 0.157 ± 0.191 | 0.235 ± 0.278 | 45.00 |
| 2 | 0.068 ± 0.172 | 0.095 ± 0.293 | 13.75 |
| 12 | 0.168 ± 0.187 | 0.258 ± 0.269 | 53.33 |
| 4 | 0.170 ± 0.209 | 0.248 ± 0.302 | 40.83 |
| 7 | 0.131 ± 0.188 | 0.195 ± 0.274 | 35.42 |
| 3 | 0.042 ± 0.131 | 0.061 ± 0.187 | 9.58 |
| 10 | 0.164 ± 0.188 | 0.250 ± 0.273 | 49.58 |
| 4 | 0.112 ± 0.185 | 0.164 ± 0.269 | 27.50 |
| 7 | 0.142 ± 0.188 | 0.213 ± 0.275 | 39.58 |
| 3 | 0.085 ± 0.175 | 0.122 ± 0.251 | 19.17 |
| | N 8 2 12 4 7 3 10 4 7 3 | $\begin{array}{c cccc} N & H \\ \hline 8 & 0.157 \pm 0.191 \\ 2 & 0.068 \pm 0.172 \\ 12 & 0.168 \pm 0.187 \\ 4 & 0.170 \pm 0.209 \\ 7 & 0.131 \pm 0.188 \\ 3 & 0.042 \pm 0.131 \\ 10 & 0.164 \pm 0.188 \\ 4 & 0.112 \pm 0.185 \\ 7 & 0.142 \pm 0.188 \\ 3 & 0.085 \pm 0.175 \end{array}$ | NHI8 0.157 ± 0.191 0.235 ± 0.278 2 0.068 ± 0.172 0.095 ± 0.293 12 0.168 ± 0.187 0.258 ± 0.269 4 0.170 ± 0.209 0.248 ± 0.302 7 0.131 ± 0.188 0.195 ± 0.274 3 0.042 ± 0.131 0.061 ± 0.187 10 0.164 ± 0.188 0.250 ± 0.273 4 0.112 ± 0.185 0.164 ± 0.269 7 0.142 ± 0.188 0.213 ± 0.275 3 0.085 ± 0.175 0.122 ± 0.251 |

Population genetic parameters (PGP).

Samples size (N).

Nei's (1973) gene diversity (H).

Shannon Information Index (I).

% of polymorphic loci (P).

Table 4 Nested analysis of molecular variances(AMOVA) for Macrophomina phaseolina isolatesfrom 10 regions of Iran.

| Source | df | Sum of squares | Mean squares | Est. Var. | % | | | | |
|--------------------|----|----------------|--------------|--------------|-----|--|--|--|--|
| ISSR | | | | | | | | | |
| Among Pops | 9 | 133.655 | 14.851 | 0.912 | 9 | | | | |
| Within Pops | 50 | 477.495 | 9.550 | 9.550 | 91 | | | | |
| Total | 59 | 611.150 | | 10.461 | 100 | | | | |
| URP-PCR | | | | | | | | | |
| Among Pops | 9 | 172.770 | 19.197 | 1.439 | 12 | | | | |
| Within Pops | 50 | 541.430 | 10.829 | 10.829 | 88 | | | | |
| Total | 59 | 714.200 | | 12.268 | 100 | | | | |
| ISSR + URP- PCR | | | | | | | | | |
| Among Pops | 9 | 306.425 | 34.047 | 2.351 | 10 | | | | |
| Within Pops | 50 | 1018.925 | 20.379 | 20.379 | 90 | | | | |
| Total | 59 | 1325.350 | | 22.729 | 100 | | | | |

The total gene diversity (H_t) and mean diversity within each population (H_s) belonging to URP-PCR + ISSR data were obtained for the populations which were 0.181 and 0.124, respectively, and the mean coefficient of gene differentiation (G_{st}) among populations was 0.313, indicating 31% of differentiation between populations (Table 5). Similarly, almost relevant results were obtained for relative differentiation (G_{st}) , total Heterozygosity (H_t) , intra-population heterozygosity (H_s) indices over populations for either individual datasets of each marker analyses.

Table 5 Genetic structure estimated by Nei's geneticparametersamongMacrophominaphaseolinapopulationsobtained from URP-PCR + ISSR data.

| Population genetic parameters | Obtained values |
|---|-------------------|
| Intra-population (H _s) | 0.124 ± 0.016 |
| Total heterozygosity (H _T) | 0.181 ± 0.034 |
| Relative differentiation (G _{ST}) | 0.313 |
| Estimate gene flow (N_m) | 1.09 |

The amount of gene flow (N_m) from URP-PCR + ISSR data between populations was 1.09, where N_m is the average number of migrants among the population. According to Wright, (1951), the $N_m > 1$ shows the existence of little differentiation among populations. No specific fragments were found in either population. This result shows that ISSR and URP-PCR divergence among populations was mainly ascribed to difference of the DNA-bands frequency instead of allele fixation.

The values of Nei's genetic distance ranged from 0.027 to 0.123 for 10 populations. The lowest value of genetic distance was obtained between Fars and Hormozgan populations and showed that the populations from these two regions were most similar (Table 6).

The relationship among populations was estimated based on Nei's genetic distance. The UPGMA dendrogram revealed that clusters were not related to the geographic distance between populations. Cluster analysis (UPGMA) demonstrated that the populations could be differentiated into six clusters. Cluster I consisted of five populations from Bushehr, Fars, Hormozgan, Khuzestan and Mazandaran, clusters II, III, IV, V and VI were comprised of Khorasan-jonoobi, Golestan, Yazd, Isfahan, and Kerman provinces, respectively (Fig. 7). In cluster I populations of Hormozgan and Fars showed 100% similarity coefficient. Geographically, both Fars and Hormozgan populations have grown at low latitude and almost in similar climates, which result in high genetic similarity between them.

| Table 6 Nei's genetic distance measured amore | ig ten p | populations of | f Macro | phomina | phaseolina |
|---|----------|----------------|---------|---------|------------|
|---|----------|----------------|---------|---------|------------|

| Population | В | Е | F | G | Н | K | KH | KJ | М | Y |
|------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| В | 0.000 | | | | | | | | | |
| Е | 0.084 | 0.000 | | | | | | | | |
| F | 0.028 | 0.064 | 0.000 | | | | | | | |
| G | 0.067 | 0.116 | 0.066 | 0.000 | | | | | | |
| Н | 0.038 | 0.070 | 0.027 | 0.062 | 0.000 | | | | | |
| K | 0.095 | 0.184 | 0.087 | 0.120 | 0.094 | 0.000 | | | | |
| KH | 0.046 | 0.106 | 0.042 | 0.063 | 0.045 | 0.064 | 0.000 | | | |
| KJ | 0.050 | 0.094 | 0.055 | 0.092 | 0.050 | 0.109 | 0.051 | 0.000 | | |
| М | 0.048 | 0.093 | 0.045 | 0.071 | 0.050 | 0.075 | 0.039 | 0.056 | 0.000 | |
| Y | 0.072 | 0.106 | 0.077 | 0.114 | 0.067 | 0.123 | 0.081 | 0.063 | 0.065 | 0.000 |



Figure 7 UPGMA cluster using Nei's genetic distance among ten populations of *Macrophomina phaseolina*. B = Bushehr, E = Isfahan, F = Fars, G = Golestan, H = Hormozgan, K = Kerman, Kh = Khuzestan, KJ = Khorasan-jonoobi, M = Mazandaran, Y = Yazd.

Discussion

Macrophomina phaseolina the causal agent of charcoal rot of sesame, is an important fungal pathogen causing economically important crop losses in oil seeds in Iran. Because of the high level of heterogeneity, the use of resistant cultivars was proposed as the most effective and practical method to control charcoal rot of sesame (Reyes-Franco et al., 2006). Understanding genetic diversity within and between populations is essential for the improvement of disease management systems and sesame breeding program against charcoal rot. In this study, the distribution of genetic diversity among 60 M. phaseolina isolates from different sesame growing regions in Iran was investigated and URP-PCR and ISSR genomic fingerprinting revealed different levels of heterogeneity in populations of this fungus.

The dendrogram obtained with of **URP-PCR** combination and ISSR fingerprints showed that isolates of M. phaseolina were separated into eight groups. The significant correlation (r = 0.83)between cophenetic matrices value generated ISSR URP-PCR from and datasets demonstrated that both molecular markers tools for are powerful analyzing the intraspecific variability among М. phaseolina isolates.

The existence of genetic diversity among isolates of *M. phaseolina* in Iran could be due to variations in the climatic conditions and cropping patterns in different parts of the country, also transportation of the pathogen (Aghakhani *et al.*, 2009; Baird *et al.* 2010)

Cluster analysis indicated that *M. phaseolina* isolates obtained from soybean, cotton, and chickpea hosts clearly differentiated into specific groups (Jana *et al.*, 2005b).

In the present study URP-PCR marker revealed a high genetic diversity within and between groups.

Different molecular approaches have been used in the analysis of genetic variation among *M. phaseolina* from various plant hosts and geographic locations.

AFLP assessment of the polymorphism of M. phaseolina isolates from sesame in China has shown a high level variations of 73.9% among the isolates (Linhai et al., 2011). ISSR evaluation of *M. phaseolina* populations from soybean and cotton grown in India and USA has shown highly significant genetic heterogeneity among and within populations and isolates could be grouped into three major clusters according to their hosts and geographical region (Jana et al., 2005a). Genetic variation of M. phaseolina isolates in India using RAPD marker has grouped the isolates into six categories at 40% genetic similarity (Aghakhani et al., 2009). Using AFLP analysis, Reyes-Franco et al. (2006) reported that genetic variation of isolates was not strongly correlated to geographic origin for Mexico and other countries.

Mahdizadeh *et al.* (2011) reported that isolates from 24 host plant species did not clearly differentiate to the specific group according to the host or geographical origins, however, usually the isolates from the same host or the same geographic origin tend to group nearer. The results of this study correspond with above findings that there exists a high genotypic diversity level among M. *phaseolina* isolates.

In populations of *M. phaseolina*, diversity parasexual could be attributed to recombination between nuclear genes or fusion of cells favoring vegetative heterokaryons (Carlile 1986). Factors such as immigration, climatic conditions, different patterns of culture, use of various different host genotypes, high selection pressure and breeding system might explain high levels of genetic diversity of the fungus (Aghakhani et al. 2009; Almeida et al. 2008; Purkayastha et al. 2008). Miao et al. (1991) reported that unstable B chromosome may be one of the mechanisms for generating variation in fungi.

According to our results, no clear relationship was found between genetic diversity and the geographical origin of the isolates using URP-PCR and ISSR molecular markers. Previous studies showed that

application of AFLP (Mayek-Perez et al., 2001), ISSR (Jana et al., 2005a) markers could beneficially be used to group isolates based on their geographical areas but there are some reports that they could not clearly be grouped according to geographical areas (Aghakhani et al. 2009; Baird et al. 2010; Khan et al. 2017; Linhai et al. 2011; Mahdizadeh et al. 2011). Sanchez et al. (2017) using SSR markers reported a clear tendency in isolates to cluster by origin, and making evident a genetic distance between Chilean and Spanish strawberry isolates.

The results of the present findings revealed considerable genetic similarities among the isolates from different geographic areas. Hamer *et al.* (1989) provided evidence at the DNA sequence level for pathogen genotype selection by a host species.

We suggest that these isolates may evolve from the same ancestral population. This result is in agreement with the results obtained by Almeida et al. (2003) and Linhai et al. (2011) on isolates from the hosts as Sorghum, Soybean, Chickpea, Corn and Sesame. In addition, the similarity among isolates of remote origin confirmed free flux genetic genetic information and exchange in the genus, reducing the probability of delimited groups formation (Mihail and Taylor, 1995).

The results showed that the Fars and Golestan populations have the highest heterozygosity among the ten populations. This might be caused by diversity of sesame varieties in the two regions.

Comparison of two molecular marker systems used in this study demonstrated that the average marker index was between 3.28 (ISSR) to 1.56 (URP-PCR). Compared with URP-PCR, ISSR markers are much more informative and powerful tools for assessing variability in *M. phaseolina*.

The low level of observed genetic differentiation could be related to level of gene flow among populations. In addition, the observation of low levels of differentiation in populations could be explained by asexual propagules and polyphagy of this fungus (Salahlou *et al.*, 2018).

In this study, both genetic differentiation and AMOVA indicated that most of the genetic diversity is within populations. One reason for high genetic diversity within populations might be parasexual recombination among isolates (Carlile, 1986). The high level of genetic variation within a fungal population could be due to the uniformity of its substrate and previous environmental experience (Ijaz *et al.*, 2013). Kendrick, (1992) reported that degree of sexual reproduction within a population is closely related to genetic diversity.

The level of gene flow (N_m) was measured at 1.09, demonstrating a very high migration rate between populations. This gene flow might be due to movement of M. phaseolina through infected seeds and soil with planting materials by means of vegetative stage and sclerotia and may be occurring through natural dispersal mechanisms involving ascospores (Kohn, 1995). Gene flow induces replacement of new alleles or genotypes in new areas via introduction, reproduction and survival of the introduced organism. The Gst and Nm indices indicated high gene flow and low genetic differentiation among populations. Gene Genetic flow and differentiation amount generated from both molecular marker systems separately were similar to those generated with URP-PCR and ISSR data. Our survey suggests that frequent gene flow and recombination between populations of M. phaseolina have significantly impressed the evolution and development of this fungus in Iran. Gordon and Martyn, (1997) reported that high level of pathogenic and genetic diversity in fungus is also asexual observed with accumulation of genetic mutation over time.

Investigation of genetic variability induced by parasexual reproduction of fungi has two crucial factors including activity of transposons and a high mutation rate of microsatellites (Purkayastha *et al.*, 2008).

In conclusion, we discovered a high level of intraspecific variation among *M. phaseolina* pathogens using URP-PCR and ISSR. Both methods are shown to be useful for population

analysis of *M. phaseolina* isolates. Comparison of the two molecular marker systems used in this study showed that ISSR markers were highly informative to study the polymorphisms among *M. phaseolina* populations causing charcoal rot of Sesame. However, to evaluate genetic diversity among races of *M. phaseolina* the utilization of these molecular marker systems, especially ISSR markers should be considered as one of the priority aims of studies in future. Our study provided primary information on the population genetics of *M. phaseolina* from different geographical regions in Iran, which is essential for sesame breeding program against the charcoal rot.

Reference

- Aboshosha, S. S., Attaalla, S. I., El-Korany, A. E. and El-argawy, E. 2007. Characterization of Macrophomina phaseolina isolates affecting sunflower growth in El-Behera governorate. International Journal of Agriculture and Biology, 9: 807-815.
- Aghakhani, M. and Dubey, S. C. 2009. Determination of genetic diversity among Indian isolates of *Rhizoctonia bataticola* causing dry root rot of chickpea. Antonie van Leeuwenhoek, 96: 607-619.
- Ahmad, I. and Burney, K. 1990. Macrophomina phaseolina infection and charcoal rot development in sunflower under field conditions. In: Ooi P, Lim G. S. and Teng, P. Proceedings S. (Eds.), of the Third International Conference in Plant Protection in the Tropics; 1990 March 20-23; Malaysia. Kuala Lumpur: Genting Highlands, pp: 58-61.
- Almeida, M. R., Abdelnoor, R. V., Arias, C. A.
 A., Carvalho, V. P., Martin, S. R. R., Benato, L. C., Pinto, M. C. and Carvalho, C.
 G. P. 2003. Genotypic diversity among brazilian isolates of *Macrophomina phaseolina* revealed by RAPD. Fitopatologia Brasileira, 28: 279-285.
- Almeida, M. R., Sosa-Gomez, D. R., Binneck, E., Marin, S. R. R., Zucchi, M.
 I., Abdelnoor, R. V. and Souto, E. R.
 2008. Effect of crop rotation on specialization and genetic diversity of

Macrophomina phaseolina. Tropical Plant Pathology, 33: 257-264.

- Babu, B. K., Saxena, A. K., Srivastava, A. K. and Arora, D. K. 2007. Identification and detection of *Macrophomina phaseolina* by using species-specific oligonucleotide primers and probe. Mycologia, 99: 797-803.
- Baird, R. E., Wadl, P. A., Allen, T., McNeill, D., Wang, X. W., Moulton, J. K., Rinehart, T. A., Abbas, H. K., Shier, T. and Trigiano, R. N. 2010. Variability of United States isolates of *Macrophomina phaseolina* based on simple sequence repeats and cross genus transferability to related genera within Botryosphaeriaceae. Mycopathologia, 170: 169-180.
- Bornet, B. and Branchard, M. 2001. Nonanchored Inter Simple Sequence Repeat (ISSR) markers: reproducible and specific tools for genome fingerprinting. Plant Molecular Biology Reporter. 19: 209-215.
- Broglie, K. E., Gaynor, J. J. and Broglie, R. M. 1986. Ethylene-regulated gene expression: molecular cloning of the genes encoding an endochitinase from *Phaseolus vulgaris*. Proceedings of the National Academy of Science of the United States of America. 1986 Sept 9, USA.
- Carlile, M. J. 1986. Genetic exchange and gene flow: their promotion and prevention. In: Rayner A. D. M. and Moore, D. (Eds.), Evolutionary Biological of the Fungi. Cambridge: Cambridge University Press; pp: 203-214.
- Chakravarthi, B. K. and Naravaneni, R. 2006. SSR Marker Based DNA Fingerprinting and Diversity Study in Rice (Oryza sativa. L.). African Journal of Biotechnology, 5: 684-688.
- Dhingra, O. D. and Sinclair, J. B. 1978. Biology and Pathology of *Macrophomina phaseolina*. Vic.osa (Brasil): Imprensa Universita' ria, Universidade Federal de Vic.osa.
- Ershad, D. 1995. Fungi of Iran. Ministry of Agriculture, Agricultural Research, Education and Extension Organization Publication. No. 10, Tehran, Iran.

Downloaded from jcp.modares.ac.ir on 2024-04-28

- Etebarian, H. R. 2006. Evaluation of Streptomyces strains for Biological Control of Charcoal Stem Rot of Melon Caused by *Macrophomina phaseolina*. Plant Pathology Journal, 5: 83-87.
- Geo, P., Yao, Q., Ding, B., Zeng, H., Zhong, Y., Fu, C. and Jin, X. 2006. Genetic diversity of *Sinojackia dolichocarpa* (Styracaceae), a species endangered and endemic to China, detected by inter-simple sequence repeat (ISSR). Biochemical Systematics and Ecology, 34: 231-239.
- Gordon, T. R. and Martyn. R. D. 1997. The evolutionary biology of Fusarium oxysporum. Annual Review of Phytopathology, 35: 111-128.
- Hamer, J. E., Farrall, L., Orbach, M. J., Valent, B. and Chumley, F. G. 1989. Host speciesspecific conservation of a family of repeated DNA sequences in the genome of a fungal plant pathogen. PNAS USA. 86: 9981-9985.
- Hlinkova, E. and Sykora, M. 1996. Changes in extracellular protein patterns induced by powdery mildew (*Erysiphe graminis* f.sp. *hordei*) infection (abstract). In: Kema, G. H. J., Niks, R. E. and Daamen, R. A. (Eds.), Proceedings of the Ninth European and Mediterranean Cereal Rusts and Powdery Mildew Conference; 1996 Sept 2-6; Wageningen, The Netherlands.
- Ijaz, S. Sadaqat, H. A. and Khan, M. N. 2013. A review of the impact of charcoal rot (*Macrophomina phaseolina*) on sunflower. Journal of Agricultural Science, 151 (2): pp. 222-227.
- Jana, T., Sharma, T. R. and Singh, N. K. 2005a. SSR-based detection of genetic variability in the charcoal root rot pathogen Macrophomina phaseolina. Mycological Research, 109: 81-86.
- Jana, T., Singh, N., Koundal, K. and Sharma, T. 2005b. Genetic differentiation of charcoal rot pathogen, *Macrophomina phaseolina*, into specific groups using URP-PCR. Canadian Journal of Microbiology, 51: 159-164.
- Kang, H. W., Park, D. S., Par, Y. J., You, C. H., Lee, B. M., Eun, M. Y. and Go, S. J. 2002. Fingerprinting of diverse genomes using

PCR with universal rice primers generated from repetitive sequence of Korean weedy rice. Molecules and Cells, 13: 281-287.

- Kendrick, B. 1992. The Fifth Kingdom. 2d ed. Waterloo, Ontario, Canada: Mycologue Publication, Focus Information Group Inc.
- Khan, A. N., Shair, F., Malik, K., Hayat, Z., Khan, M. A., Hafeez, F. Y. and Hassan, M. N. 2017. Molecular Identification and Genetic Characterization of *Macrophomina phaseolina* Strains Causing Pathogenicity on Sunflower and Chickpea. Frontiers in Microbiology, 8: 1309. doi: 10.3389/fmicb. 2017.01309.
- Kohn, L. M. 1995. The clonal dynamic in wild and agricultural plant pathogen populations. Canadian Journal of Botany, 73: S1231-S140.
- Lee, S. B. and Taylor, J. W. 1990. Isolation of DNA from fungal mycelia and single spores.
 In: Innis, M. A., Gelfand, D. H., Sinsky, J. J. and White, T. J. (Eds.), PCR Protocols: A Guide to Methods and Applications.
 London: Academic Press. pp: 282-287.
- Linhai, W., Yanxin, Z., Donghua, L., Junbin, H., Wenliang, W., LvHaixia, L. and Xiurong Z. 2011. Variations in the isolates of Macrophomina phaseolina from sesame in China based on amplified fragment length polymorphism (AFLP) and pathogenicity. African Journal of Microbiology Research, 5: 5584-5590.
- Lynch, M. A. and Walsh, J. B. 1998. Genetics and Analysis of Quantitative Traits. Sinauer Associates Inc, Sunderland, MA.
- Mahdizadeh, V., Safaie, N. and Mohamadi-Goltapeh, E. 2011. Diversity of *Macrophomina phaseolina* Based on Morphological and Genotypic Characteristics in Iran. Plant Pathology Journal, 27: 128-137.
- Maleki, M., Shojaeiyan, A. and Monfared, S. R. 2018. Population structure, morphological and genetic diversity within and among melon (*Cucumis melo* L.) landraces in Iran. Journal of Genetic Engineering and Biotechnology, 16 (2): 599-606.
- Mantel, N. 1967. The detection of disease clustering and a generalized regression approach. Cancer Research, 27: 209-220.

Downloaded from jcp.modares.ac.ir on 2024-04-28

- Mayek-Perez, N., Lpez-Castaeda, C., Gonzglez-Chavira, M., Garcia-Espinosa, R., Acosta-Gallegos, J., de la Vega, O. M. and Simpson, J. 2001. Variability of Mexican isolates of Macrophomina phaseolina based on pathogenesis and AFLP genotype. Physiological Molecular and Plant Pathology, 59: 257-264.
- Menzies, J. G., Bakkeren, G., Matheson, F., Procunier, J. D. and Woods, S. 2003. Use of inter-simple sequence repeats and amplified fragment length polymorphisms to analyze genetic relationships among small graininfecting species of Ustilago. Phytopathology, 93: 167-175.
- Miao, V. P, Covert, S. F. and VanEtten, H. D. 1991. A fungal gene for antibiotic resistance on a dispensable ("B") chromosome. Science, 254: 1773-1776.
- Mihail, J. D. and Taylor, S. J. 1995. Interpreting variability among isolates for *Macrophomina phaseolina* in pathogenicity, pycnidium production and chlorate utilization. Canadian Journal of Botany, 73: 1596-1603.
- Nei, M. 1973. Analysis of gene diversity in subdivided populations. PNAS USA, 70: 3321-3323.
- Peakall, R. and Smouse, P. E. 2006. GENALEX6: genetic analysis in excel. Population genetic software for teaching and research. Molecular Ecology Notes, 6: 288-295.
- Pecina, V., Alvarado, M. J., Alanis, H. W., Almaraz, R. T. and Vandemark, G. J. 2000 Detection of double-stranded RNA in *Macrophomina phaseolina*. Mycologia, 92: 900-907.
- Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J. and Tingey, S. 1996. The comparison of RFLP, RAPD, AFLP and SSR (microsattlite) markers for germplasm analysis. Molecular Breeding, 3: 225-238.
- Purkayastha, S., Kaur, B., Arora, P., Bisyer, I., Dilbaghi, N. and Chaudhury, A. 2008. Molecular genotyping of *Macrophomina phaseolina* isolates: comparison of microsatellite primed PCR and repetitive element sequence-based PCR. Journal of Phytopathology, 156: 372-381.

- Radwan, N. A. M. 2000. Implication of certain physical and chemical treatments to improve diseases resistance of barley to powdery mildew. Ph. D. Thesis, Cairo University.
- Reyes-Franco, M. C., Hernandez-Delgado, S., Beas-Fernandez, R., Medina-Fernandez, M., Simpson, J. and Mayek-Perez, N. 2006. Pathogenic and genetic variability within *Macrophomina phaseolina* from Mexico and other countries. Journal of Phytopathology, 154: 447-453.
- Rholf, F. J. 2000. NTSYS-pc Numerical taxonomy and multivariate system version 2.1. Exeter publishing, Setauket.
- Safaie, N., Alizadeh, A., Saidi, A., Rahimian, H. and Adam, G. 2005. Molecular characterization and genetic diversity among Iranian populations of *Fusarium graminearum*, the causal agent of wheat head blight. Iran. Journal of Plant Pathology, 41: 171-189.
- Salahlou, R., Safaie, N. and Shams-Bakhsh, M., 2018. Genetic diversity of *Macrophomina phaseolina* populations, the causal agent of sesame charcoal rot using inter-simple sequence repeat markers. Journal of Agricultural Science and Technology, 18 (1): 277-278.
- Sanchez, S., Chamorro, M., Henriquez, J., Grez, J., Díaz, I., Santos, B. D. L. and Gambardella, M. 2017. Genetic and biological characterization of *Macrophomina phaseolina* (Tassi) Goid. causing crown and root rot of strawberry. Chilean journal of agricultural research, 77 (4): 325-331.
- Slatkin, M. and Barton, N. H. 1989. A comparison of three indirect methods for estimating average levels of gene flow. Evolution, 43: 1349-1368.
- Yeh, F. C, Boyle, T. J. B., Ye, Y. Z. and Xiyan, J. M. 1999. POPGENE version 1.32. Microsoft Window-based freeware for population genetics analysis. University of Alberta and Center for International Forestry Research. (Last accessed March 2005).
- Zietkiewicz, E., Rafalski, A. and Labuda, D. 1994. Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. Genomics, 20: 176-183.

استفاده از نشانگرهای ISSR و URP-PCR در بررسی تنوع ژنتیکی جدایههای Macrophomina phaseolina روی کنجد در ایران

رضا صلاحلو'، ناصر صفایی^{۱*} و مسعود شمسبخش^۲

۱- گروه بیوتکنولوژی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران. ۲- گروه بیماریشناسی گیاهی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران. پست الکترونیکی نویسنده مسئول مکاتبه: nsafaie@modares.ac.ir دریافت: ۲۶ مهر ۱۳۹۷؛ پذیرش: ۲۵ خرداد ۱۳۹۸

چکیده: بهمنظور بررسی تنوع Macrophomina phaseolina عامل پوسیدگی ذغالی کنجد، شصت جدایه از ده منطقه جغرافیایی جمعآوری شده، با استفاده از نشانگرهای مولکولی ISSR و URP-PCR مورد تجزیه و تحلیل قرار گرفتند. جدایهها به هشت گروه در سطح تشابه ژنتیکی 78 درصد تقسیم شدند. این نتایج نشان داد که از پنج آغازگر ISSR مورد مطالعه در مجموع ۱۰۵ لوکوس که ۷۷.۱۱ درصد چند شکلی بودند و هشت آغازگر URP-PCR تولید ۱۳۵ لوکوس که ۶۶.۸۴ درصد چند شکلی بودند. این روشها تنوع ژنتیکی قابل توجهی را در بین جدایههای ایران نشان داد اما هیچگونه همبستگی بین تنوع ژنتیکی و مناطق جغرافیایی جدایهها مشاهده نشد. تجزیهوتحلیل واریانس مولکولی نشان داد که تنوع ژنتیکی نسبتاً زیاد، ناشی از تفاوت جدایهها در داخل مناطق جغرافیایی است. نتایج یافتههای حاضر نشان داد که تمایز ژنتیکی کم (G_{ST}) و جریان ژن بالا (M) در میان جمعیتها تأثیر قابل توجهی بر ظهور و سیر تکاملی M. داشته است.

واژگان کلیدی: پوسیدگی ذغالی، Sesamum indicum ، اثر انگشت دی ان آ، ژنتیک جمعیت