Research Article



Larvicidal and antifeedant activity of some indigenous plants of Meghalaya against 4th instar *Helicoverpa armigera* (Hübner) larvae

Dibyendu Paul and Moumita Choudhury*

Department of Environmental Studies, North Eastern Hill University, Shillong- 793022. Meghalaya, India.

Abstract: In the present study, seven indigenous, widely distributed plants of Meghalaya, namely, Pinus kesiya Royle (Pinaceae), Lantana camara Linn. (Verbenaceae), Litsea cubeba Lour. (Lauraceae), Gaultheria fragrantissima Wall. (Ericaceae), Mikania micrantha Kunth.(Asteraceae), Ambrosia artemisiifolia Linn.(Asteraceae) and Eupatorium riparium Regel (Asteraceae) were screened for their larvicidal and antifeedant activity against fourth instar larvae of the cotton bollworm, Helicoverpa armigera (Hübner) under laboratory conditions. The crude extracts of all the seven plants demonstrated a dose dependent increase in bioactivity. However the bioactivity of four plants namely, L. camara, G. fragrantissima, L. cubeba and *P. kesiya* was significantly higher ($p \le 0.05$) than the negative (solvent) control and extracts of A. artemisiifolia, E. riparium and M. micarantha. Methanol extract of L. camara caused highest oral toxicity with larval mortality ranging between 27.77% and 53.33% across the test concentration (0.25%, 0.5% and 1% w/v) while extract of G. fragrantissima demonstrated the highest feeding deterrence with reduction in larval feeding by50.92% and 70.61% at 0.1% and 0.5% respectively. Crude extract of L. cubeba leaves demonstrated high oral toxicity and feeding deterrence while extract of the needles of P. kesiya showed moderate level of oral toxicity as well as feeding deterrence at the highest tested concentration. Phytochemical analysis of the extracts of these four plants, revealed the presence of five different classes of phytocompounds each of which is known to have deleterious effect on insect pests. Thus it may be concluded that four out of the seven plants possess insecticidal property and can be further investigated for the development of a potent botanical insecticide.

Keywords: plant extract, oral toxicity, antifeedant activity, Helicoverpa armigera, Meghalaya

Introduction

Helicoverpa armigera Hübner (Lepidoptera: Noctuidae) is a polyphagous migratory noctuid

which is widespread in Asia, Africa and Oceania (Lammers and Macleod, 2007). It is known to cause serious damage to hundreds of economically important crops all over the world (Setiawati *et al.*, 2000; Fakrudin *et al.*, 2004). In India it is reported to be feeding on 182 plant species across 47 families (Manjunath *et al.*, 1985) and causes an annual loss of about Rs. 2,000 crores (Ignacimuthu and Jayaraj, 2003).

Handling Editor: Saeid Moharramipour

^{*}Corresponding author, e-mail: moumita.choudhury@gmail.com Received: 25 January 2016, Accepted: 12 June 2016 Published online: 28 June 2016

Fifty percent of all insecticides used in India and China are to control H. armigera alone (Lammers and Macleod, 2007) but the continuous and indiscriminate use of insecticides over the years has resulted in the H. armigera developing resistance to certain molecules belonging to different classes of insecticides in various parts of the world (McCaffery, 1998; Chaturvedi, 2007; Yang et al., 2013). Thus alternatives to the synthetic pesticides are being sought.

The search for alternatives to synthetic pesticides has focused the interest of the pest managers on plant derived pest control agents. Plant-based pesticides or botanicals have many advantages: firstly, they have multifarious control mechanisms against pests (Sivagnaname and Kalyanasundaram, 2004) which reduces the possibility of the development of resistance in pests (Liu *et al.*, 2000); secondly, they are target-specific and hence not harmful to humans and beneficial insects; and lastly, they are not persistent in nature and hence environment friendly (Shaalan, 2005).

In the present investigation an attempt has been made to screen seven widely distributed plants of Meghalaya, for their insecticidal activity against fourth instar larvae of H. armigera, which has been reported as a major pest of tomato and chickpea in the state (Thakur et al., 2006). The effect of many different plants and their extracts on H. armigera has been studied by several authors (Pandey et al., 1983; Jotwani and Srivastava, 1984; Hongo and Karel, 1986; Sahayaraj, 1998; Sundararajan and Kumuthakalavalli 2001; Koul et al., 2002; Kathuria and Kaushik, 2005; Ramya et al., 2008; Wambua et al., 2011; Jeyashankar et al., 2012; Arivoli and Tennyson, 2013). While extracts of certain plants like Ocimum basilicum, Gynandropsis gynandra, Acorus calamus, Lantana camara, and Toddalia asiatica demonstrated larvicidal effect on H. armigera (Pandey et al., 1983; Sundararajan and Kumuthakalavalli, 2001), others like neem seed kernel extract were seen to have indirect effects like causing larvalpupal intermediaries and abnormal adults (Jotwani and Srivastava, 1984) and feeding deterrency (Hongo and Karel, 1986). Majority of the plants tested against different larval instars of *H. armigera* have been reported to demonstrate antifeedant properties (Sahayaraj, 1998; Koul *et al.*, 2002; Kathuria and Kaushik, 2005; Ramya *et al.*, 2008; Wambua *et al.*, 2011; Jeyashankar *et al.*, 2012; Arivoli and Tennyson, 2013).

Although extensive research has been conducted on the effect of different plant extracts on H. armigera, there is limited literature available on the efficacy of plants like Lantana camara, Pinus kesiya, Litsea cubeba, Gaultheria fragrantissima, Mikania micrantha. Ambrosia artemisiifolia and Eupatorium riparium, which have a wide distribution in the state of Meghalaya and find application in medicinal practices of the local tribal population (Neogi et al., 1989; Chhetri, 2008; Hynniewta and Kumar, 2008; Kayang et al., 2008; Sinha et al., 2008; Sohkhlet, 2014). The present study is aimed at determining the oral toxicity and antifeedant activity of the above mentioned plants against fourth instar larvae of H. armigera (Hübner).

Materials and Methods

Collection of plants:

The seven plants selected for this study were collected from in and around Shillong city in Meghalaya. The selection of the plants was based on their local abundance, insecticidal properties and uses in traditional practices by the indigenous tribes of the state (Table 1). The samples were generally collected during the flowering and fruiting stage of the plants except *P. kesiya*. In case of *P. kesiya*, samples were collected from young plants aged between 10 to 12 years, mainly during spring and summer seasons when fresh needles emerged. The collected plants were identified by Dr. P. B. Gurung, Department of Botany, N. E. H. U., Shillong.

Sl. no.	Scientific name	Local Name	Site of Collection	Plant Part Used
1	Ambrosia artemisiifolia Linn. (Asteraceae)	Jaiawlong/kynbat Japan rit	N. E. H. U. campus, Shillong	Aerial parts
2	Eupatorium riparium Regel (Asteraceae)	Kynbat latnaiong	N. E. H. U. campus, Shillong	Leaves
3	Gaultheria fragrantissima Wall. (Ericaceae)	Sla thynrait	Mylliem, Upper Shillong.	Leaves
4	Lantana camara Linn. (Verbenaceae)	Soh Pyngkhlieh	N. E. H. U. campus	Aerial parts
5	Litsea cubeba (Lour.) Pers. (Lauraceae)	Dieng mosu/Dieng-si-ing	N. E. H. U. campus	Leaves
6	Mikania micrantha Kunth. (Asteraceae)	-	N. E. H. U. campus	Aerial parts
7	Pinus kesiya Royle (Pinaceae)	Dieng kseh	N. E. H. U. campus	Needles and tender branches.

Table 1 Details of the plant parts used and the collection sites of the seven plants selected for the study.

Preparation of plant extracts:

The plants were brought to the laboratory immediately after collection and washed with tap water thoroughly followed by a final rinse with dechlorinated water, following which, they were shade dried at room temperature (21 ± 2) °C) for 48-72 hours, depending on the plant. The dried plants were ground to coarse powder (~2mm) using an electric blender. The crude extracts were prepared using standard protocol (Harborne, 1998; Houghton and Raman, 1998; Kathuria and Kaushik, 2005; Handa et al., 2008; Deepa and Remadevi, 2011). For the preparation of extracts, 250 g of each of the plant powders was extracted with 2.5 litres methanol using a Soxhlet apparatus for 48 hours. Prior to extraction with methanol, the plant material was defatted with petroleum ether. The extracts were taken to dryness under reduced pressure using a rotary-vacuum evaporator and stored in airtight screw capped borosil containers at -20 °C for future use. Prior to performance of a bioassay, a standard stock solution of 2.5% w/v concentration was prepared by dissolving 2.5 g of the extract in 10 mL acetone and volume was made up to 100 ml by adding deionized water. From the stock 0.25%, 0.5% 1% solution, and w/v concentration was prepared for ingestion toxicity test and 0.1%, 0.2% and 0.5% w/v concentration for feeding deterrence test. Final volume for each of the test concentrations was 20 ml.

Test insect:

A laboratory culture of *H. armigera* larvae was maintained on a chickpea based semi-synthetic diet as suggested by Singh and Rembold (1992) under laboratory conditions $(21 \pm 2 \text{ °C}, 80 \pm 100 \text{ cm})$

5% R.H., and photoperiod of 12 L: 12 D). For the initial establishment of the colony in the laboratory, different instars of H. armigera larvae were collected from tomato crops grown in Mawiongsun village under Mawrynkneng tehsil in East Khasi Hills, District. The collected larvae were maintained on tomato leaves and fruits under laboratory conditions $(21 \pm 2 \text{ °C}, 80 \pm 5\% \text{ R}$. H. and photoperiod of 12 L: 12 D) in individual containers to prevent cannibalism and contamination until pupation. Pupae were transferred to clean containers with sterilized filter paper to facilitate moth emergence. Upon adult emergence, the male and female moths were paired and two pairs were released into individual mating chambers (2.5x1.6 feet). The adults were fed on a diet of 10% honey solution and provided with cotton strips as oviposition medium (Kaushik and Kathuria, 2004). From the first generation onwards, the laboratory colony was maintained on a chickpea based semi-synthetic diet. From the cultures, newly molted one-day old IV instar larvae were used for the bioassays.

Ingestion toxicity bioassay:

The larvicidal activity of plants was studied by oral application of the extracts through leaf dip method (Ramya *et al.*, 2008). Freshly collected tomato leaves were individually dipped in the three different concentrations (0.25%, 0.5% and 1% w/v) of each of the extracts and air dried. A single treated leaf was kept in a petri plate lined with moist filter paper and a single 6 hour starved fourth instar *H. armigera* larvae was introduced into the petri plate. Leaves treated with acetone were used as negative control while those treated with 100 ppm of Alphamethrin 10% EC (trade name: GEM)

were used as positive control. Alphamethrin 10% EC is a pyrethroid insecticide which demonstrates both contact and stomach toxicity against a wide range of insect pests (Indofil, 2016). Larval mortality was recorded after 24 hours of exposure. A total of 10 larvae were individually exposed to each treatment and each treatment was replicated thrice. The total number of subjects per treatment was 30 larvae. The mortality data were represented as corrected mortality using Abbott's correction (Abbot, 1925).

 $Ma\% = [(M_t - M_c) / (100 - M_c)] \times 100$

Ma% = corrected mortality(%), M_t = mortality in treatment(%), M_c = mortality in control(%).

Feeding deterrence bioassay:

The antifeedant activity of crude extracts was assayed using leaf disc method (Rani and Rajasekharreddy, 2009; Li et al., 2014). Discs of size 2.75cm² were punched from freshly collected tomato leaves and treated on each side with 15 µl of the test solution emulsified with 0.1% Triton X-100. The extracts were tested at three different concentrations-0.1%, 0.2% and 0.5% w/v. Leaf discs treated with acetone solution and emulsifier (0.1%) were used as control. The leaf discs were air dried and arranged in a petri plate with one treated and one control leaf disc per plate. A fourth instar larva of H. armigera was then introduced at the center of the petri plate, such that it was equidistant from the treated and the control discs. The experiment was thus conducted with one larva per petri plate with ten larvae per treatment and each treatment was replicated three times. After 6 hours, the leaf discs were removed and the area consumed by the larvae was measured using a graph sheet method. The feeding deterrence index was calculated by using the formula given by Bomford and Isman (1996):

$$FDI = \frac{C - T}{C + T} \times 100$$

Where, C = area of consumption in the control; T = area of consumption in the treatment.

Phytochemical analysis

The presence of different classes of phytochemicals in the plants demonstrating high oral toxicity and antifeedant activity was investigated qualitatively using standard procedures as described by Trease and Evans (1989), Sofowara (1993) and Harborne (1998).

Alkaloids

0.5 gm of the methanol extract was mixed with 8 ml of 1 % HCl, warmed and filtered. In a test tube, 2 ml of the filtrate was taken and a few drops of Dragendroffs reagent (solution of Potassium Bismuth Iodide) was added along the side of the test tube. Formation of red precipitate indicated presence of alkaloids.

Flavonoids

0.5 gm of the extract was shaken with petroleum ether to remove the lipid layer. The defatted residue was dissolved in 20 ml of 80 % ethanol and filtered. Three ml of the filtrate was mixed with 4 ml of 1 % potassium hydroxide solution in a test tube and the colour was observed. A dark yellow color indicated the presence of flavonoids.

Phenols

0.5 gm of the methanol extract was dissolved in 5 ml distilled water and then few drops of neutral 5% ferric chloride solution were added. A dark green colour indicated the presence of phenolic compounds.

Phytosterols

To 2 ml of the extract, 2 ml of chloroform was added followed by 2 ml of concentrated sulphuric acid. Formation of red colour in the chloroform layer indicated the presence of steroids.

Saponins

0.5 gm of the extract was dissolved in distilled water in a test tube and heated over a boiling water bath. The test tube was allowed to cool and then shaken vigorously. Formation of persistent froth indicated presence of saponins.

Tannins

0.5 gm of the extract was dissolved in 20 ml distilled water in a test tube and boiled. The solution was filtered and 1% aqueous ferric chloride was added to the filtrate. The appearance of intense green, purple, blue or black colour indicated the presence of tannins.

Terpenoids

5ml of the extract was mixed in 2 ml of chloroform, followed by the careful addition of 3 ml of concentrated sulphuric acid. A layer of reddish brown colouration at the interface indicated the presence of terpenoids.

Data analysis

The data obtained from the two bioassays were subjected to arcsine transformation prior to statistical analysis. The transformed data were then statistically analysed by one-way ANOVA. Separation of means and comparison between the different treatments was performed by Tukey's test at $P \le 0.05$. SPSS version 20 was used for the analysis.

Results

Ingestion toxicity bioassay

The larvicidal activity of methanolic extract of the seven plant species is presented in Table 2. All the seven plants demonstrated a dose dependent increase in oral toxicity, with percentage mortality of fourth instar larvae of H. armigera being highest at test concentration of 1% w/v. When tested at the concentration of 0.25%, methanol extract of all the seven plants demonstrated an average mortality of 15.78% which was statistically similar (p > 0.05) to the mortality rate of the larvae in the negative (solvent) control. However, at concentration of 0.5% and 1% w/v, the crude extracts of L.camara, L.cubeba and P.kesiya caused significantly higher mortality ($p \le 0.000$) than the negative control. The larvicidal activity demonstrated by the extract of aerial parts of Lantana camara against fourth instar larvae of H. armigera, was the highest amongst all the seven plants with percent corrected mortality

ranging from 27.77% to 53.33% across the test concentration. Its larvicidal activity was significantly higher than the negative control and the other plants ($p \le 0.05$) except *L. cubeba* (p = 0.672) and *P.kesiya* (p = 0.315). The methanol extract of the leaves of Litsea cubeba demonstrated the second highest oral toxicity against fourth instar larvae of H. armigera, with corrected larval mortality ranging between 20.37% and 51.48% across the test concentration. Of the remaining five plants, extract of Pinus kesiya caused 37.77% larval mortality at the highest concentration of 1% w/v and it was significantly higher (p=0.014) than the larval mortality in negative control, thereby making it the third best plant after L. camara and L. cubeba in terms of oral toxicity against fourth instar larvae of H. armigera. However, it is to be noted that the synthetic insecticide, Alphamethrin 10% EC, which was used as a reference/positive control in this bioassay, caused 100% larval mortality within 24 hours of exposure and its activity was significantly higher ($p \le 0.000$) than the activity of the plant extracts.

Feeding deterrence bioassay

The antifeedant activity of crude extracts of the selected plants was studied at three different concentrations. The feeding deterrence activity of the plants was assessed on the basis of the feeding deterrence index (FDI). Higher antifeedant/ feeding deterrence index indicates lower feeding by the test organism. All the seven plants demonstrated dose dependent increase in feeding deterrence but irrespective of the test concentration of the plant extracts, the antifeedance index of the negative (solvent) control was significantly lower ($p \le 0.0001$) in comparison to that of the plants (Table 3). Of the seven plants, the crude extract of G. fragrantissima demonstrated the highest antifeedant activity, causing 50.92% to 70.61% reduction in feeding by the fourth instar larvae of H. armigera, across the test concentration and thus its FDI was significantly higher (p \leq 0.05) than the other six plants. Apart from G. fragrantissima, crude extract of L. cubeba, P.

kesiya and *L. camara*, also caused high feeding deterrence, which was significantly higher than the remaining three plants with $p \le 0.05$. While the FDI on exposure to *L. cubeba* extract was 27.17% to 56.78% across test concentrations, *P. kesiya* extract reduced larval feeding by 36.07% to 49.39%; and, *L. camara* extract reduced larval feeding in the range of 18.66 % to 40.73% across the test concentrations.

Phytochemical analysis:

Based on the outcome of the two bioassays, crude extract of four out of the seven plants demonstrated high larvicidal and antifeedant activity. Hence, qualitative analysis of the methanol extracts of these four plants, namely, *G fragrantissima*, *L. camara*, *L. cubeba* and *P. kesiya*, was carried out for determination of the major phytochemical constituents present in them (Table 4). The outcome of the phytochemical analysis revealed that flavonoids, phenols and tannins were present in the extracts of all the four plants whereas alkaloids were detected only in extract of *P. kesiya*. Phytosterols were found in *L. camara* extract while terpenoids tested positive in the extracts of both *L. camara* and *L. cubeba*. Saponins were absent in the methanol extract of *L. camara* but present in the extracts of the other three plants.

 Table 2 The larvicidal (oral) toxicity of the crude extracts of the seven selected plants against fourth instar

 Helicoverpa armigera larvae.

Plant Name	Concentration of Extract (% w/v)			
	0.25%	0.5%	1%	
Ambrosia artemisiifolia	13.7 ± 5.48 ^b	13.70 ± 5.48^{def}	24.07 ± 5.25 ^{cd}	
Eupatorium riparium	7.04 ± 6.12 ^b	7.04 ± 6.12^{ef}	$13.33 \pm 5.77^{\text{ de}}$	
Gaultheria fragrantissima	13.33 ± 5.77^{b}	20.00 ± 10.00 ^{cde}	24.07 ± 5.25 ^{cd}	
Lantana camara	27.77 ± 6.93^{b}	44.81 ± 5.01^{b}	53.33 ± 8.01 ^b	
Litsea cubeba	20.37 ± 9.45^{b}	35.55 ± 3.85 bcd	51.48 ± 7.88^{b}	
Mikania micrantha	7.50 ± 6.61^{b}	7.87 ± 6.85^{ef}	18.52 ± 6.41^{d}	
Pinus kesiya	20.37 ± 9.45^{b}	34.44 ± 5.09^{bc}	37.77 ± 3.85 bc	
Negative control	3.33 ± 5.77 ^b	3.33 ± 5.77 f	3.33 ± 5.77^{e}	
Positive (reference)control	100^{a}	100 ^a	100 ^a	

Mean \pm SD represents mean percent corrected mortality of 3 replicates of 10 individuals each. Within columns, Means followed by the same alphabet do not differ significantly at 5% level of significance using Tukey's HSD test. Note: Chemical insecticide-Alphamethrin 10% EC at a concentration of 100 ppm was used as reference (positive) control while acetone was used as negative control.

Table 3 The antifeedant activity (feeding	deterrence) of the	e crude e	extracts of	f each	of the	seven	plants	against
fourth instar larvae of Helicoverpa armiger	ra.							

Plant Name	(Concentration of Extract (%w/v)			
	0.1%	0.2%	0.5%		
Ambrosia artemisiifolia	12.67 ± 1.44^{e}	12.42 ± 6.51^{cd}	17.12 ± 7.75^{d}		
Eupatorium riparium	12.83 ± 0.83^{e}	15.8 ± 9.85^{bc}	17.31 ± 5.31^{d}		
Gaultheria fragrantissima	52.27 ± 2.19^{a}	50.92 ± 11.21^{a}	70.61 ± 9.04^{a}		
Lantana camara	18.66 ± 2.95^{de}	35.44 ± 6.83^{ab}	40.73 ± 8.55^{bc}		
Litsea cubeba	$27.17 \pm 5.68^{\circ}$	46.61 ± 7.16^{a}	56.78 ± 4.93^{ab}		
Mikania micrantha	21.57 ± 1.31^{cd}	23.99 ± 6.03^{ab}	25.26 ± 5.92^{cd}		
Pinus kesiya	36.07 ± 1.05^{b}	43.57 ± 6.7^{a}	49.39 ± 5.25^{b}		
Negative (solvent) control	$6.10 \pm 1.21^{\rm f}$	5.50 ± 1.64^{d}	5.50 ± 1.64^{e}		

Mean \pm SD represents mean percent feeding deterrence of 3 replicates of 10 individuals each. Within columns, means followed by the same alphabet do not differ significantly at 5% level of significance using Tukey's HSD test. Acetone was used as negative control.

Table 4 The major phytochemical constituents presentin the methanol extracts of the four plantsdemonstrating insecticidal activity.

Classes of Phytochemicals	G. fragrantissima	L. camara	L. cubeba	P. kesiya
Alkaloids	-	-	-	+
Flavonoids	+	+	+	+
Phenols	+	+	+	+
Phytosterols	-	+	-	-
Saponins	+	-	+	+
Tannins	+	+	+	+
Terpenoids	-	+	+	-

(+) present; (-) absent.

Discussion

The ingestion toxicity bioassay revealed that larvicidal activity of the crude methanolic extract of the plants was much lower than that of the synthetic insecticide, Alphamethrin 10% EC. However, four out of the seven plants caused significantly higher ($p \le 0.05$) larval mortality as well as feeding deterrence in comparison to the solvent (negative) control indicating potent insecticidal activity against the notorious pest, *H. armigera*.

The results of the present study indicated that at higher concentrations, Lantana camara L. could act both as a potent oral toxicant and feeding deterrent against H. armigera larvae, and this result is in agreement with the findings of Prasad and Roy (2011), who had concluded from their histopathological study that extracts of L. camara could act as stomach poison in addition to some antifeedant activity against fourth instar H. armigera larvae. In a study by Murugesan and co-workers (2012), it was reported that essential oil of L. camara at a concentration range of 2500-10000 ppm caused 20-50% larval mortality after 24 hours exposure against third instar larvae of teak defoliator, Hyblaea puera while in another study, aqueous crude extract of L. camara leaves at a concentration of 40% caused 100% mortality of fourth instar larvae of Spodoptera litura (Deshmukhe et al., 2011). Both these studies found that larvicidal activity of L. camara increased with increase in its concentration which corroborated with our present findings.

Also, several authors have studied the antifeedant activity of L. camara (Deka et al., 1998; Ogendo et al., 2003; Murugesan and Murugesh, 2009). In a study by Arivoli and Tennyson (2013), at a concentration of 1%, the ethyl acetate crude extract of L. camara showed 25-50% antifeedance against third instar larvae of Spodoptera litura while the hexane and dichloromethane extracts showed < 25% antifeedance. Our study indicated higher activity of methanolic extract of L. camara, causing 40.74% feeding deterrence against fourth instar H. armigera larvae, at a much lower concentration of 0.5% w/v. However, it may be noted that different test organisms were used in the investigations conducted by other authors and many studies have shown that even closely related insect species can show widely di erent susceptibilities to the same extract or compound (Isman, 1993), which could be one of the reasons for the variation between the outcome of the present study and the previous studies.

Like L. camara, the crude extract of L. *cubeba* Pers. leaves showed significantly high (p ≤ 0.05) larvicidal and antifeedant activity against fourth instar larvae of H. armigera. Litsea cubeba is an important medicinal plant which finds wide application in traditional Chinese medicine (Kong et al., 2015). Apart from having antifungal properties against several pathogens (Nor Azah and Susiarti, 1999), it has also been found to have strong repellent, contact and fumigant toxicity as well as deterrent effect against stored product pests (Liu et al., 2007; Ko et al., 2009). Also, its essential oil has been found to be moderately effective as a contact against third instar toxicant larvae of Trichoplusia ni (Jiang et al., 2009) while Feng and co-workers (2012) reported that its ethanolic extract had strong feeding deterrence activity against third instar larvae of Spodoptera litura. To the best of our knowledge, the present study reports for the first time the toxicity of crude extract of Litsea cubeba against Helicoverpa armigera and a notable outcome is that at higher concentrations, it was moderately toxic (~43.51%) and demonstrated relatively high

antifeedance (~51.69%) against H. armigera.

Another important finding from this study is the larvicidal and antifeedant activity displayed by Pinus kesiva. This is the first record of the insecticidal activity of needle extracts of P. kesiya, although, it finds application in the traditional pest management practices of the indigenous tribes of Meghalaya (Sinha et al., 2008; Sokhlet, 2014). The methanolic extract of P. kesiya needles, at the highest tested concentration, caused close to 40% larval mortality apart from deterring larval feeding by Thus, insecticidal ~50%. the activity demonstrated by P. kesiya needle extract against fourth instar H. *armigera* larvae was significantly ($p \le 0.05$) higher than the negative (solvent) control. In fact, the FDI demonstrated by P. kesiya extract was comparable to the antifeedancy displayed by ethanolic and hexane extracts of Eucalyptus camaldulensis and Tylophora indica against fifth instar H. armigera (Kathuria and Kaushik, 2005). In a study by Kanis and co-workers (2009), a direct correlation was found between the lignin content of the acetone extracts of Pinus caribaea Morelet and their larvicidal activity against Aedes aegypti. Thus, although P. kesiya demonstrated moderate efficacy against H. armigera larvae in the present study, it may be a good candidate for future research.

During this study, the outcome of feeding deterrence bioassay was very encouraging with crude extract of G. fragrantissima demonstrating stronger antifeedant activity than plants like Tephrosia vogelii Hook (Leguminosae) and Solanum pseudocapsicum (Solanaceae) which have been reported to show promising antifeedant activity against larval stages of H. armigera by Jeyashankar and co-workers (2012) and Arivoli and Tennyson (2013), respectively. Although no earlier reports on the insecticidal activity of crude extracts of G. fragrantissima were found but several authors have studied the larvicidal, pupicidal, antifeedant and repellent activities of the essential oil of Gaultheria species (Senthilkumar and Venkatesalu, 2012; Ranyaphi et al., 2012; Jeyasankar, 2012; 2014). Palanimuthu et al., Gaultheria *fragrantissima* is rich in essential oil which is a constituent of several insecticidal and insect repellent preparations (Ranyaphi *et al.*, 2012) and therefore future investigation on its bioactivity could enable the development of an effective antifeedant against larval stages of polyphagous pests like *H. armigera*.

An important finding of the present study is that three out of four plants demonstrating high antifeedant activity also caused maximum larval mortality. Similar findings have been reported by several authors (Chen et al., 1996; Koul et al., 2004; Ling et al., 2008). According to a study by Lingathurai and co-workers (2011), chloroform extract of Acalypha fruticosa Forssk leaves demonstrated maximum antifeedance and oral toxicity against third instar larvae of Plutella xylostella L.; the authors attributed the two different modes of action of the extract to the presence of five different phytochemical groups namely, terpenoids, tannins, coumarins, anthraquinones and saponins. In the present study too, the qualitative analysis revealed the presence of phytochemical groups like phenols, flavonoids, tannins, terpenoids, saponins and alkaloids in the methanol extract of G. fragrantissima, L. camara, L. cubeba and P. kesiya.

The insecticidal activity of plants is presence of various attributed to the phytochemical groups (Kabaru and Gichia, 2001) and the occurrence of more than one major class of phytocompounds is responsible for the different modes of action of plant extracts against the target pests (Park et al., 2002; Lingathurai et al., 2011). The extract of the four plants, namely, G. fragrantissima, L. camara, L. cubeba and P. kesiya, tested positive for phenolic compounds, flavonoids and tannins. All three groups of phytocompounds have been reported to affect herbivorous insect's growth and development either by feeding inhibition or through post-ingestive phenomena (Coley et al., 1985; Barbehenn et al., 2001; Hoffman-Campo et al., 2001; Lago et al., 2002; Treutter, 2006; Jadhav et al., 2012). In addition, extracts of L. camara and L. cubeba also tested positive for terpenoids. Terpenoids

in plants can act mainly as antifeedant and growth disruptor and possess considerable toxicity toward insects (Kubo and Nakanishi, 1978; Khalid et al., 1989). Saponins on the other hand are a class of phytochemicals which are reported to be insecticidal by many investigators (Marston and Hostettmann, 1985; Jeong et al., 2004; Sparg et al., 2004; McGaw et al., 2008). In the present study, saponins were found to be present in the methanol extract of G. fragrantissima, L. cubeba and P. kesiya. Thus, the insecticidal and antifeedant activity demonstrated by the methanol extracts of G. fragrantissima, L. camara, L. cubeba and P. kesiva could be the result of composite effect of all these classes of phytocompounds.

However, the present study is a preliminary investigation which indicates that crude methanol extracts of the four plants possess insecticidal property. Future research has to be conducted with these plants to understand their exact mode of action/s as well as isolate and identify the bioactive compound/s responsible for the toxicity demonstrated towards the target pest.

Conclusion

From the present study it can be concluded that out of seven selected plants, four plants namely, *L. camara*, *G. fragrantissima*, *L. cubeba* and *P. kesiya* have demonstrated promising insecticidal activity against *H. armigera* larvae. Further research on the bioactivity of these commonly found plants can lead to the development of a cost effective, eco-friendly formulation for crop protection, which will be beneficial to farmers of states like Meghalaya where organic farming is being encouraged by the Central and the State governments.

Acknowledgement

The authors are thankful to Dr. P. B. Gurung, Herbarium section, Department of Botany, N. E. H. U., Shillong, for his help in identification of the plant samples.

References

- Abbott, W. S. 1925. A method for computing the effectiveness of an insecticide. Journal of Economic Entomology, 18: 265-267.
- Arivoli, S. and Tennyson, S. 2013. Antifeedant activity, developmental indices and morphogenetic variations of plant extracts against *Spodoptera litura* (Fab) (Lepidoptera: Noctuidae). Journal of Entomology and Zoology Studies, 1 (4): 87-90.
- Barbehenn, R. V., Bumgarner, S. L., Roosen, E. F. and Martin, M. M. 2001. Antioxidant defenses in caterpillars: role of the ascorbate-recycling system in the midgut lumen. Journal of Insect Physiology, 47 (4-5): 349-357.
- Bomford, M. K. and Isman, M. B. 1996. Desensitization of fifth instar *Spodoptera litura* to azadirachtin and neem. Entomologia Experimentalis et Applicata, 81: 301-313.
- Chaturvedi, I. 2007. Status of insecticide resistance in the cotton bollworm, *Helicoverpa armigera* (Hübner). Journal of Central European Agriculture, 8 (2): 171-182.
- Chen, H., Liu, Y. H., Den, X. P. and Zhao, Z. M. 1996. Effects of the extracts of balsam pear leaves on feeding, growth and survival of larvae of *Bombyx mori* and *Pieris rapae*. Southwest China Journal of Agricultural Science, 9: 68-71.
- Chhetri, R. 2008. Trends in ethnodomestication of some wild plants in Meghalaya, Northeast India. Indian Journal of Traditional Knowledge, 5 (3): 342-347.
- Coley, P. D., Bryant, J. P. and Chapin, S. 1985. Resource availability and plant anti herbivore defence. Science, 230: 895-899.
- Deepa, B and Remadevi, O. K. 2011. Insecticidal Activity of the Phytoextracts derived from different parts of the trees of Fabaceae family against *Hyblaea puera* Cramer (Lepidoptera: Hyblaeidae). Biological Forum, 3 (2): 1-8.
- Deka, M. K., Singh, K. and Handique, R. 1998. Antifeedant and repellent effects of pongam (*Pongamia pinnat*a) and wild sage (*Lantana camara*) on tea mosquito bug (*Helopeltis theivora*). Indian Journal of Agricultural

Science, 68: 274-276.

- Deshmukhe, P. V., Hooli, A. A. and Holihosur, S. N. 2011. Effect of *Lantana camara* (L.) on growth, development and survival of tobacco caterpillar (*Spodoptera litura* Fabricius). Karnataka Journal of Agricultural Science, 24 (2): 137-139.
- Fakrudin, B., Kumar, V., Krishnareddy, K., Patil, B. and Kuruvinashetti, M. 2004. Morphometric differences between pyrethroid resistant and susceptible populations of cotton bollworm, *Helicoverpa armigera*. Resistant Pest Management Newsletter, 13: 18-19.
- Feng, X., Jiang, H., Zhang, Y., He, W. and Zhang, L. 2012. Insecticidal activities of ethanol extracts from thirty Chinese medicinal plants against *Spodoptera exigua* (Lepidoptera: Noctuidae). Journal of Medicinal Plants Research, 6 (7): 1263-1267.
- Handa, S. S. 2008. An Overview of Extraction Techniques for Medicinal and Aromatic Plants, In: Handa, S. S., Khanuja, S. P. S., Longo, G. and Rakesh, D. D. (Eds.), Extraction Technologies for Medicinal and Aromatic Plants, International Centre for Science and High Technology, Trieste, pp. 22-30.
- Harborne, J. B. 1998. Phytochemical Methods-A Guide to Modern Techniques of Plant Analysis. Chapman and Hall Publishers. London. pp. 1-32.
- Hoffman-Campo, C. B., Harborne, J. B. and McCaffery, A. R. 2001. Preingestive and post-ingestive effects of soya bean extracts and rutin on *Trichoplusia ni* growth. Entomologia Experimentalis et Applicata, 98: 181-194.
- Hongo, H. and Karel, A. K. 1986. Effect of plant extract on insect pest of common beans. Zeitschrift Fur Angewandte Entomologie, 102: 164-169.
- Houghton, P. J. and Raman, A. 1998. Laboratory Handbook for The Fractionation of Natural Extracts. Chapman and Hall Publishers. London.
- Hynniewta, S. R. and Kumar, Y. 2008. Herbal remedies among the *Khasi* traditional healers and village folks in Meghalaya. Indian Journal of Traditional Knowledge, 7

(4): 581-586.

- Ignacimuthu, S. and Jayaraj, S. 2003. Ecofriendly approaches for sustainable pest management. Current Science, 84: 10-25.
- Indofil. 2016. Indofil industries limited. Available on: https://indofilcc.com/businessarea/agricultural-chemicals/insecticides/gem /(Accessed on 2nd May, 2016).
- Isman, M. B. 1993. Growth inhibitory and antifeedant effects of azadirachtin on six noctuids of regional economic importance. Pesticide Science, 38: 57-63.
- Jadhav, D. R., Mallikarjuna, N., Rathore, A. and Pokle, D. 2012. Effect of some flavonoids on survival and development of *Helicoverpa* armigera (Hbner) and Spodoptera litura (Fab) (Lepidoptera: Noctuidae). Asian Journal of Agricultural Sciences, 4 (4): 298-307.
- Jeong, S. I., Kim, K. J., Choo, Y. K., Keum, K. M., Choi, B. K. and Jung, K. Y.2004. *Phytolacca americana* inhibits the high glucose-induced mesangial proliferation via suppressing extracellular matrix accumulation and TGF-b production. Phytomedicine, 11: 175-181.
- Jeyasankar, A. 2012. Antifeedant, insecticidal and growth inhibitory activities of selected plant oil on black cutworm, *Agrotis ipsilon* (Hufnagel) (Lepidoptera: Noctucidae). Asian Pacific Journal of Tropical Diseases, 2: 347-351.
- Jeyasankar, A., Premalatha, S. and Elumalai, K. 2012. Biological activities of Solanum pseudocapsicum (Solanaceae) against cotton bollworm, Helicoverpa armigera Hübner and armyworm, Spodoptera litura Fabricius (Lepidotera: Noctuidae). Asian Pacific Journal of Tropical Biomedicine, 2 (12): 981-986.
- Jiang, Z. L., Akhtar, Y., Bradbury, R., Zhang, X. and Isman, M. B. 2009. Comparative toxicity of essential oils of *Litsea pungens* and *Litsea cubeba* and blends of their major constituents against the cabbage looper, *Trichoplusia ni*. Journal of Agricultural and Food Chemistry, 57: 4833-4837.
- Jotwani, M. G. and Srivastava, K. P. 1984. A review of neem research in India in relation to insects. Proceedings of International

Downloaded from jcp.modares.ac.ir on 2024-05-03]

Neem Conference, Ranicholzhausen, Germany, pp. 43-56.

- Kabaru J.M. and Gichia L. 2001. Insecticidal activity of extracts derived from different parts of the mangrove tree *Rhizophora mucronata* (rhizophoraceae) Lam. against three anthropods. African Journal of Science and Technology (AJST). Science & Engineering Series 2 (2): 44-49.
- Kanis, L., Antonio, R., Antunes, É., Prophiro, J. and Silva, O. 2009. Larvicidal effect of dried leaf extracts from *Pinus caribaea* against *Aedes aegypti* (Linnaeus, 1762) (Diptera: Culicidae). Revista da Sociedade Brasileira de Medicina Tropical, 42 (4): 373-376.
- Kathuria, V. and Kaushik, N. 2005. Feeding Inhibition of *Helicoverpa armigera* (Hübner) by *Eucalyptus camaldulensis* and *Tylophora indica* extracts. Insect Science, 12: 249-254.
- Kaushik, N. and Kathuria, V. 2004. *Helicoverpa* (Heliothis) and botanical pesticides: an overview. In: Prasad, D. and Singh, A. (Eds.), Advances in Plant Protection Sciences, Akansha Publishing House, Darya Ganj, New Delhi, India, pp: 156-166.
- Kayang, H., Kharbuli, B. and Syiem, D. 2008. *Gaultheria fragrantissima* Wall.-an untapped economic plants species of Meghalaya. Natural Product Radiance, 7 (5): 400.
- Khalid, S. A., Duddeck, H. and Gonzalez-Sierra, M. 1989. Isolation and characterization of an antimalarial agent of the neem tree *Azadirachta indica*. Journal of Natural Products, 52: 922-926.
- Ko, K., Juntarajumnong, W. and Chandrapatya, A. 2009. Repellency, fumigant and contact toxicities of *Litsea cubeba* (Lour.) Persoon against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). Kasetsart Journal, 43:56-63.
- Koul, O., Kaur, H., Goomber, S. and Wahab,
 S., 2004. Bioefficacy and mode of action of rocaglamide from *Aglaia elaeagnoidea* (syn *A. roxburghiana*) against gram pod borer, *Helicoverpa arm igera* (Hubner). Journal of Applied Entomology, 128: 177-181.
- Koul, O., Multani, J. S., Singh, G. and Wahab, S. 2002. Bioefficacy of Toosendanin from

Melia dubia (syn. *M. azedarach*) against gram pod-borer *Helicoverpa armigera* (Hubner). Current Science, 83: 1387-1391.

- Kubo, I. and Nakanishi, K. 1978. Some terrpenoid insect antifeedants from tropical plants-I. Proceedings of 4th International Congress of Pesticide Chemistry, Zurich, Switzerland, p. 284.
- Lago, J. H. G., Brochini, C. B. and Roque, N. F. 2002. Terpenoids from *Guarea guidonia*. Phytochemistry, 60: 333-338.
- Lammers, J. W. and Macleod, A. 2007. Report of a pest risk analysis: *Helicoverpa armigera (Hübner, 1808)*. European Union. Retrieved from https://secure.fera. defra.gov.uk/phiw/riskRegister/plant health/ documents/helicoverpa.pdf (Accessed on 13 Jun, 2013).
- Li, W., Hu, J., Yang, J., Yuan, G., Guo, X. and Luo, M. 2014. Feeding deterrence of common spices against *Helicoverpa armigera* larvae. Advances in Bioscience and Biotechnology, 5: 1025-1031.
- Ling, B., Wang, G., Ya, J., Zhang, M. and Liang, G. 2008. Antifeedant activity and active ingredients against *Plutella xylostella* from *Momordica charantia* leaves. Agricultural Sciences in China. 7: 466-1473.
- Lingathurai, S., Vendan,S. E, Paulraj, M. G. and Ignacimuthu, S. 2011. Antifeedant and larvicidal activities of *Acalypha fruticosa* Forssk. (Euphorbiaceae) against *Plutella xylostella* L. (Lepidoptera: Yponomeutidae) larvae. Journal of King Saud University (Science), 23: 11-16.
- Liu, S., Shi, J., Cao, H., Jia, F., Liu, X. and Shi, G. 2000. Survey of pesticidal component in plant. In: Dianmo, L. (Ed.), Entomology in China in 21st Century, Proceedings of Conference of Chinese Entomological Society, Science & Technique Press, Beijing, China. pp: 1098-1104.
- Liu, Z., Goh, S. and Ho, S. 2007. Screening of Chinese medicinal herbs for bioactivity against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). Journal of Stored Products Research, 43 (3): 290-296.
- Manjunath, T., Bhatnagar, V., Pawar, C. and

Downloaded from jcp.modares.ac.ir on 2024-05-03

J. Crop Prot.

Sithanantham, S. 1985. Economic importance of *Heliothis* spp. in India and assessment of their natural enemies and host plants. Proceedings of the Workshop on Biological Control of *Heliothis*: Increasing the Effectiveness of Natural Enemies, New Delhi, p: 197-228.

- Marston, A. and Hostettmann, K. 1985. Plant molluscicides. Phytochemistry, 2: 639-652.
- McCaffery, A. R. 1998. Resistance to insecticides in Heliothine lepidoptera: A Global View. Philosophical Transactions of the Royal Society B: Biological Sciences, 353 (1376): 1735-1750.
- McGaw, L. J., Lall, N., Meyer, J. J. and Eloff, J. N. 2008. The potential of South African plants against mycobacterium infections. Journal of Ethnopharmacology, 119: 482-500.
- Murugesan, N. and Murugesh, T. 2009. Bioefficacy of some plant products against brinjal fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Pyrallidae). Journal of Biopesticides, 2 (1):60-63.
- Murugesan, S., Rajeshkannan, C., Suresh Babu, D., Sumathi, R. and Manivachakam, P. 2012. Identification insecticidal properties in common weed - *Lantana camara* Linn. by gas chromatography and mass spectrum (GC-MS-MS). Advances in Applied Science Research, 3 (5): 2754-2759.
- Neogi, B., Prasad, M. N. V. and Rao, R. R. 1989. Ethnobotany of some weeds of Khasi and Garo Hills, Meghalaya, Northeastern India. Economic Botany, 43 (4): 471-479.
- Nor Azah M. A. and Susiarti, S. 1999. *Litsea cubeba* (Lour.) Parson in essential oil plants.
 In: L. Oyen, P. A. and Dung, N. X. (Eds.), Plant Resources of South East Asia, Vol. 19, Backhuys Publishers., Leiden, pp: 123-126.
- Ogendo, J. O., Belmain, S. R., Deng, A. L. and Walker, D. J. 2003. Comparison of toxic and repellent effects of *Lantana camara* L. with *Tephrosia vogelii* Hook. and a synthetic pesticide against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae) in stored maize grain. Insect Science Application, 23 (2): 127-135.

- Palanimuthu, A., Kadarkarai, M., Anbazhagan, P. and Sujitha, R. 2014. Larvicidal, pupicidal and repellent activities of *Gaultheria* Oil (Plantae: Ericaceae) against the filarial vector, *Culex quinquefasciatus* (Insecta: Diptera: Culicidae). Journal of Entomology and Zoology Studies, 2 (4):290-294.
- Pandey, U. K., Kumar, A. S., Kumar, A. S. and Mamta, P. 1983. Evaluation of some plant origin insecticides against gram caterpillar, *Helicoverpa armigera* Hubner. Indian Journal of Entomology, 45 (2): 211.
- Park, S. B., Lee, E. S., Choi, S. W., Jeong, Y. C., Song, C. and Cho, Y. W. 2002. Insecticidal and acaricidal activity of piperonndine and piperoctadealidne derived from dried fruits of *Piper longum* L. Crop Protection, 21: 249-251.
- Prasad, A. and Roy, S. 2011. Histoarchitechtural alterations in the midgut tissues of fourth instar larvae of grampod borer, *Helicoverpa* armigera (Hub) fed with leaf extract of plant, *Lantana camara* (L). International Journal of Pharmacy and Biological Sciences, 2 (4): B (613)-B (620).
- Ramya, S., Rajasekaran, C., Sundararajan, G, Alaguchamy, N. and Jayakumararaj, R. 2008. Antifeedant activity of leaf aqueous extracts of selected medicinal plants on VI instar larva of *Helicoverpa armigera* (Hübner), Ethnobotanical Leaflets, 12: 938-943.
- Rani, P. U. and Rajasekharreddy, P. 2009. Toxic and antifeedant activities of Sterculia Foetida (L.) seed crude extract against *Spodoptera litura* (F.) and *Achaea Janata* (L.). Journal of Biopesticides, 2 (2): 161-164.
- Ranyaphi, R. A., Mao, A. A. and Borthakur, S.
 K. 2012. Invitro plant regeneration of wintergreen (*Gaultheria fragrantissima* Wall.): assessment of multiple nutrient formulations and cytokinin types. Indian Journal of Biotechnology, 11: 197-204.
- Sahayaraj, K. 1998. Antifeedant effect of some plant extracts on the Asian armyworm, *Spodoptera litura* (Fabricius). Current Science, 74: 523-525.
- Senthilkumar, A. and Venkatesalu, V. 2012. Larvicidal potential of *Acorus calamus* L.

essential oil against filarial vector mosquito *Culex quinquefasciatus* (Diptera: Culicidae). Asian Pacific Journal of Tropical Disease, 2 (4): 324-326.

- Setiawati, W., Somantri, A. and Duriat, A. 2000. Effect of population density and infestation of *Helicoverpa armigera* Hübner on tomato yield loss and its control. Journal of Horticulture, 10: 112-120.
- Shaalan, E., Canyon, D., Younes, M., Abdel-Wahab, H. and Mansour, A. 2005. A review of botanical phytochemicals with mosquitocidal potential. Environment International, 31: 1149-1166.
- Singh, A. K. and Rembold, H. 1992. Maintenance of the cotton bollworm, *Heliothis armigera* Hübner (Lepidoptera: Noctuidae) in laboratory culture-I. Rearing on semi-synthetic diet. International Journal of Tropical Insect Science, 13: 333-338.
- Sinha. B., Dey, S. and Kalita, J. 2008. How the hill farmers control pests using locally available resources: Lessons from the upland areas of North East India. Available on SSRN: http://ssrn.com/abstract = 1303990. doi:10.2139/ssrn.1303990 (Accessed on March 21st 2012).
- Sivagnaname, N., and Kalyanasundaram, M. 2004. Laboratory evaluation of methanolic extract of *Atlantia monophylla* (Family: Rutaceae) against immature stages of mosquitoes and non-target organisms. Memórias do Instituto Oswaldo Cruz, 99 (1): 115-118.
- Sofowora A. 1993. Medicinal plants and Traditional Medicine in Africa. Spectrum Books Ltd, Ibadan, Nigeria. pp. 289.

Sohkhlet, M. 2014. Evaluation of traditional

insect-pest management practices in agricultural crops of Khasi and Jaintia hills, Meghalaya. Ph. D. Dissertation, North Eastern Hill University, Shillong. 41 pp.

- Sparg, S. G., Light, M. E. and Staden, J. V. 2004. Biological activities and distribution of plant saponins. Journal of Ethnopharmacology, 94: 219-243.
- Sundararajan, G. and Kumuthakalavalli, R. 2001. Antifeedant activity of aqueous extract of *Gnidia glauca* Gilg. and *Toddalia asiatica* Lam. on the gram pod borer, *Helicoverpa armigera* (Hbn). Journal of Environmental Biology, 22 (1):11-14.
- Thakur, N., Shylesha A. and Rao, R. 2006. Major insect pests of agri-horticultural crops and their management in north eastern hill region. In: Dwivedi, S. (Ed.), Integrated Pest Management and Biocontrol 1st ed. Pointer Publ. Pvt. Ltd., New Delhi, pp: 196-99.
- Trease, G. E. and Evans, W. C. 1989. Pharmacology 11th ed., Bailliere Tindall Ltd., London, pp: 60-75.
- Treutter, D. 2006. Significance of flavonoids in plant resistance: a review. Environmental Chemistry Letters, 4 (3): 147-157.
- Wambua L. M., Deng A. L., Ogendo J. O., Owuoche J. and Bett P. K. 2011. Toxic, antifeedant and repellent activity of aqueous crude extracts of *Tephrosia vogelii* Hook on the larval stages of *Helicoverpa armigera* Hubner. Baraton Interdisciplinary Research Journal, 1: 19-29.
- Yang, Y., Li, Y. and Wu, Y. 2013. Current status of insecticide resistance in *Helicoverpa armigera* after 15 years of Bt cotton planting in China. Journal of Economic Entomology, 106 (1): 375-81.

اثرات لاروکشی و ضدتغذیهای برخی گیاهان بومی مگالایا علیه لارو سن چهارم کرم غوزه پنبه (Hübner) Helicoverpa armigera

ديبيندو پاول و موميتا چودهاری*

گروه مطالعات محیطی، دانشگاه هیل شمال شرقی، شیلونگ-۷۹۳۰۲۲، مگالایا، هندوستان. * پست الکترونیکی نویسنده مسئول مکاتبه: moumita.choudhury@gmail.com دریافت: ۵ بهمن ۱۳۹۴؛ پذیرش: ۲۳ خرداد ۱۳۹۵

چکیده: در این مطالعه، اثرات لاروکشی و ضدتغذیهای هفت گیاه بومی شامل Pinus kesiya Royle (Pinaceae), Lantana camara Linn. (Verbenaceae), Litsea cubeba Lour. (Lauraceae), Gaultheria fragrantissima Wall. (Ericaceae), Mikania micrantha Kunth.(Asteraceae), Ambrosia *eupatorium riparium* Regel (Asteraceae) و *artemisiifolia* Linn.(Asteraceae) که در مگالایای کشور هند گسترش وسیعی دارند، عیله لاروهای سن چهارم کرم غوزه پنبه Helicoverpa armigera (Hübner) در شرایط آزمایشگاهی مورد بررسی قرار گرفت. تأثیر عصارههای تمام گیاهان با افرایش غلظت افزايش يافتند. اما با اين حال تأثير چهار گياه شامل I. cubeba ، G fragrantissima L. camara، و P. kesiya بهطور معنى دارى نسبت به كنترل منفى (حلال) و عصاره هاى P. kesiya بهطور معنى دارى نسبت به كنترل منفى and M. micarantha بالاتر بود. عصاره متانولي L. camara موجب بالاترين مركومير مابين ۲۷/۷۷ و G درصد در غلظتهای مورد آزمایش (۲۵/۰۰ \wedge ۱ و ۱ درصد w/v) شد درحالی که عصاره Gfragrantissima بالاترین تأثیر بازدارنده تغذیه در حدود ۵۰/۹۲ و ۷۰/۶۱ درصد بهترتیب در غلظتهای ۰/۱ و ۵/۰ درصد بود. عصاره برگهای L. cubeba دارای سمّیت گوارشی و ضدتغذیهای زیادی در بالاترین غلظت بود. این درحالی است که عصاره برگهای سوزنی کاج P. kesiya در بالاترین غلظت اثرات سمّیت گوارشی و ضدتغذیهای متوسطی داشت. تجزیه فیتوشیمی این چهار گیاه نشان داد که عصاره گیاهان فوق حاوی پنج گروه مختلف از ترکیبات شیمیایی هستند که هر کدام از آنها برای آفات مضر و زیانبار هستند. بنابر این می توان نتیجه گیری نمود که چهار گیاه از هفت گیاه مورد مطالعه دارای تأثیر حشره کشی مناسبی هستند که لازم است برای تولید حشره کشهای گیاهی مؤثر مورد بررسیهای بیشتری قرار گیرند.

واژگان كليدى: عصارەھاى گياھى، سمّيت گوارشى، فعاليت ضدتغذيەاى، كرم غوزە پنبە، مگالايا