

## Research Article

# Development, reproduction and life table parameters of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on five main host plants

Yaghoub Fathipour<sup>1\*</sup>, Fatemeh Bagheri<sup>1</sup>, Abdoolnabi Bagheri<sup>2</sup> and Bahram Naseri<sup>3</sup>

1. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

2. Plant Protection Research Department, Hormozgan Agricultural and Natural Resources Research and Education Center, Agricultural Research Education and Extension Organization (AREEO), Bandar Abbas, Iran.

3. Department of Plant Protection, Faculty of Agriculture, University of Mohaghegh Ardabili, Ardabil, Iran.

**Abstract:** The cotton bollworm, *Helicoverpa armigera* (Hübner), is a destructive pest on a wide range of economic crops in many parts of the world. In this research, demographic parameters of *H. armigera* were determined on five host plant species including cowpea (cv. Mashhad), navy bean (cv. Dehghan), chickpea (cv. Hashem), soybean (cv. 033) and corn (cv. Single cross 704). All experiments were carried out under laboratory conditions at  $25 \pm 1^\circ\text{C}$ ,  $65 \pm 5\%$  RH and a photoperiod of 16: 8 (L: D) h. The results revealed that females and males had the shortest development time on chickpea (36.16 and 34.98 d, respectively) and the longest development time on corn (42.00 and 42.95 d, respectively). The highest daily and total fecundity of *H. armigera* were observed on cowpea and the lowest ones were on corn. The values of the intrinsic rate of increase ( $r_m$ ) on the above-mentioned host plants were 0.180, 0.144, 0.161, 0.161 and 0.126  $\text{day}^{-1}$ , respectively. Also, the values of the net reproductive rate ( $R_0$ ) were 365.66, 294.28, 365.67, 239.69 and 147.40 female offspring, respectively. The longest mean generation time ( $T$ ) ( $37.90 \pm 0.26$  d) and doubling time ( $DT$ ) ( $5.62 \pm 0.17$  d) were observed on corn. Our findings revealed that cowpea and corn were the most susceptible and resistant host plants to *H. armigera*, respectively.

**Keywords:** cotton bollworm, *Helicoverpa armigera*, corn, cowpea

## Introduction

The cotton bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) is a cosmopolitan and destructive pest, feeding on a diverse array of host plant species (Fathipour *et al.*, 2019). It frequently attacks over 200 crop species including cotton,

maize, chickpea, pigeon pea, sorghum, tomato and soybean (Sedaratian *et al.*, 2014). The neonate larvae feed on different parts of host plants such as flower, bud, stem and leaves (Garcia, 2006). It has been shown that chemical control cannot be a permanent tool for the control of this pest either due to the environmental problems which restrict the long-term application of pesticides or growing resistance to the different classes of chemicals (Sedaratian *et al.*, 2013). Use of genetically resistant host plants is an important approach which helps to diminish

Handling Editor: Ali Asghar Talebi

\*Corresponding author: fathi@modares.ac.ir

Received: 01 December 2019, Accepted: 20 July 2020

Published online: 04 August 2020

the damage of *H. armigera*. Also, it can be integrated successfully with other non-chemical managing approaches (Fathipour and Sedaratian, 2013; Fathipour *et al.*, 2019).

Different host plants can affect the feeding behavior of insect pests positively or negatively (Jafari *et al.*, 2020) through targeting their development time, survival rate, reproduction and life table parameters (Golizadeh *et al.*, 2009; Soufbaf *et al.*, 2010a, b; Soufbaf *et al.*, 2012). Demographic parameters provide a simple way to extract detailed information about the population growth potential of an insect pest in the current and the next generations (Frei *et al.*, 2003). Also, understanding the demographic parameters of a pest helps us to assess the level of resistance to insect pests in different host plants. Such information can be utilized appropriately in integrated pest management (IPM) programs.

Although, valuable information exists on the demographic parameters of *H. armigera* on each of these host plants separately (Ali *et al.*, 2009; Naseri *et al.*, 2009a,b; Soleimannejad *et al.*, 2010; Naseri *et al.*, 2011; Karimi *et al.*, 2012), no information is available regarding comparative demography of *H. armigera* on these host plants. In this study, we aimed to study the demographic parameters of *H. armigera* on the above-mentioned host plants using artificial diet prepared from their seeds. The artificial diet provides suitable conditions for successive rearing of *H. armigera* to assess its potential on these host plants. Such information would be useful in the integrated pest management of *H. armigera* especially in areas where these plant species are planted at the same time.

## Materials and Methods

### Plant seeds

The seeds of five different host plants including cowpea (cv. Mashhad), chickpea (cv. Hashem), soybean (cv. 033), navy bean (cv. Dehghan) and corn (cv. Sc704) were obtained from Seed and Plant Improvement Institute, Karaj, Iran.

### Rearing methods and experimental conditions

The eggs of *H. armigera* were prepared from a laboratory culture kept on an artificial diet at the Tabriz University, Iran. Five separate stock cultures of *H. armigera* were reared for two generations on artificial diets based on the seeds of five host plants. The rearing was performed in a growth chamber set at  $25 \pm 1$  °C,  $65 \pm 5\%$  RH, and a photoperiod of 16: 8 (L: D) h.

The artificial diets were prepared as follows: powdered seed of each host plant (250 g), wheat germ (30 g) as protein and carbohydrate sources, sorbic acid (1.1 g) as an antimicrobial agent, ascorbic acid (3.5 g) as a vitamin source, sunflower oil (5 ml) as a preservative, agar (14 g) as a moisturizer, methyl-p-hydroxyl benzoate (2.2 g), formaldehyde 37% (2.5 g) and distilled water (650 ml) (Teakle, 1991).

To obtain the same-aged eggs, 15-20 pairs of adult moths (both sexes) of *H. armigera* emerged from the pupae were transferred to the mating cages (14 cm diameter and 19 cm height) for 72 h. The mating cages were covered at the top with a fine mesh net. A small cotton roll soaked in 10% honey solution was applied for supplying the adults. The eggs laid on the net were harvested for running the experiment.

### Determining demographic parameters

Seventy eggs were used to determine survivorship and development of *H. armigera* on each host plant. All eggs were checked daily and the number of emerged larvae was recorded. The newly hatched larvae were transferred individually into plastic containers using fine camel hair brush. The plastic containers were 8 cm in diameter and 5 cm in height with a hole covered by a fine mesh net for ventilation. Fresh artificial diet was provided as required and observations were made daily to record the mortality/survival of larvae in the same instar or molting to the next instar up to adult emergence. The larval instars were recognized by measuring the head capsules or presence of exuviae from molting. The fifth instar larvae were kept in the plastic containers for prepupation and pupation. The

pupae were also checked daily until all adults emerged or the pupae died. At pupation, the gender was determined and recorded.

After adults' emergence, a pair of female and male moths was transferred into the oviposition cages (as described above). A small cotton wick soaked in 10% honey solution was placed in the oviposition cages to nourish the adults. The adults were introduced into new oviposition cages daily and the laid eggs were counted and harvested once a day. The adult longevity, oviposition period and fecundity were determined until the death of the last female in the cohort.

The life table parameters of *H. armigera* on their hosts were determined using the formulae explained by Carey (1993) and the net reproductive rate ( $R_0$ ), intrinsic rate of increase ( $r_m$ ), finite rate of increase ( $\lambda$ ), mean generation time ( $T$ ) and doubling time ( $DT$ ) were estimated. In addition, age-specific survivorship ( $l_x$ ) and life expectancy ( $e_x$ ) were calculated.

### Data analysis

To estimate the variance for  $r_m$  and other life table parameters, the jackknife method was used (Meyer *et al.*, 1986). The technique is based on the repeated recalculation of the required estimator and missing out each sample in turn (Maia *et al.*, 2000). The jackknife

pseudo-values for hosts were tested for normality and then subjected to an analysis of variance (ANOVA) (PROC GLM, SAS Institute). Based on the results, mean comparison was done based on Tukey test ( $P < 0.05$ ) (SAS version 9.4).

### Cluster analysis

A dendrogram was constructed for the studied hosts based on the demographic parameters estimated for *H. armigera* on these host plants using the statistical software SPSS (SPSS version 16.0).

## Results

### Development of immature stages

Except incubation time, the other immature stages had significant differences on different hosts studied. The larvae fed on the corn diet had a longer larval period ( $23.52 \pm 0.43$  days) than those fed on the other host plants (Table 1). Also, significant differences in the length of development time and total life span of female individuals of *H. armigera* were found on these hosts. The shortest ( $35.21 \pm 0.35$  days) and longest ( $42.72 \pm 0.28$  days) development time (egg to adult emergence) of *H. armigera* was observed on chickpea and corn, respectively.

**Table 1** Duration (mean  $\pm$  SE) of different life stages of *Helicoverpa armigera* on five different host plants.

| Host plant | Incubation period (day) |                    | Larval period (day) | Pre-pupal period (day) | Pupal period (day) | Development time (day) |                   | Longevity (day)   |      | Total lifespan (day) |      |
|------------|-------------------------|--------------------|---------------------|------------------------|--------------------|------------------------|-------------------|-------------------|------|----------------------|------|
|            |                         |                    |                     |                        |                    |                        |                   | Female            | Male | Female               | Male |
| Cowpea     | 3.09 $\pm$ 0.08a        | 17.20 $\pm$ 0.28c  | 3.02 $\pm$ 0.07b    | 13.90 $\pm$ 0.14b      | 37.07 $\pm$ 0.24c  | 12.56 $\pm$ 0.99b      | 11.61 $\pm$ 0.93a | 49.33 $\pm$ 0.82b |      | 48.56 $\pm$ 0.91a    |      |
| Soybean    | 2.92 $\pm$ 0.05a        | 18.31 $\pm$ 0.30b  | 3.03 $\pm$ 0.06b    | 14.40 $\pm$ 0.13a      | 38.62 $\pm$ 0.24b  | 11.09 $\pm$ 0.67b      | 10.34 $\pm$ 0.82a | 49.38 $\pm$ 0.76b |      | 49.31 $\pm$ 0.76a    |      |
| Corn       | 3.06 $\pm$ 0.08a        | 23.52 $\pm$ 0.43a  | 3.93 $\pm$ 0.11a    | 13.90 $\pm$ 0.31b      | 42.72 $\pm$ 0.28a  | 11.88 $\pm$ 0.59b      | 9.67 $\pm$ 1.15a  | 52.50 $\pm$ 1.10a |      | 51.67 $\pm$ 0.68a    |      |
| Navy bean  | 3.00 $\pm$ 0.06a        | 17.95 $\pm$ 0.39bc | 2.94 $\pm$ 0.05b    | 13.68 $\pm$ 0.12b      | 37.90 $\pm$ 0.29bc | 12.70 $\pm$ 0.48b      | 11.92 $\pm$ 0.90a | 49.41 $\pm$ 0.62b |      | 49.54 $\pm$ 0.95a    |      |
| Chickpea   | 2.97 $\pm$ 0.06a        | 15.92 $\pm$ 0.22d  | 3.05 $\pm$ 0.12b    | 13.61 $\pm$ 0.14b      | 35.21 $\pm$ 0.35d  | 15.11 $\pm$ 0.46a      | 11.95 $\pm$ 0.68a | 49.88 $\pm$ 0.52b |      | 48.11 $\pm$ 1.06a    |      |
| F          | 0.32                    | 36.60              | 6.99                | 2.45                   | 36.55              | 6.56                   | 0.82              | 2.66              |      | 1.11                 |      |
| df         | 4, 150                  | 4, 150             | 4, 149              | 4, 149                 | 4, 149             | 4, 130                 | 4, 108            | 4, 141            |      | 4, 110               |      |
| P          | 0.868                   | < 0.001            | < 0.001             | 0.049                  | < 0.001            | < 0.001                | 0.516             | 0.035             |      | 0.356                |      |

The means in a column followed by the same letters are not significantly different ( $P < 0.05$ , Tukey's test).

**Adult longevity and fecundity**

The effect of five different diets on the adult longevity and reproduction of *H. armigera* are given in Table 1 and 2. The female adult longevity was significantly different on the diets examined (Table 1). The longest female adult longevity ( $15.11 \pm 0.46$  d) was found in the larvae feeding on chickpea. However, no significant difference in the longevity of male adults was found on the five diets tested.

Significant difference was found among five diets in terms of the total life span of female

individuals (Table 1). The longest ( $52.50 \pm 1.10$  days) and shortest ( $49.33 \pm 0.82$  days) total life span of females were observed on corn and cowpea, respectively. However, no significant differences in the total life span were found among male individuals. The oviposition period and fecundity of *H. armigera* are summarized in Table 2. The females reared on cowpea had the highest daily fecundity ( $93.1 \pm 9.32$  eggs) (Table 2) and total fecundity ( $632.82 \pm 50.75$  eggs) (Table 2).

**Table 2** Pre- and post-oviposition and oviposition periods and fecundity of *Helicoverpa armigera* emerging from the larvae reared on five different host plants.

| Host plant | Pre-oviposition period (day) | Oviposition period (day) | Post-oviposition period (day) | Fecundity (Egg/Female) |                      |
|------------|------------------------------|--------------------------|-------------------------------|------------------------|----------------------|
|            |                              |                          |                               | Daily Total            |                      |
| Cowpea     | $2.86 \pm 0.15c$             | $6.33 \pm 0.51b$         | $5.38 \pm 0.61a$              | $93.11 \pm 9.32a$      | $632.82 \pm 50.75a$  |
| Soybean    | $3.06 \pm 0.23bc$            | $6.33 \pm 0.37b$         | $3.93 \pm 0.49bc$             | $71.84 \pm 6.34b$      | $560.25 \pm 47.75ab$ |
| Corn       | $3.80 \pm 0.26a$             | $5.47 \pm 0.47b$         | $3.38 \pm 0.44c$              | $55.74 \pm 6.20b$      | $352.00 \pm 26.09c$  |
| Navy bean  | $3.42 \pm 0.17ab$            | $6.04 \pm 0.30b$         | $3.69 \pm 0.35bc$             | $69.00 \pm 5.24b$      | $466.24 \pm 36.61bc$ |
| Chickpea   | $2.68 \pm 0.10c$             | $7.86 \pm 0.40a$         | $4.92 \pm 0.35ab$             | $75.67 \pm 1.55ab$     | $590.58 \pm 39.73ab$ |
| F          | 5.96                         | 5.02                     | 3.41                          | 3.92                   | 5.64                 |
| df         | 4, 103                       | 4, 104                   | 4, 96                         | 4, 98                  | 4, 89                |
| P          | < 0.001                      | 0.001                    | 0.012                         | 0.005                  | < 0.001              |

The means in a column followed by the same letters are not significantly different ( $P < 0.05$ , Tukey's test).

**Survivorship and life expectancy**

Age-specific survivorship ( $l_x$ ) at the age of adult emergence of *H. armigera* on cowpea, soybean, corn, navy bean and chickpea was 0.86, 0.87, 0.59, 0.85 and 0.85, respectively (Fig. 1). Life expectancy ( $e_x$ ) at the first day of the adult emergence on the above-mentioned diets was 10.88, 10.15, 9.55, 10.28 and 11.18 days, respectively (Fig. 2). At the beginning of life, the longest and shortest life expectancy was recorded on soybean (43.07 days) and corn (37.82 days), respectively. Maximum age (the death of the last female) was observed on corn (58 days).

**Reproductive parameters**

The values of the reproductive parameters of *H. armigera* on five diets are presented in Table 3. The highest gross fecundity rate was observed on cowpea (778.02 eggs) and the lowest one


was on corn (417.33 eggs) (Table 3). The gross fertility also showed significant differences among five different host plants tested. The highest value of the gross fertility was observed on chickpea (850.86 eggs). The lowest (135.78 eggs) and highest (933.59 eggs) net fecundity rates were observed on corn and chickpea (Table 3), respectively. The net fertility rate ranged from 681.52 to 50.24 eggs on chickpea and corn, respectively (Table 3).

**Life table parameters**


The life table parameters of *H. armigera* on the different diets are shown in Table 4. The net reproductive rate ( $R_0$ ) was significantly different on different host plants tested (Table 4). The lowest  $R_0$  was observed on corn ( $147.40 \pm 6.69$  female offspring/ female) and the highest one was on cowpea ( $365.66 \pm 24.52$  female offspring/ female). Furthermore,

the intrinsic rate of increase ( $r_m$ ) and finite rate of increase ( $\lambda$ ) of *H. armigera* were significantly different on the five diets (Table 4). Among five different host plants, the highest value of  $r_m$  ( $0.180 \pm 0.003 \text{ d}^{-1}$ ) and  $\lambda$  ( $1.192 \pm 0.004 \text{ d}^{-1}$ ) were estimated on cowpea and the lowest value of these two parameters were observed on corn ( $0.126 \pm 0.002 \text{ d}^{-1}$  and  $1.131 \pm 0.004 \text{ d}^{-1}$ , respectively). The results

revealed significant differences among five diets in terms of the mean generation time ( $T$ ) and doubling time ( $DT$ ) (Table 4). The longest mean generation time ( $T$ ) and doubling time ( $DT$ ) were observed on corn ( $37.90 \pm 0.26 \text{ d}$  and  $5.62 \pm 0.17 \text{ d}$ , respectively) and the shortest ones were obtained on cowpea ( $31.62 \pm 0.22 \text{ d}$  and  $3.92 \pm 0.08 \text{ d}$ , respectively).



**Figure 1** Survival rate ( $l_x$ ) of *Helicoverpa armigera* reared on five different host plants.



**Figure 2** Life expectancy ( $e_x$ ) of *Helicoverpa armigera* reared on five different host plants.

**Table 3** The mean values of reproduction parameters of *Helicoverpa armigera* on five different host plants.

| Host plant | Gross fecundity rate (Eggs per female) | Gross fertility rate (Eggs per female) | Net fecundity rate (Eggs per female) | Net fertility rate (Eggs per female) |
|------------|--|--|--------------------------------------|--------------------------------------|
| Cowpea     | 778.02                                 | 544.62                                 | 644.38                               | 451.07                               |
| Soybean    | 1074.57                                | 848.91                                 | 843.34                               | 666.24                               |
| Corn       | 417.33                                 | 154.41                                 | 135.78                               | 50.24                                |
| Navy bean  | 1173.64                                | 809.81                                 | 862.25                               | 594.95                               |
| Chickpea   | 1165.57                                | 850.86                                 | 933.59                               | 681.52                               |

**Table 4** The mean ( $\pm$  SE) population parameters of *Helicoverpa armigera* on five different host plants.

| Host plant | $R_0$ (female offspring per female) | $r_m$ (day <sup>-1</sup> ) | $\lambda$ (day <sup>-1</sup> ) | $T$ (day)          | $DT$ (day)       |
|------------|-------------------------------------|----------------------------|--------------------------------|--------------------|------------------|
| Cowpea     | 365.66 $\pm$ 24.52a                 | 0.180 $\pm$ 0.003a         | 1.192 $\pm$ 0.004a             | 31.62 $\pm$ 0.22c  | 3.92 $\pm$ 0.08b |
| Soybean    | 239.69 $\pm$ 14.74c                 | 0.161 $\pm$ 0.004b         | 1.177 $\pm$ 0.004ab            | 33.28 $\pm$ 0.19b  | 4.23 $\pm$ 0.10b |
| Corn       | 147.40 $\pm$ 6.69d                  | 0.126 $\pm$ 0.002c         | 1.131 $\pm$ 0.004c             | 37.90 $\pm$ 0.26a  | 5.62 $\pm$ 0.17a |
| Navy bean  | 294.28 $\pm$ 11.52b                 | 0.164 $\pm$ 0.005b         | 1.178 $\pm$ 0.005ab            | 32.31 $\pm$ 0.71bc | 4.14 $\pm$ 0.16b |
| Chickpea   | 359.67 $\pm$ 15.52a                 | 0.161 $\pm$ 0.004b         | 1.175 $\pm$ 0.004b             | 33.28 $\pm$ 0.19b  | 4.27 $\pm$ 0.11b |
| F          | 22.01                               | 15.35                      | 17.83                          | 28.49              | 21.54            |
| df         | 4, 71                               | 4, 94                      | 4, 103                         | 4, 103             | 4, 103           |
| P          | < 0.001                             | < 0.001                    | < 0.001                        | < 0.001            | < 0.001          |

Means in a column followed by the same letter(s) are not significantly different ( $P < 0.05$ , Tukey's test).

## Discussion

Host plants and diets taken from them may be different in terms of their nutritive properties and this difference may influence survival, development and reproductive rates of herbivorous insects feeding on them. Herbivorous insect pests usually have shorter development time and greater total reproduction on susceptible host plants in comparison with resistant ones. Such host plants can be rapidly colonized by insect pests (van Lenteren and Noldus, 1990). Although, developmental rates and reproduction provide clues concerning the ability of a host to support an insect to complete life cycle, these data should be linked to other parameters to show host suitability (Liu *et al.*, 2004). Results of the present study showed that the performance of the cotton bollworm differed significantly on the five host plants studied. It provided valuable information on the effect of different host plants on the performance of *H. armigera* which can be utilized properly in integrated pest management programs.

We found that the larvae reared on corn-based artificial diet moulted five times during their development time and had six larval instars similar to that reported previously by Ali *et al.* (2009) and Soleimannejad *et al.* (2010). Existence of different number of larval instars for the cotton bollworm shows that the number of larval instars in *H. armigera* can be affected by the quality of host plants. The results also revealed that the egg incubation period was not affected by host plant quality and *H. armigera*

had the same incubation period on different host plants. In contrast with the incubation period, length of the larval stage and development time varied significantly on the different host plants and the longest larval period was observed on corn. This may stem from poor nutritional quality or existence of high amount of the secondary metabolites in this cultivar which makes it less suitable to be fed properly by *H. armigera*. Short life cycle, high reproductive potential and rapid population growth indicate suitability of a host plant to be colonized by an insect pest (Singh and Parihar, 1988). This would be reflected ultimately in the population size. These findings are in agreement with our previous outcomes (Bagheri, 2011) in which we showed that the *H. armigera* feeding on corn had the lowest values of efficiency of conversion of ingested food (ECI) and efficiency of conversion of digested food (ECD). The value of ECD and ECI show the activity of digestive enzymes in an insect to utilize properly ingested food (Lazarevic *et al.*, 2004). Plants can synthesize the secondary substances that are involved in blocking of insects' digestive enzymes and can increase the resistance of host plants to insect pests (Chougule *et al.*, 2003). It seems that corn contains some enzyme inhibitors which can slow down the activity of digestive enzymes of *H. armigera* and prolong its developmental time.

Difference in the longevity of the cotton bollworm feeding on the five different host plants might have resulted due to the difference in the feeding rate or difference in the feeding efficiency of this pest in the larval stage on

these host plants. When food quality decreases only a limited number of females are able to survive. The insufficient feeding in the larval stages may cause the number of eggs in the adult stage to decrease. This may explain the lower fecundity of *H. armigera* on the corn host plant compared with the others.

Our research demonstrated significant differences in the life table parameters of *H. armigera* on five different host plants studied. The lowest value of the net reproduction rate ( $147.40 \pm 6.69$  female offspring) was obtained on corn which was greater than that documented by Liu *et al.* (2004) on corn (44.5 female offspring) and similar to that reported by Reddy *et al.* (2004) on sunflower (143.77 female offspring). At the same time, Dabhi and Patel (2007) showed that the  $R_0$  value of *H. armigera* on chickpea was 361.84 female offspring, which is close to our findings on chickpea in the present study, indicating the values of  $R_0$  estimated in the current study were similar to those reported by other researchers (Dabhi and Patel, 2007). The net reproduction rate is an important parameter in population dynamics (Richard, 1961). Since this parameter indicates the physiological ability of an insect in relation to its reproductive capacity (Fathipour and Naseri, 2011), its comparison often provides considerable insight beyond that available from the independent analysis of individual life history parameters.

The intrinsic rate of increase is a key factor in the life table, influenced by different factors such as fecundity, survival and especially generation time. Furthermore, it adequately summarizes the physiological qualities of an insect to increase its population. Therefore, it would be a most appropriate index to evaluate the performance of an insect on different host plants or in other words it can show level of the resistance in different host plants to insect pests. Higher value of  $r_m$  on cowpea indicates that the cotton bollworm has a higher reproductive potential on this host plant. Similarly, lower value of  $r_m$  on corn can be attributed to its lower fecundity and longer development time on this host plant.

In cluster analysis, the grouping of different host plants within the same cluster might be due to high correspondence of physiological traits of these host plants. In contrast, the separate clusters might represent significant variability in host plant suitability. By comparing the demographic parameters of *H. armigera* on five different host plants we found that the cluster A embraced highly susceptible host plants (cowpea and chickpea) whereas clusters B and C contained moderate (soybean and navy bean) and resistant (corn) host plants, respectively. The corn host plant grouped in cluster C and diverged from the others. It was the most resistant host plant to *H. armigera* because it retarded the development time of *H. armigera* and decreased its fecundity.

Our results confirmed the lower nutritional attribute of the corn host plant compared with the other species. Insufficient amount of primary metabolites necessary for growth and development of the cotton bollworm renders this host plant improper for *H. armigera*. Also, this plant may contain some phytochemicals that act as antixenosis and antibiosis agents, resulting in poor colonization of the cotton bollworm on this host plant. The antibiosis effects of this host plant may have led to the reduction in survival fitness of *H. armigera* and prolonged its development time. The increased development time of an insect pest can increase its exposure and predation by natural enemies. Understanding the extent of resistance in different host plants, population growth potential and biology of an insect pest on a host plant are important tools used in the integrated pest management (IPM) programs. It is known that the kind and amount of nutrients can affect the survival and reproduction of insect pests through targeting the capability of digestion and assimilation. Therefore, knowing how the quality of host plants influence the demographic parameters of *H. armigera* can help to increase our knowledge regarding its population dynamics and management. Our findings provided insights into the life cycle of *H. armigera* and its potential on different host plants. This information in combination with



other semi-field and field experiments can be useful in development and implementation of management programs against *H. armigera*.

### Acknowledgments

This research was supported by the Department of Entomology, Tarbiat Modares University, which is greatly appreciated.

### References

- Ali, A., Choudhury, R. A., Ahmad, Z., Rahman, F., Khan, F. R. and Ahmad, S. K. 2009. Some biological characteristics of *Helicoverpa armigera* on chickpea. *Tunisian Journal of Plant Protection*, 4: 99-106.
- Bagheri, F. 2011. Effect of seeds of different host plants on demographic parameters, nutritional indices and digestive enzymes of *Helicoverpa armigera* (Lepidoptera: Noctuidae). M. Sc. Dissertation, Tarbiat Modares University, Tehran. 102 pp.
- Carey, J. R. 1993. *Applied Demography for Biologists with Special Emphasis on Insects*. Oxford University Press, New York.
- Chougule, N. P., Hivrae, V. K., Chhabda, P. J., Giri, A. P. and Kachole, M. S. 2003. Differential inhibition of *Helicoverpa armigera* gut proteinases by proteinase inhibitors of pigeon pea (*Cajanus cajan*) and its wild relatives. *Phytochemistry*, 64(3): 681-687.
- Dabhi, M. V. and Patel, C. C. 2007. Life expectancy of *Helicoverpa armigera* on chickpea. *Journal of SAT Agricultural Research*, 5(1): 1-2.
- Fathipour, Y. and Naseri, B. 2011. Soybean cultivars affecting performance of *Helicoverpa armigera* (Lepidoptera: Noctuidae). In: Ng, T. B. (Ed.), *Soybean - Biochemistry, Chemistry and Physiology*. InTech, Rijeka, Croatia, pp. 599-630. Available from: <http://www.intechopen.com/articles/show/title/soybean-cultivars-affecting-performance-of-helicoverpa-armigera-lepidoptera-noctuidae>.
- Fathipour, Y. and Sedaratian, A. 2013. Integrated management of *Helicoverpa armigera* in soybean cropping systems. In: El-Shemy, H. (Ed.), *Soybean-Pest Resistance*. InTech, Rijeka, Croatia, pp. 231-280.
- Fathipour, Y., Sedaratian, A., Bagheri, A. and Talaei-Hassanlouei, R. 2019. Increased food utilization indices and decreased proteolytic activity in *Helicoverpa armigera* larvae fed sublethal *Bacillus thuringiensis*-treated diet. *Physiological Entomology*, 44(3-4): 178-186.
- Frei, A., Gu, H., Bueno, J. M., Cardona, C. and Dorn, S. 2003. Antixenosis and antibiosis of common beans to *Thrips palmi* Karny (Thysanoptera: Thripidae). *Journal of Economic Entomology*, 96(5): 1577-1584.
- Garcia, F. J. M. 2006. Analysis of the spatio-temporal distribution of *Helicoverpa armigera* Hb. in a tomato fields using a stochastic approach. *Biosystems Engineering*, 93(3): 253-259.
- Golizadeh, A., Kamali, K., Fathipour, Y. and Abbasipour, H. 2009. Life table of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) on five cultivated brassicaceous host plants. *Journal of Agricultural Science and Technology*, 11: 115-124.
- Jafari, K., Fathipour, Y., Bagheri, A. and Talebi, A. A. 2020. Tritrophic interactions in a wheat (*Triticum aestivum*), aphid (*Rhopalosiphum padi*) and parasitoid (*Aphidius matricariae*) system. *Crop Protection*, p.105076.
- Karimi, S., Fathipour, Y., Talebi, A. A. and Naseri, B. 2012. Evaluation of canola cultivars for resistance to *Helicoverpa armigera* (Lepidoptera: Noctuidae) using demographic parameters. *Journal of Economic Entomology*, 105(6): 2172-2179.
- Lazarevic, J., Peric-mataruga, V., Vlahovic, M., Mrdakovic, M. and Cvetanovic, D. 2004. Effects of rearing density on larval growth and activity of digestive enzymes in *Lymantria dispar* L. (Lepidoptera: Lymantriidae). *Folia Biologica*, 52: 105-112.
- Liu, Z., Li, D., Gong, P. and Wu, K. 2004. Life table studies of the cotton bollworm,

- Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae), on different host plants. Environmental Entomology, 33(6): 1570-1576.
- Maia, A. H. N., Luiz, A. J. B. and Campanhola, C. 2000. Statistical inference on associated fertility life table parameters using jackknife technique: computational aspects. Journal of Economic Entomology, 93(2): 511-518.
- Meyer, J. S., Ingersoll, C. G., McDonald, L. L. and Boyce, M. S. 1986. Estimating uncertainty in population growth rates: jackknife vs. bootstrap techniques. Ecology, 67(5): 1156-1166.
- Naseri, B., Fathipour, Y., Moharramipour, S. and Hosseiniaveh, V. 2009a. Comparative life history and fecundity of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on different soybean varieties. Entomological Science, 12: 51-58.
- Naseri, B., Fathipour, Y., Moharramipour, S. and Hosseiniaveh, V. 2009b. Life table parameters of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae) on different soybean cultivars. Journal of Entomological Society of Iran, 29: 25-40.
- Naseri, B., Fathipour, Y., Moharramipour, S. and Hosseiniaveh, V. 2011. Comparative reproductive performance of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) reared on thirteen soybean varieties. Journal of Agricultural Science and Technology, 13: 17-26.
- Reddy, K. S., Rao, G. R., Rao, P. A. and Rajasekhar, P. 2004. Life table studies of the capitulum borer, *Helicoverpa armigera* (Hübner) infesting sunflower. Journal of Entomological Research, 28(1): 13-18.
- Richard, O. W. 1961. The theoretical and practical study of natural insect populations. Annual Review of Entomology, 6: 147-162.
- Sedaratian, A., Fathipour, Y. and Talaei-Hassanlou, R. 2014. Deleterious effects of *Bacillus thuringiensis* on biological parameters of *Habrobracon hebetor* parasitizing *Helicoverpa armigera*. BioControl, 59(1): 89-98.
- Sedaratian, A., Fathipour, Y., Talaei-Hassanlou, R. and Jurat-Fuentes, J. L. 2013. Fitness costs of sublethal exposure to *Bacillus thuringiensis* in *Helicoverpa armigera*: a carryover study on offspring. Journal of Applied Entomology, 137(7): 540-549.
- Singh, O. P. and Parihar, S. B. B. 1988. Effect of different hosts on the development of *Heliothis armigera* Hub. Bulletin of Entomological Research, 29(2): 168-172.
- Soleimannejad, S., Fathipour, Y., Moharramipour, S. and Zalucki, M. P. 2010. Evaluation of potential resistance in seeds of different soybean cultivars to *Helicoverpa armigera* (Lepidoptera: Noctuidae) using demographic parameters and nutritional indices. Journal of Economic Entomology, 103(4): 1420-1430.
- Soufbaf, M., Fathipour, Y., Zalucki, M. P. and Hui, C. 2012. Importance of primary metabolites in canola in mediating interactions between a specialist leaf-feeding insect and its specialist solitary endoparasitoid. Arthropod-Plant Interactions, 6: 241-250.
- Soufbaf, M., Fathipour, Y., Karimzadeh, J. and Zalucki, M. 2010a. Bottom-up effect of different host plants on *Plutella xylostella* (Lepidoptera: Plutellidae): A life-table study on canola. Journal of Economic Entomology, 103: 2019-2027.
- Soufbaf, M., Fathipour, Y., Karimzadeh, J. and Zalucki, M. 2010b. Development and age-specific mortality of diamondback moth on *Brassica* host plants: pattern and causes of mortality under laboratory conditions. Annals of the Entomological Society of America, 103: 574-579.
- Teakle, R. E. 1991. Laboratory culture of *Heliothis* species and identification of disease. In: Zalucki, M. P. (Ed.), *Heliothis: Research Methods and Prospects*. Springer Verlag, pp. 22-29.
- Van Lenteren, J. C. and Noldus, L. P. J. J. 1990. Whitefly-plant relationship: behavioral and biological aspects. In: Gerling, D. (Ed.), *Whitefly: Their Bionomics, Pest Status and Management*. Intercept, Andover, U. K, pp. 47-89.

## رشدونمو، تولیدمثل و پارامترهای جدول زندگی *Helicoverpa armigera* (Lepidoptera: Noctuidae) روی بذر پنج میزبان مختلف گیاهی

یعقوب فتحی پور<sup>۱\*</sup>، فاطمه باقری<sup>۱</sup>، عبدالنبی باقری<sup>۲</sup> و بهرام ناصری<sup>۳</sup>

- ۱- گروه حشره شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران.
  - ۲- بخش تحقیقات گیاه پزشکی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی هرمزگان، سازمان تحقیقات، آموزش و ترویج کشاورزی، بندرعباس، ایران.
  - ۳- گروه گیاه پزشکی، دانشکده گیاه پزشکی، دانشگاه محقق اردبیلی، اردبیل، ایران.
- پست الکترونیکی نویسنده مسئول مکاتبه: fathi@modares.ac.ir  
دریافت: ۱۰ آذر ۱۳۹۸؛ پذیرش: ۳۰ تیر ۱۳۹۹

**چکیده:** کرم غوزه پنبه (*Helicoverpa armigera* (Hübner) یکی از آفات مهم اقتصادی بسیاری از محصولات کشاورزی در بسیاری از مناطق دنیا است. در این تحقیق، پارامترهای دموگرافی این آفت روی بذر پنج میزبان گیاهی شامل لوبیا چشم بلبلی (رقم مشهد)، لوبیا سفید (رقم دهقان)، نخود (رقم هاشم)، سویا (رقم 033) و ذرت (رقم سینگل کراس 704) در شرایط آزمایشگاهی، دمای  $25 \pm 1$  درجه سلسیوس، رطوبت نسبی  $65 \pm 5$  درصد و دوره نوری ۱۶ ساعت روشنایی و ۸ ساعت تاریکی تعیین شد. کوتاه ترین طول دوره رشدی قبل از بلوغ ۳۶/۱۶ و ۳۴/۹۸ روز به ترتیب برای ماده و نر روی نخود بود و طولانی ترین مدت زمان این دوره برای نر و ماده روی ذرت بود (به ترتیب ۴۲/۰۰ و ۴۲/۹۵ روز). بیشترین تخم ریزی روزانه و کل روی لوبیا چشم بلبلی و کمترین آن روی ذرت مشاهده شد. براساس این نتایج، نرخ ذاتی افزایش جمعیت ( $r_m$ ) روی میزبان های ذکر شده در بالا به ترتیب ۰/۱۸۰، ۰/۱۴۴، ۰/۱۶۱، ۰/۱۶۱ و ۰/۱۲۶ در روز و نرخ خالص تولیدمثل ( $R_0$ ) روی این میزبان ها به ترتیب ۳۶۵/۶۶، ۲۹۴/۲۸، ۳۶۵/۶۷، ۳۳۹/۶۹ و ۱۴۷/۴۰ ماده/ماده بود. طولانی ترین متوسط مدت زمان یک نسل ( $T$ ) و مدت زمان دو برابر شدن جمعیت ( $DT$ ) این آفت به ترتیب ۳۷/۹۰ و ۵/۶۲ روز روی ذرت بود. این نتایج نشان داد در بین میزبان های گیاهی در این تحقیق لوبیا چشم بلبلی و ذرت به ترتیب حساس ترین و مقاوم ترین میزبان ها برای رشدونمو *H. armigera* بودند.

**واژگان کلیدی:** کرم غوزه پنبه، *Helicoverpa armigera*، باروری، ذرت (cv. Single cross 704)، لوبیا چشم بلبلی (cv. Mashhad)