

# Morphological and molecular identification of *Fusarium* spp. associated with carnation *Dianthus caryophyllus* in Mahallat, Iran

Samira Shahbazi<sup>1</sup>, Fatemeh Zaker Tavallaie<sup>2\*</sup> and Zoha Daroodi<sup>3</sup>

1. Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute, Alborz, Iran.

2. Department of Plant Production, Faculty of Agriculture, Higher Education Complex of Shirvan, Shirvan, Iran.

3. Department of Plant Protection, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran.

Abstract: Carnation is an ornamental plant of economic importance worldwide. Fusarium fungus is one of the critical infectious agents of Carnation in Mahallat city. Some species of Fusarium cause wilting and crown and root rot in carnation. In this study, 20 fungal isolates of Fusarium were isolated from diseased carnation plants and rhizosphere samples from Carnation greenhouses in Mahallat. The isolates were identified morphologically using Leslie and Summerell's identification key. Amplification of the *TEF-1* $\alpha$  gene region was done using ef1 and ef2 primers. Amplified fragments were purified and sequenced, and the nucleotide sequences were introduced in the Fusarium ID database for molecular identification of isolates based on the *TEF-1* $\alpha$  gene. A phylogenetic tree was also drawn. The results showed that Fusarium isolates were categorized into five species: F14 was identified as Fusarium brachygibbosum morphologically and F. solani molecularly. F6 was morphologically and molecularly identified as F. solani, F10 identified as Fusarium culmorum. All crown and root isolates (F11, F12, F13, F15, F16, F17, F18, F9, and F20) were identified as F. oxysporum. The rhizosphere isolates of F1, F2, F3, F4, F5, F7, F8, and F9 were identified as Fusarium proliferatum. This study shows that the use of the TEF-1 $\alpha$  gene for molecular identification of Fusarium isolates is a relatively accurate, fast, and straightforward method for detecting species and can be used in plant pathological studies. Identifying the pathogen is the first step to controlling the disease.

Keywords: Carnation, *Fusarium*, molecular identification, *TEF-1* $\alpha$  gene

#### Introduction

Mahallat city in Markazi province is an important center for producing flowers and ornamental plants in Iran. Carnation *Dianthus caryophyllus* L. belongs to the family Caryophyllaceae and is one of Iran's four central cut flowers for export. The genus Dianthus includes more than 300 species,

several of which are sold as ornamental garden plants, pots, or cut branches. *Fusarium* species are economically important as pathogens on most crops in the world. Also, some of the *Fusarium* species are known as saprophytes. Diseases caused by *Fusarium* spp. in carnations include wilting, root, and crown rot that cause damage to carnation greenhouses every year. *Fusarium* wilt is one of the most destructive diseases in most carnation-growing regions of the world (Kermajany *et al.*, 2017). This disease was first reported in Iran in 1996 by Varamin (Etebarian, 1996). Fattahi *et al.* (2013) identified the *F.oxysporum*, *F. proliferatum*, *F.* 

Handling Editor: Naser Safaie

<sup>\*</sup> Corresponding author: zakertavallaie@cheshirvan.ac.ir Received: 19 February 2021, Accepted: 05 June 2021 Published online: 05 July 2021

J. Crop Prot.

*solani*, and *F. equiseti* in carnation greenhouses in Iran and reported wilt for the first time disease caused by *F. proliferatum* and *F. solani* on carnation in Iran.

Accurate identification of *Fusarium* species in carnation greenhouses is the first step in combating diseases caused by this pathogenic fungus. Morphological specification of fungi is based on the size, shape, absence, or presence of asexual reproduction structures in different culture media (Leslie and Summerell, 2006). Molecular identification can be made using DNA-based procedures, such as conserved sequencing regions. The TEF-1 $\alpha$  gene encoding translation elongation factor  $1\alpha$  (TEF-1 $\alpha$ ) is suitable in sequence analysis for taxonomic and phylogenetic studies (Geiser et al., 2004). It appears that this gene in Fusarium is consistently single-copy. It shows a high level of sequence polymorphism among closely related species. That is why TEF-1 $\alpha$  is a marker of choice as a single-locus recognition tool in Fusarium (Geiser et al., 2004).

study aimed This to identify the morphological and molecular characteristics of Fusarium species associated with carnation in Mahallat. According to our study, molecular identification of carnation fusarium has not been performed in Iran so far. So, carnation plants with disease and infected rhizosphere symptoms were collected from infected greenhouses in Mahallat city in Markazi province. After culturing the samples and purifying the fungus, morphological identification was performed using the Leslie and Summerell identification key (Leslie and Summerell, 2006). Molecular identification was also based on the TEF-1 $\alpha$  gene. (Geiser *et al.*, 2004).

#### **Materials and Methods**

## Sample collection and isolation of *Fusarium* species

The *Fusarium* Isolates were isolated from the diseased carnation plants and their rhizosphere in Mahalat during the growing season in 2019. From among 20 *Fusarium* isolates obtained

from Carnation grown in greenhouses, isolates F1 to F10 were obtained from the rhizosphere, and isolated F11 to F20 were isolated from the crown and roots of infected plants.

The infected plant tissues were surface sterilized with ethanol 80 % and transferred onto potato dextrose agar medium (Peeled potato; 250 g, Dextrose; 20 g, Distilled water1000 mL at pH 7.0) in Petri dishes. The Petri dishes were incubated at room temperature  $(28 \pm 2 \text{ °C})$  observed periodically to grow colonies. The pure colonies were obtained by the single spore method. The pure colonies were transferred to the PDA Petri plate to maintain the culture.

#### Morphological identification

The Carnation leaf culture medium (CLA and KCL-agar) was used to identify Fusarium species. Morphological identification of isolates was made based on the morphological characteristics of macroconidia, the presence or absence of microconidia. If so, their shape and position at the end of conidiophore (single, chain, and false heads), type of phialides (monophialide or polyphialide), presence or absence of chlamydospores, as well as colony characteristics were determined on potato dextrose agar (PDA) medium, including mycelium pigmentation and rate of colony growth using Leslie and Summerell identification key (Leslie and Summerell, 2006). The pathogenicity test was performed according to the method described by Moeini et al. (2014).

#### Molecular identification of *Fusarium* species

DNA extraction was performed from fungal mycelium using the extraction kit and instructions of Pars Toos Company. DNA concentration was estimated using a UV-vis spectrophotometer (Thermo Spectronic UV1). It was then stored at 20 °C. A standard polymerase chain reaction (PCR) was used to amplify the TEF-1 $\alpha$  gene region using ef1 (5-ATGGGTAAGGA(A/G)GACAAGAC-3) as the forward primer and ef2 (5-GGA(G/A)GTACCAGT(G/C)ATCATGTT-3)

as a reverse primer (Geiser *et al.*, 2004). These primers amplify a 700 bp region of TEF-1 $\alpha$  in all known fusaria.

The PCR was performed in ThermoCycler (BIORAD® T100TM, Germany) as the following program: Initial denaturation at 94 °C for 5 min followed by 30 cycles at 94 °C for 1 min, 55 °C for 50 s, and 72 °C for 3 min and a final extension at 72 C for 10 min. Amplified PCR products were separated on an agarose gel (1.5% w/v) in 1X TAE buffer at 100 V. The 100bp DNA marker (MBI Fermentas. Lithuania) was used to estimate DNA band size. Negative control was used to test the absence of contamination.

#### **DNA Sequencing**

Isolation of 700 bp band of DNA was done under UV. Separated bands were purified and sequenced by the South Korean Macrogene Company (http://dna.macrogen.com, Korea). The sequences for TEF-1 $\alpha$  areas were first reviewed with the BioEdit software, and then sequences were presented in the Gene Bank database (www.ncbi.nlm.nih.gov/BLAST). The sequences were registered on the GenBank of (https://www.ncbi.nlm.nih.gov/ NCBI site WebSub/). The nucleotide sequences also were introduced in the Fusarium ID database (http://isolate.Fusariumdb.org/blast.php) for molecular identification of isolates based on translation elongation factor 1 alpha (TEF-1a gene). Phylogenetic analysis was done using the ClustalW (https://www.genome.jp/tools-bin/ clustalw) program.

#### **Phylogenetic analysis**

The newly obtained elongation factor 1-alpha (EF1- $\alpha$ ) sequences of *Fusarium* spp. (from F1 to F20), together with the other sequences of the *Fusarium* genus already used by Laurence *et al.* (2011), Herron *et al.* (2015), Papizadeh *et al.* (2018), and Al-Hatmi *et al.* (2018) were selected for phylogenetic analyses. The dataset was updated by investigations in the database for acquiring correct sequences. The outgroup taxa for the present dataset were taken based on Hirooka *et al.* (2011). All sequences were

aligned using the Q-INS-i algorithm of MAFFT version 7 (http://mafft.cbrc.jp/ alignment/ server/) (Katoh and Standley, 2013). The online version of Gblocks 0.91b (Castresana, 2000) was applied to remove ambiguous parts of the alignment, with all three options (including allow smaller final blocks, allow gap positions within the final blocks, and allow less strict flanking position) for a fewer stringent selection (http://molevol.cmima.csic.es/ castresana/Gblocks server.html). The most suitable substitution model for the dataset was chosen using the Akaike information criterion (AIC) using PAUP\* /MrModeltest v2.2 (Nylander, 2004). Asymmetrical model including a gamma distribution for rates (GTR + G) was picked out for EF1- $\alpha$  analysis. Bayesian inference (BI) was carried out using MrBayes v3.1.2 (Ronquist and Huelsenbeck, 2003), choosing a random beginning tree and running the chains for 4 million for EF1- $\alpha$ . After casting off burn-in samples, the residual samples were reserved for additional analyses. The Markov Chain Monte Carlo (MCMC) method within a Bayesian framework was utilized to assess the posterior probabilities of the phylogenetic trees (Larget and Simon, 1999) by the 50% majority rule. The obtained phylogenetic tree was visualized via Dendroscope V.3.2.8 (Huson and Scornavacca, 2012).

#### Results

### Morphological identification of *Fusarium* isolates

Fusarium isolates were identified morphologically. Based on structures of macroconidia. microconidia, and other morphological characters (Fig. 1), the isolates of F1, F2, F3, F4, F5, F7, F8, and F9 were identified as F. proliferatum. The isolates of F11, F12, F13, F15, F16, F17, F18, F9, and F20 were identified as F. oxysporum. Also, the isolates F6, F10, and F14 as F. solani, F. brachygibbosum, culmorum, and F. respectively. All F. oxysporum isolates and F. brachygibbosum were isolated from the roots

and crowns of wilted plants. At the same time, *F. proliferatum* and *F. culmorum* were isolated from the rhizosphere. Moreover, molecular identification recognized the *F. brachygibbosum* as *F. oxysporum*. The vascular browning, root, and crown rot symptoms were

visible in diseased plants in the pathogenicity test (Fig. 2).

Figs. 1 and 3 show the morphological characteristics of these isolates. Table 1 shows the results of morphological characterization of *Fusarium* spp. Isolates (Table 1).





Figure 1 Colones of Fusarium spp. A: Fusarium proliferatum, B: Fusarium oxysporum, C: Fusarium culmorum, D: Fusarium solani and E: Fusarium brachygibbosum.



**Figure 2** Necrosis and browning of the vessels in the crown region of diseased carnation in the greenhouse in pathogenicity test of *Fusarium* isolates A: F14, B: F13, C: F17, and D: F19.



**Table 1** Morphological characteristics of Fusarium spp. isolated from rhizosphere, crown, and root of Dianthuscaryophyllus.

Entries	F. proliferatum	F. oxysporum	F. solani	F. culmorum
Pigmentation on PDA	white but may become	white to pale violet	White to cream	pale orange but
	purple-violet with age			becomes brown
Growth rate	R	R	R	R
Macroconidia	+	+	+	+
Apical cell	Cu	Ta and cu	Bl and Ro	Bl and Ro
Basal cell	Pd	Fs	Fs or pd	Nwfs
Septate	3-5	3	5-7	3-4
Size (µm)	27-90 × 3-6	$40-85 \times 3.5-9$	30-72 × 5-8	30-60 × 5-8
Microconidia	+	+	+	-
Form	Lc, Fh	Fh	Fh	-
Phialides	Mono, Poly	Mono	Mono	-
Shape	Cl, Py	Ov, Ki	Ov, Re, Fu	-
Septate	0	0	0-2	-
Size (µm)	7-16.5 × 2-5	6.9-8.9 × 3-5	7-15 × 2-4	-
Clamydospore	-	+	+	+

+: Presence, -: Absence, Cu: Curved, Bl: Blunt, Ro: Rounded, Ta: Tapered, Pd: Poorly developed, Fs: Foot shape, Nwfs: Notched and without a distinct foot shape, Lc: Long chains, Fh: False heads, Cl: Club, Py: Pyriform, Ov: Oval, Re: Reniform, Fu: Fusiform, Ki: Kidney shaped.

Molecular identification of Fusarium species

Primers ef1/ef2 successfully identified and amplified a 700bp fragment containing the *TEF-1* $\alpha$  gene in *Fusarium* species. The presence of a clear band in fungal isolates indicates the correct amplification of this gene. Also, no amplicon was recorded in the negative control (Fig. 4).

Molecular identification results based on the TEF-1 $\alpha$  gene with average identities of 99.33%

supported morphological identification results. The only exception was the F14 isolate, which was morphologically identified as *F. brachygibbosum* and molecularly as F. solani, using the *Fusarium* ID site with 93.87% identity. This isolate in the BLAST site with 93% identities was recognized as *F. metavorans. Fusarium* species were identified based on the translation-elongation factor -1 alpha gene (TEF-1 $\alpha$ ), according to table 2.



Figure 4 Amplification of conserved region of TEF-1a gene. C: 100 bp DNA ladder, F1-F20: fusarium isolates.

Table 2 Characterization	of Fusarium	isolates and their	<ul> <li>GenBank accession</li> </ul>	n number.
--------------------------	-------------	--------------------	---------------------------------------	-----------

Fungi	Host	Isolation source	Morphological	Molecular identification based	GenBank
ID Host		isolation source	identification	on TEF-1A-1α (% similarity)	Accession number
F1	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (99.07%)	MT723967
F2	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (99.07%)	MT723968
F3	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (99.69%)	MT723969
F4	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (99.24%)	MT723975
F5	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (97.92%)	MT723977
F6	D. caryophyllus	Rhizosphere	F. solani	F. solani (98.78%)	MT723976
F7	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (98.74%)	MT723978
F8	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (99.4%)	MT723979
F9	D. caryophyllus	Rhizosphere	F. proliferatum	F. proliferatum (97.53%)	MT723980
F10	D. caryophyllus	Rhizosphere	F. culmorum	F. culmorum (98.87%)	MT753012
F11	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (98.58%)	MT753011
F12	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (99.84%)	MT723981
F13	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (100%)	MT723971
F14	D. caryophyllus	Crown and root	F. brachygibbosum	F. solani (93.87%)	MT681918
F15	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (99.84%)	MT723970
F16	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (99.84%)	MT723982
F17	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (100%)	MT723972
F18	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (99.85%)	MT723983
F19	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (99.7%)	MT723973
F20	D. caryophyllus	Crown and root	F. oxysporum	F. oxysporum (100%)	MT723974

The TEF-1 $\alpha$  partial gene sequences of *Fusarium* isolates were analysed to conduct a phylogenetic tree. As shown in Figure 5, *Fusarium* isolates were divided into five general groups. The first group consisted only of F14 isolate that was identified *F. brachygibbosum* morphologically and *F. solani* molecularly. This isolate was far from the rest. The second group consisted of F6 that identified as *F. solani* morphologically and

molecularly. The third group consisted of F10 identified as *F. culmorum*. The fourth group included F1, F2, F3, F4, F5, F7, F8, and F9, all of which were identified as *F. proliferatum*. The fifth group comprised F11, F12, F13, F15, F16, F17, F18, F9, and F20, identified as *F. oxysporum*. Thus, phylogenetic analysis was able to differentiate *Fusarium* species based on the TEF- $1\alpha$  gene.



**Figure 5** Bayesian 50% majority rule consensus tree deduced using EF1- $\alpha$  sequences of the *Fusarium* spp. under the GTR + G model. The newly generated sequences are in color frameworks.

#### Discussion

Since Fusarium species carnation is among the most severe pathogens in greenhouses, accurate identification of carnation related Fusarium species can be the first step in combating Fusarium disease and preventing damage to this plant. The purpose of this study is the identification of Fusarium spp. associated with carnation. In this research, molecular identification of Fusarium isolates detected the four species, including F. oxysporum, F. proliferatum, F. solani, and F. culmorum, in the order of frequency. F. oxysporum is known as a wilting agent in carnation. F. oxysporum seems to be one of the most harmful pathogens for greenhouse crops in Iran. Also, Sahampoor et al. (2020) oxysporum mentioned F. f.sp. radiciscucumerinum as an important pathogen in greenhouse cucumber. They considered the use of Trichoderma to be effective in laboratory control of this pathogenic fungus, and by using gamma rays, they increased the efficiency of Trichoderma in the control of F. oxysporum. Fattahi et al. (2013) also reported wilt disease caused by F. proliferatum and F. solani on carnation in Iran. Although there are many nonpathogenic species of Fusarium, all F. oxysporum species were isolated from diseased plants were pathogens based on the results of the pathogenicity test. Of the 20 Fusarium isolates collected in Mahalat, F.oxysporum was the most prevalent with 45% (9 of 20 isolates), followed by F. proliferatum with a frequency of 40% (8 of 20 isolates). The least frequent isolate was F. culmorum, with a frequency of 5% (1 of 20 isolates). The use of the TEF-1 $\alpha$  gene, which encodes an essential part of the 1a translation elongation factor, has been suggested as a useful phylogenetic tool for detecting Fusarium species. This gene is highly informative at the species level in Fusarium, and nonorthologous copies of the gene have not been detected in the genus. The primers of ef1 and ef2 can successfully amplify this region for all Fusarium species (Geiser et al., 2004).

Camacho López *et al.* identified *Fusarium* oxysporum, *F. proliferatum*, *F. globosum*, *F.* solani, *F.incarnatum*, *F. equiseti*, and *F.* tricinctum that were associated with diseased carnation plants in Baja California. Molecular identification of these isolates was performed based on the TEF-1 $\alpha$  gene. (Camacho López *et al.*, 2014). Rezaee *et al.* (2018) identified races of *Fusarium solani* f. sp. cucurbitae using specific primers based on TEF-1 $\alpha$  gene. Rabiei-Motlagh *et al.* (2017) showed that PCR assay using *Avr1*, *Avr2*, and *Avr3* genes could distinguish between nonpathogenic and low virulence strains of *Fusarium oxysporum* f. sp. *lycopersici.* 

Considering the correlation between molecular identification and morphological results, it seems that the molecular method can be faster and more straightforward for identifying these pathogens. In this research, the TEF-1 $\alpha$  gene was well able to identify and distinguish species. The results of this study can help identify diseases caused by *Fusarium* in carnation greenhouses.

#### **Conflict of Interests**

The authors declare that they have no conflicting interests.

#### **Authors' Contributions**

Samira Shahbazi has collected, isolated, and purified fungal isolates, contributed to writing the discussion (Research project collaborator); Fatemeh Zaker Tavallaie designed experiment, Performed molecular identification and gene registration in NCBI and wrote the manuscript (Executor of the research project) and Zoha daroodi performed morphological identification, DNA extraction and PCR amplification (Research project collaborator).

#### Acknowledgments

This work was supported by the Higher Education Complex of Shirvan (Project Title: "Morphological and molecular identification of isolated *Fusarium* from Dianthus and its pathogenicity test" with grant number 3474.

#### References

- Al-Hatmi, A. M., Ahmed, S. A., Van Diepeningen, A. D., Drogari-Apiranthitou, M., Verweij, P. E., Meis, J. F. and De Hoog, G. S. 2018. *Fusarium metavorans* sp. nov.: the frequent opportunist 'FSSC6'. Medical Mycology, 56(1): S144-S152.
- Camacho López, M. D., Valenzuela Solano, C. and Hernández Martínez, R. 2014. Fusarium spp. associated with carnation (*Dianthus* caryophyllus L.) in Baja California, Mexico. Revista Mexicana de Ciencias Agrícolas. 5: 1409-1415. https://doi: 10.29312/remexca. v0i8.1096.
- Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Molecular Biology and Evolution, 17: 540-552.
- Etebarian, H. R. 1996. Occurrence of *Fusarium* wilts of camation in Varamin area. Iranian Journal of Plant Pathology, 32: 223-232. (in Persian).
- Fattahi, B., Rahanandeh, H., Chavoshi, S., Zamanizadeh, H. R., Bayat, H. and Moshayedi, M. 2013. Report primary of *Fusarium proliferatum* and *Fusarium solani* agents of *Dianthus caryophyllus* wilting in Markazi province in Iran. International Journal of Biosciences, 12: 1-7. https://doi: http://dx.doi.org/10.12692/ijb/3.
- Geiser, D. M., Jime'nez-Gasco, M. M., Kang, S., Makalowska, I., Veeraraghavan, N., Ward, N., Zhang, T. J., Kuldau, G. A. and O'Donnell, K. K. 2004. *FUSARIUM*-ID v. 1.0: A DNA sequence database for identifying *Fusarium*. European Journal of Plant Pathology, 110: 473-479. https://doi.org/10.1023/B:EJPP.0000032386. 75915.a0.
- Herron, D. A., Wingfield, M. J., Wingfield, B. D., Rodas, C. A., Marincowitz, S. and Steenkamp, E. T. 2015. Novel taxa in the *Fusarium fujikuroi* species complex from Pinus spp. Studies in Mycology, 80: 131-150.
- Hirooka, Y., Rossman, A.Y. and Chaverri, P. 2011. A morphological and phylogenetic

revision of the Nectria cinnabarina species complex. Studies in Mycology, 68: 35-56.

- Huson, D. H. and Scornavacca, C. 2012. Dendroscope 3: An interactive tool for rooted phylogenetic trees and networks. Systematic Biology, 61: 1061-1067.
- Katoh, K. and Standley, D. M. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution, 30: 772-780.
- Kermajany, Z., Jamali Zavareh, A. and Fadaei Tehrani, A. 2017. *In vitro* inhibitory effects of five strains of Trichoderma on the growth of *Fusarium oxysporum* f. sp. *dianthi*. Biological Control of Plant Pests and Diseases, 1: 121-125. https://doi: 10.22059/jbioc.2017.63360.
- Larget, B. and Simon, D. L. 1999. Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. Molecular Biology and Evolution, 16: 750-759.
- Laurence, M. H., Summerell, B. A., Burgess, L. W. and Liew, E. C. Y. 2011. Fusarium burgessii sp. nov. representing a novel lineage in the genus Fusarium. Fungal Diversity, 49(1): 101-112.
- Leslie, J. F. and Summerell, B.A. 2006. The *Fusarium* Laboratory Manual. Blackwell Publishing. 1a. Ed. USA, 388 p. https://doi:10.1002/9780470278376.
- Moeini, A., Torabi, M. and Shahriari, D. 2014. Evaluation of *Trichoderma* spp. isolates for biological control of fusarium wilt of carnation in greenhouse. Applied Phytopathology, 3(2): 131-144. (in Persian).
- Nylander, J. A. A. 2004. MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre: Uppsala University.
- Papizadeh, M., van Diepeningen, A. D., Zamanizadeh, H. R., Saba, F. and Ramezani, H. 2018. Fusarium ershadii sp. nov., a pathogen on Asparagus officinalis and Musa acuminata. European Journal of Plant Pathology, 151(3): 689-701.
- Rabiei-Motlagh, E., Rouhani, H., Shokouhifar, F., Falahati Rastegar, M., and Taheri, P.

2017. Molecular identification of Formae speciales and racial identity in Iranian strains of *Fusarium oxysporum* f. sp. lycopersici: detection of avirulence genes. Journal of Crop Protection, 6(1): 67-77.

- Rezaee, S., Gharanjik, S. and Mojerlou, S. 2018. Identification of *Fusarium solani* f. sp. cucurbitae races using morphological and molecular approaches. Journal of Crop Protection, 7(2): 161-170.
- Ronquist, F. and Huelsenbeck, J. P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics, 19: 1572-1574.
- Sahampoor, L., Zaker Tavallaie, F., Fani, S. R. and Shahbazi, S. 2020. In vitro efficiency of *Trichoderma harzianum* mutants in biocontrol of *Fusarium oxysporum* f. sp. radicis-cucumerinum. Journal of Crop Protection, 9(2): 285-300.

### شناسایی مورفولوژیکی و مولکولی فوزاریومهای مرتبط با میخک Dianthus caryophyllus در ایران، محلات

سمیرا شهبازی'، فاطمه ذاکر تولایی ٔ و ضحی درودی ٔ

**چکیده:** میخک یک گیاه زینتی با اهمیت اقتصادی در سراسر جهان است. قارچ *Fusarium* یکی از عوامل مهم آلودگی میخک در شهرستان محلات است. بعضی از گونههای Fusarium باعث پژمردگی و پوسیدگی طوقه و ریشه در میخک میشوند. در این مطالعه، ۲۰ جدایه قارچی از جنس Fusarium از گیاهان میخک دارای علائم بیماری و همچنین ریزوسفر این گیاهان از گلخانههای گل میخک در محلات جداسازی شدند. جدایهها از نظر مورفولوژی با استفاده از کلید شناسایی Leslie و Summerell شناسایی شدند. تکثیر ناحیه ژنی TEF-1۵ با استفاده از آغازگرهای ef1 و ef2 انجام شد. قطعات تکثیر شده جداسازی و توالییابی شدند. این توالیهای نوکلئوتیدی در پایگاه داده Fusarium ID جهت شناسایی مولکولی جدایهها براساس ژن TEF-1a وارد شدند و درخت فیلوژنتیکی نیز رسم شد. نتایج نشان داد که جدایههای فوزاریوم به پنج گروه تقسیم میشوند: F14 از نظر ریختشناسی بهعنوان Fusarium brachygibbosum و از لحاظ مولکولی F. solani شناخته شد. F6 از نظر ریختشناسی و مولکولی بهعنوان F. solani بهعنوان F. culmorum شناسایی شد. سایر جداشدههای طوقه و ریشه F13 ،F13 ،F13 ،F13 ،F13 ،F13 ،F13 ،F11 و F10 ,F12 ،F13 ،F13 ،F12 ،F11 ) شناخته شدند. جدایههای ناحیه ریزوسفر (F1، F3، F3، F4، F3، F1 و F9) به عنوان F. proliferatum تشخیص داده شدند. این مطالعه نشان داد که استفاده از ژن TEF-1a برای شناسایی مولکولی جدایههای Fusarium یک روش نسبتاً دقیق، سریع و ساده برای تشخیص گونههای قارچ است و میتواند در مطالعات آسیبشناسی گیاه مورد استفاده قرار گیرد. شناسایی پاتوژن اولین قدم برای کنترل بیماری است.

واژگان کلیدی: میخک، فوزاریوم، شناسایی مولکولی، ژن TEF-1a