# Influence of different host plant cultivars on nutritional indices of *Plutella xylostella* (Lepidoptera: Plutellidae)

Roja Kianpour<sup>1</sup>, Yaghoub Fathipour<sup>1</sup>\*, Javad Karimzadeh<sup>2</sup> and Vahid Hosseininaveh<sup>3</sup>

1. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P. O. Box: 14115-336, Tehran, Iran.

2. Isfahan Research Centre for Agriculture and Natural Resources, Isfahan, Iran.

3. Department of Plant Protection, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran.

**Abstract:** The diamondback moth, *Plutella xylostella* (L.) is a serious and economically important pest of cruciferous crops worldwide. The nutritional indices of this pest on four host plants including two canola (SLM<sub>046</sub> and RGS<sub>003</sub>) and two cabbage cultivars (Green-Cornet and Glob-Master) were determined under laboratory conditions ( $25 \pm 0.5$  °C,  $65 \pm 5\%$  RH and 16: 8 L: D h). The highest relative consumption rate (RCR) and relative growth rate (RGR) were recorded on RGS<sub>003</sub> (95.68 ± 14.52 mg/mg/day) and SLM<sub>046</sub> (0.382 ± 0.041 mg/mg/day), respectively. The efficiency of the conversion of ingested food (ECI) and efficiency of conversion of digested food (ECD) were significantly higher on SLM<sub>046</sub> (2.298 ± 0.646 and 2.471 ± 0.719%, respectively). However, there was no significant difference among cultivars regarding approximate digestibility (AD) index. The highest feeding performance of the larvae of *P. xylostella* was found on SLM<sub>046</sub>. The nutritional indices are appropriate tools in host plant resistance evaluation and they also could provide profound understanding of the behavioral and physiological bases of insect-host plant interactions.

Keywords: nutritional indices, canola, cabbage, Plutella xylostella

## Introduction

The diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) is the most important pest of Brassicaceae crops in the world. Importance of *P. xylostella* is reflected in (i) being the most widely distributed of all the Lepidoptera (Talekar and Shelton, 1993;), (ii) its high reproductive potential (over 20 generations per year in tropics), (iii) lack or disruption of its natural enemies (Hui *et al.*, 2010) and (iv) its genetic elasticity facilitating rapid development of resistance to chemical insecticides (Sayyed *et al.*, 2004; Sarfraz *et al.*, 2007; Ahmad *et al.*, 2012). Insecticide application is the primary method of

control of *P. xylostella*, but high tolerance to most of insecticides and associated environmental problems may result in outbreaks of the pest by destruction of its natural enemies (Ahmad *et al.*, 2012). These drawbacks have increased interest in other control methods such as biological control and resistant cultivars of host plants. Therefore, the mentioned methods are useful and desirable tools in most pest management programs because they are effective and non-toxic to natural enemies, human and the environment (Haseeb *et al.*, 2004; Xu *et al.*, 2004; Fathipour and Sedaratian, 2013).

Study of biological, physiological and ecological characteristics of insect pests on different host plants is a way to recognize the host plant resistance to these herbivores (Sarfraz *et al.*, 2007). The role of quality of herbivores' food in insect-plant relationship has been determined in some studies (David and Gardiner, 1962; Sheiker



Handling Editor: Dr. Ali Asghar Talebi

<sup>\*</sup> Corresponding author, e-mail: fathi@modares.ac.ir Received: 16 August 2013, Accepted: 30 September 2013

*et al.*, 2001; Xue *et al.*, 2010). The rates of food ingestion, growth and utilization efficiency are important components of herbivores' performance. From a nutritional point of view, utilization efficiency reflects the quality of food consumed (Naseri *et al.*, 2010; Baghery *et al.*, 2013).

Few studies are available regarding the nutritional indices of P. xylostella. Some of these studies have presented the effects of chinaberry extract (Chen et al., 1996), antifeedants (Perera et al., 2000) and neem-based insecticides (Ahmad et al., 2012) on these indices. Chen et al., (1996) showed that the food consumption was reduced at high concentrations (2 and 4%) of chinaberry fruit extract. Also, some antifeedants significantly decreased the food consumption of P. xylostella larvae (Perera et al., 2000. Ahmad et al., (2012) suggested that high concentrations (15 and 20 ppm) of Neemazal, one of the neem-based insecticides, significantly affected the nutritional indices of the larvae of P. xylostella by reducing consumption and utilization rate of available food.

Growth, development and reproduction of insects strongly depend on the quality and quantity of food consumed (Scriber and Slansky, 1981). This information could help to assess the crop loss properly and design a suitable integrated pest management (IPM) program to keep the population of a herbivorous pest below an economically important level. Determining the nutritional indices of an insect on a particular host is a tool for evaluating host plant resistance mechanisms that could improve pest management programs. In this study, the effects of different cultivars of host plants on the nutritional indices of *P. xylostella* were studied.

#### **Materials and Methods**

#### **Insect and plants**

Seeds of four different cultivars of two host plants including canola, *Brassica napus* L. (SLM<sub>046</sub>, RGS<sub>003</sub>) and cabbage, *Brassica oleracea var*. *capitata* (Green-Cornet, Glob-Master) were obtained from the Seed and Plant Improvement Institute (Karaj, Iran). These seeds were individually planted in the greenhouse of Tarbiat Modares University in plastic pots, without any fertilizers and pesticides. The leaves of cruciferous cultivars were used for experiments when the plants were 4 weeks old. The original population of *P. xylostella* was collected from the cabbage fields of University of Tehran (Karaj, Iran), during September 2010. The collected specimens were maintained in a growth chamber at  $25 \pm 0.5$  °C,  $65 \pm 5\%$  RH and 16: 8 h (L:D). The F<sub>2</sub> generation of *P. xylostella* was used in all experiments.

#### **Feeding assays**

Newly molted fourth instar larvae of P. xylostella were used to measure their nutritional indices on different host plants tested. The larvae were individually transferred into plastic containers (diameter: 10 cm, depth: 1.5 cm) with a hole on its top that was covered by fine mesh net for ventilation. The experiment was started with 20 replications per cultivar. Every 8 hours, the weights of the fourth instar larvae, food given, unconsumed food and feces produced were recorded until the pupal stage. To calculate the dry weight of the larvae, feces and leaves of each cultivar, extra specimens (10 specimens for each treatment) were weighed, oven-dried (until completely dried at 60 °C), and then re-weighed to establish a percentage of their dry weight. The nutritional indices were calculated based on the dry weights using the formulae presented in Waldbauer (1968) (Table 1).

#### Statistical analysis

Normal distribution of the data obtained from feeding assays was tested by Kolmogrov-Smirnov test. The data for the effects of different cultivars of cruciferous plants (set as completely randomized design) on nutritional indices were subjected to one-way analysis of variance (ANOVA) to determine the similarities or significant differences. Statistical differences among means were evaluated using Tukey test at  $\alpha = 0.05$ .

#### Results

Significant differences among host plants for larval weight, food consumed and feces produced by the fourth instar larvae of *P. xylostella* are presented in Table 2. The highest and lowest values of larval weight were on Green-Cornet (0.980 mg) and

Glob-Master (0.110 mg), respectively (F = 107.39; df = 3, 60; P < 0.001). Food consumed by the fourth instars were highest on Green-Cornet (48.08 mg) and lowest on Glob-Master (21.52 mg) (F = 5.37; df = 3, 60; P < 0.03). The amounts of feces produced ranged from 0.522 (RGS<sub>003</sub>) to 1.184 mg (SLM<sub>046</sub>) (F = 3.39; df = 3, 60; P < 0.05).

Different host plants affected the relative growth rate (RGR) of the larvae of *P. xylostella* (Table 3). The lowest and highest RGR values belonged to the larvae reared on Green-Cornet and SLM<sub>046</sub>, respectively (F = 11.62; df = 3, 59; P < 0.01). The data on the effects of different host plants on the relative consumption rate (RCR) are given in Table 3. The RCR was significantly different among cultivars tested (F = 13.57; df = 3, 59; P < 0.01) and its value ranged from 95.68 mg/mg/day on RGS<sub>003</sub> to 21.74 mg/mg/day on Green-Cornet. The efficiency of conversion of digested food (ECD) as the function of different cultivars of cruciferous plants is summarized in Table 3. Among all of the cultivars tested, the ECD was not significantly different except on SLM<sub>046</sub> (F = 4.67; df = 3, 59; P <0.01). The effects of different cruciferous cultivars on the conversion of ingested food efficiency (ECI) are indicated in Table 3. The highest value of ECI was related to the larvae reared on SLM<sub>046</sub> and it was significantly different from other cultivars tested (F = 4.93; df = 3, 59; P < 0.01). As shown in Table 3, the fourth larval instar had no significant difference regarding approximate digestibility (AD) on different cultivars (F = 2.28; df = 3, 59; P > 0.05).

Table 1 The formulae of the nutritional indices calculated for *Plutella xylostella* (Waldbauer, 1968).

Abbreviation	Nutritional index	Formula	
RCR	Relative consumption rate	$E/(A \times T)$	
RGR	Relative growth rate	$P/(A \times T)$	
AD	Approximate digestibility	[(E-F)/E] × 100	
ECI	Efficiency of conversion of ingested food	(P/E) × 100	
ECD	Efficiency of conversion of digested food	$[P/(E-F)] \times 100$	

P = dry weight gain (mg), A = initial and final mean dry weights of the larvae during feeding period (mg), E = dry weight of food ingested (mg), T = duration of feeding period (days), F = the dry weight of fees produced (mg).

**Table 2** Mean (+SE) larval weight of *Plutella xylostella*, food consumed and feces produced by fourth instar larvae on different cultivars.

Cultivars	Larval weight (mg)	Food consumed (mg)	Feces produced (mg)
SLM <sub>046</sub>	$0.786\pm0.049b$	$35.23\pm2.730ab$	$1.184 \pm 0.207a$
RGS <sub>003</sub>	$0.179\pm0.009c$	$30.51\pm4.061ab$	$0.522 \pm 0.086b$
Green-Cornet	$0.980\pm0.058a$	$48.08 \pm 4.552a$	$0.928\pm0.194ab$
Glob-Master	$0.110\pm0.005c$	$21.52\pm3.875b$	$0.713 \pm 0.096ab$

Means followed by the same letter within the same column are not significantly different ( $\alpha = 0.05$ , Tukey).

<b>Table 3</b> Nutritional indices (mean $\pm$ SE) of fourth instar larvae of <i>Plutella xylostella</i> on different cult
--

Cultivars	RCR (mg/mg/day)	RGR (mg/mg/day)	ECD (%)	ECI (%)	AD (%)
SLM <sub>046</sub>	$28.838\pm2.325b$	$0.382\pm0.0412a$	$2.471\pm0.719a$	$2.298 \pm 0.646a$	$95.867 \pm 0.715 a$
RGS <sub>003</sub>	$95.685 \pm 14.521a$	$0.335\pm0.025a$	$0.563\pm0.111b$	$0.539\pm0.103b$	$97.557 \pm 0.618 a$
Green-Cornet	$21.743\pm2.379b$	$0.139\pm0.016b$	$0.808 \pm 0.166b$	$0.782\pm0.152b$	$97.866 \pm 0.549a$
Glob-Master	$93.099 \pm 15.749a$	$0.216\pm0.027ab$	$0.387\pm0.109b$	$0.356\pm0.093b$	$95.321 \pm 1.377a$

Means followed by the same letters in the same columns are not significantly different ( $\alpha = 0.05$ , Tukey). AD = approximate digestibility, ECI = efficiency of conversion of ingested food, ECD = efficiency of conversion of digested food, RCR = relative consumption rate, RGR = relative growth rate.

#### Discussion

Resistant cultivars are recognized as a key factor in IPM programs (Endo et al., 2007; Fathipour and Sedaratian, 2013). The quality and quantity of food consumed by a pest can biological affect its and physiological parameters (Nation 2000; Golizadeh et al., 2009). Recent demographic studies suggested SLM<sub>046</sub> and Green-Cornet as susceptible cultivars and RGS<sub>003</sub> and Glob-Master as resistant cultivars to P. xylostella (Soufbaf et al., 2010; Kianpour et al., 2013). The larvae of P. xylostella fed on the susceptible cultivars (SLM<sub>046</sub> and Green-Cornet) had lower RCR values than those fed on the resistant ones (RGS<sub>003</sub> and Glob-Master). The larvae fed on SLM<sub>046</sub> and Green-Cornet had more weight with lower amount of food consumed. On the other hand, the larvae fed on RGS<sub>003</sub> and Glob-Master consumed more food to reach suitable weight because these host plants may have not good quality or the values of nutritional elements were not enough (Slansky and Scriber, 1982). The highest RGR on SLM<sub>046</sub> showed its high quality and suitability as a host plant for the larvae of *P. xylostella*.

More than 95% of amount of total food consumed by the fourth instar larvae of P. xylostella was digested. It might be linked to high ability of digestive enzymes of the larvae to digest food consumed. It seems that P. xylostella has a good conformity with the range of host plants and high ability to digest eaten food. On the other hand, about 2% of food eaten was converted to biomass on susceptible cultivar (SLM<sub>046</sub>). Indeed, the high amount of AD could be explained in two ways: (i) increase in RGR due to energy of digested food or (ii) repelling most of eaten food as feces. The data showed that the second explanation is more acceptable, showing that a great percentage of eaten food was converted to feces (Xue et al., 2010). Considering ECD and ECI values, about 2% of the ingested food was converted to biomass in the larvae that were fed on  $SLM_{046}$ . The lower ECI on other cultivars might be related to decrease in efficiency of converting

the ingested food into growth. The larvae feeding on  $SLM_{046}$  had the highest value of ECD, showing that these larvae have more efficiency for the conversion of digested food to their biomass. Any changes in ECI and ECD values represent the proportion of ingested food converted into energy (Koul *et al.*, 2004; Baghery *et al.*, 2013).

Analysis of the nutritional indices can lead to understand the behavioral and physiological bases of insect- plant interactions (Lazarevic and Peric-Mataruga, 2003). Lower fitness of P. xylostella on some cultivars may depend on presence of some secondary phytochemicals in the cruciferous plants, e.g. glucosinolate (GS), or may depend on the absence of some primary nutrients such as nitrogen, which is necessary growth and development of insect for herbivores (Gols et al., 2008; Soufbaf et al., 2012). Some studies showed that high level of glucosinolates is the main factor that reduces the performance of P. xylostella (Li et al., 2000; Gols et al., 2008; Soufbaf et al., 2012). On the other hand, Soufbaf et al., (2012) reported that P. xylostella developed poorly on the resistant cultivar of canola, which has lower value of nitrogen in its leaf tissues.

In conclusion, this study revealed that among different measures to be applied for evaluating host plants resistance in IPM programs, the nutritional indices could have an important place and they might be used in combination with other methods for getting more reliable results. This conclusion has been supported by comparing the results of the nutritional indices obtained in this study with the life table parameters and digestive enzymes of *P. xylostella* (unpublished data) in which in most cases they were in conformity with one another.

## References

- Ahmad, N. A., Ansari, S. and Hasan, F. 2012. Effects of neem based insecticides on *Plutella xylostella* (Linn.). Crop Protection, 34: 18-24.
- Baghery, F., Fathipour, Y. and Naseri, B. 2013. Nutritional indices of *Helicoverpa armigera*

(Lepidoptera: Noctuidae) on seeds of five host plants. Applied Entomology and Phytopathology, 8: 19-27.

- Chen, C. C., Chang, S. J., Cheng, L. L. and Hou, R. F. 1996. Effect of Chinaberry extract on feeding, growth and fecundity of diamondback moth, *Plutella xylostella* Linn. (Lep., Yponomeutidae). Journal of Applied Entomology, 120: 341-345.
- David, W. A. L. and Gardiner, B. O. C. 1962.Observation on the larva and pupa of *Pieris brassicae* (L.) in the laboratory culture. Bulletin of Entomological Research, 53: 414-436.
- Endo, N., Hirakawa, I., Wada, T. and Tojo, S. 2007. Induced resistance to the common cutworm, *Spodoptera litura* (Lepidoptera: Noctuidae) in three soybean cultivars. Applied Entomology and Zoology, 42: 199-204.
- Golizadeh, A., Kamali, K., Fathipour, Y. and Abbasipour, H. 2009. Life table of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) on five cultivated brassicaceous host plants. Journal of Agricultural Science and Technology, 11: 115-124.
- Fathipour, Y. and Sedaratian, A. 2013. Integrated Management of *Helicoverpa armigera* in Soybean Cropping Systems. In: El-Shemy, H. (Ed.), Soybean-Pest Resistance. InTech, Rijeka, Croatia, pp. 231-280.
- Gols, R., Bukovinszky, T., van Dam, N. M., Dicke, M., Bullock, J. M. and Harvey, J. A. 2008. Performance of generalist and herbivores specialist and their endoparasitoids differs on cultivated and brassica population. Journal wild of Chemical Ecology, 34: 132-143.
- Haseeb, M., Liu, T. X. and Jones, W. A. 2004.
  Effects of selected insecticides on *Cotesia plutellae* (Hymenoptera: Braconidae), an endolarval parasitoid of *Plutella xylostella* (Lepidoptera: Plutellidae). Biological Control, 49: 33-46.
- Hui, W., Juan, W., Hong-shan, L., Hua-guo, D. and Xiao-jun, G. U. 2010. Sub-lethal effects of fenvalerate on the development, fecundity, and juvenile hormone esterase

activity of diamondback moth, *Plutella xylostella* (L.). Agricultural Sciences in China, 9: 1612-1622.

- Kianpour, R., Fathipour, Y., Karimzadeh, J. and Hosseininaveh, V. 2013. Bottom-up effects of brassicaceous genotypes on life table parameters of diamondback moth *Plutella xylostella*. Crop Protection, (in Press).
- Koul, O., Singh, G., Singh, R., Singh, J., Daniewski, W. M. and Berlozecki, S. 2004. Bioefficacy and mode-of-action some limonoids of salanin group from Azadirachta indica A. Juss and their role in multicomponent а system against lepidopteran larvae. Journal of Bioscience, 29: 409-416.
- Lazarevic, J. and Peric-Mataruga, V. 2003. Nutritive stress effects on growth and digestive physiology of *Lymantria dispar* larvae. Yugoslav Medical Biochemistry, 22: 53-59.
- Li, Q., Eigenbrode, S. D., Stringam, G. R. and Thiagarajah, M. R. 2000. Feeding and growth of *Plutella xylostella* and *Spodoptera eridania* on *Brassica juncea* with varying glucosinolate concentrations and myrosinase activities. Journal of Chemical Ecology, 26: 2401-2419.
- Naseri, B., Fathipour, Y., Moharramipour. S. and Hosseininaveh, V. 2010. Nutritional indices of the cotton bollworm, *Helicoverpa armigera*, on 13 soybean varieties. Journal of Insect Science, 10: 1-14.
- Nation, J. L. 2002. Insect Physiology and Biochemistry, CRC Press, Boca Raton, Florida.
- Perera, D. R., Armstrong, G. and Senanayake, N. 2000. Effect of antifeedants on the diamondback moth (*Plutella xylostella*) and its parasitoid *Cotesia plutellae*. Pest Management Science, 56: 486-490.
- Sayyed, A. H., Omar, D. and Wright, D. J. 2004. Genetics of spinosad resistance in a multiresistant field selected population of *Plutella xylostella*. Pest Management Science, 60: 827-832.
- Scriber, J. M. and Slansky, F. 1981. The nutritional ecology of immature insects. Annual Review of Entomology, 26: 183-211.

Downloaded from jcp.modares.ac.ir on 2024-05-03

- Sheikher, C., Sharma, R. and Sharma, K. C. 2001. Consumption and utilization of cabbage by cabbage butterfly *Pieris brassicae* (L.). Journal of Insect Science, 14: 33-35.
- Slansky, F. and Scriber, J. M. 1981. Selected bibliography and summary of quantitative food utilization by immature insects. Bulletin of the Entomological Society of America, 20: 43-56.
- Soufbaf, M., Fathipour, Y., Karimzadeh, J. and Zalucki, M. P. 2010. Bottom-up effect of different host plants on *Plutella xylostella* (Lepidoptera: Plutellidae): a life-table study on canola. Journal of Economic Entomology, 103: 2019-2027.
- Soufbaf, M., Fathipour, Y., Zalucki, M. P. and Hui, C. 2012. Importance of primary metabolites in canola in mediating interactions between a specialist leaf-feeding insect and its specialist solitary endoparastoid. Arthropod-Plant Interactions, 6: 241-250.

- Talekar, N. S. and Shelton, A. M. 1993. Biology, ecology, and management of the diamondback moth. Annual Review of Entomology, 38: 275-301.
- Waldbauer, G. P. 1968. The consumption and utilization of food by insects. Advances in Insect Physiology, 5: 229-288.
- Xu, Y., Liu, T., Leibee, G. L. and Jones, W. A. 2004. Effects of selected insecticides on *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of *Plutella xylostella* (Lepidoptera: Plutellidae). Biocontrol Science and Technology, 14: 713-723.
- Xue, M., Pang, Y. M., Wang, H. T., Li, Q. L. and Liu, T. X. 2010. Effects of four host plants on biology and food utilization of the cutworm *Spodoptera litura*. Journal of Insect Science, 10: 22.
- Zhu, J. H., Zhang, F. P. and Ren, H. G. 2005. Development and nutrition of *Prodenia litura* on four food plants. China Bulletin of Entomology, 42: 643-646.

# تأثیر میزبانهای مختلف گیاهی روی شاخصهای تغذیهای شب پره پشتالماسی Plutella (Lepidoptera: Plutellidae)

روجا کیان پور'، یعقوب فتحی پور<sup>۲\*</sup>، جواد کریمزاده اصفهانی<sup>۳</sup> و وحید حسینینوه<sup>†</sup>

۱- گروه حشرهشناسی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، صندوق پستی ۳۳۶–۱۴۱۱۵، تهران، ایران.
 ۲- مرکز تحقیقات کشاورزی و منابع طبیعی استان اصفهان، اصفهان، ایران.
 ۳- گروه گیاهپزشکی، پردیس کشاورزی و منابع طبیعی، دانشگاه تهران، کرج، ایران.
 \* پست الکترونیکی نویسنده مسئول مکاتبه: ۱۳۹۲
 دریافت: ۲۵ مرداد ۱۳۹۲؛ پذیرش: ۸ مهر ۱۳۹۲

چکیده: شبپره پشتالماسی . Plutella xylostella L یکی از آفات مخرب محصولات خانواده کلمیان در جهان است. در این مطالعه، شاخصهای تغذیه ی لاروهای شبپره پشتالماسی روی دو میزبان گیاهی کلزا (ارقام SLM<sub>046</sub> و RGS<sub>003</sub>) و کلم پیچ (ارقام Green-Cornet و Glob-Master) در دمای ۵/۰ ± ۲۵ درجه سلسیوس، رطوبت نسبی ۵ ± ۶۵ درصد و دوره نوری ۱۶ ساعت روشنایی و ۸ ساعت تاریکی مورد مطالعه قرار گرفت. بیشترین میزان نرخ مصرف نسبی (RCR) و نرخ رشد نسبی (RGR) لاروهای سن چهارم شب پره پشتالماسی روی RGS<sub>003</sub> بهمیزان ۲۴/۵۲ ± ۸۹/۹۹ (میلی گرم/میلی گرم/روز) و SLM<sub>046</sub> شب پره پشتالماسی روی RGS<sub>003</sub> بهمیزان ۲۴/۵۲ ± ۸۹/۹۹ (میلی گرم/میلی کارایی تبدیل غذای خورده شده (ECI) و کارایی تبدیل غذای هضم شده (ECD) (بهترتیب ۶۶۴۶ ± ۲۲۹۸ و ۲/۲۹۸ و ۲/۴۹۰ درصد) روی رقم SLM<sub>046</sub> مشاهده شد. میزان شاخص تقریبی هضمشوندگی (AD) تفاوت معنیداری روی ارقام مختلف کلزا و کلم نداشت. این نتایج نشان داد که لاروهای شبپره میتالماسی کارایی تغذیه ای بهتری را روی رقم SLM<sub>046</sub> دارند. تجزیه و تحلیل شاخصهای تغذیه ای

**واژگان کلیدی**: شاخصهای تغذیه، Plutella xylostella، کلم، کلزا