

## Transcript analysis of some pathogenicity-related elements in an Iranian A\* isolate of *Xanthomonas citri* subsp. *citri*

Raana Roohani<sup>1, 2</sup>, Masood Soltani Najafabadi<sup>3</sup>, Seyed Mehdi Alavi<sup>1\*</sup>, Naser Farrokhi<sup>4</sup> and Masoud Shams-bakhsh<sup>2\*</sup>

1. Department of Plant Biotechnology, National Institute for Genetic Engineering and Biotechnology (NIGEB), P. O. Box: 14155-6343, Tehran, Iran

2. Plant Pathology Dept., Faculty of Agriculture, Tarbiat Modares University (T. M. U.), Tehran-Iran

3. Oil Seeds Research Department, Seed and Plant Improvement Institute (SPII), Karaj, Iran

4. Department of Agronomy and Plant Breeding, Faculty of Agricultural Engineering, Shahrood University of Technology, P. O. Box: 361999516, Shahrood, Iran

**Abstract:** Asiatic citrus canker is a devastating disease resulting in drastic economic losses in citriculture worldwide. Amongst three different types of the disease, i.e. A, A\* and A<sup>w</sup>, the A\* type is genetically less known. In order to comprehend the behavior of the Asiatic citrus canker A\*-type strain (*Xanthomonas citri* subsp. *citri*) in the vicinity of the host cells, a targeted semi-quantitative transcript analysis approach via RT-PCR was carried out. A subset of sixteen genes, as representative of different steps involved in phytopathogenicity, was analyzed on the culture medium (as uninduced) and compared with the subset isolated from the infected Mexican lime (*Citrus aurantifolia* L.) plants (as induced). The results showed that certain genes were up-regulated in induced condition, suggesting a putative role in bacteria-host interaction. Furthermore, the transcripts in induced condition could be classified into constitutive, early- and late-responsive genes, demonstrating their functional relevance during the host-pathogen interaction.

**Keywords:** RT-PCR, *Xcc*, citrus canker, Mexican lime

### Introduction

*Xanthomonas* spp. are Gram-negative bacteria belonging to *Gammaproteobacteria* in Xanthomonadaceae family, with 27 species, causing serious diseases in nearly 400 plant species in a host and/or even tissue specific manner (Qian *et al.*, 2008; Ryan *et al.*, 2011). Amongst others, some pathotypes (A, A\* and A<sup>w</sup>) of *Xanthomonas citri* namely asiatic citrus canker (ACC) affect a wide range of hosts

including most commercial citrus species and many close relatives of Rutaceae (Verniere *et al.*, 1998; Cubero and Graham, 2002; Qian *et al.*, 2008; Vojnov *et al.*, 2010). Consequently and as a result of quarantine restrictions, citrus market and trade are affected enormously endangering the zest for its production (Brunings and Gabriel, 2003; Schaad *et al.*, 2005; Guo *et al.*, 2011). Hence, control of the disease poses a continuous challenge to the citrus orchardists (Traore *et al.*, 2009; Bui Thi Ngoc *et al.*, 2010). As in the war field, understanding the enemy and its tactics helps to better change the course and improve the controlling strategies. Here, a transcript analysis was performed to better appreciate the pathogenicity-related elements of ACC A\*-type strain.

Handling Editor: Dr. Mohammad Javaheri

\* **Corresponding author**, e-mail: mealavi@nigeb.ac.ir, shamsbakhsh@modares.ac.ir

Received: 12 May 2012; Accepted: 28 November 2012

Among different pathotypes of ACC, most scientific findings were focused on A with a wide range of hosts (Shiotani *et al.*, 2009; Bock *et al.*, 2010; Vojnov *et al.*, 2010; Ryan *et al.*, 2011), while little is known about A\* and A<sup>w</sup> with narrow range of host range specificity (Verniere *et al.*, 1998; Bui Thi Ngoc *et al.*, 2010). The main plant that gets affected by A\* is Mexican lime (*Citrus aurantifolia* L.), whilst A<sup>w</sup> causes canker in Mexican lime and alemow (*Citrus macrophylla* L.). Recently, the genetic diversity of a fairly large collection of Iranian *Xanthomonas citri* subsp. *citri* (*Xcc*-A\*) strains was evaluated (Soltaninejad *et al.*, 2010a, b; Rezaei *et al.*, 2012) and it was noted that A\* is genetically more divergent than A as previously reported by Bui Thi Ngoc *et al.*, (2010). Interestingly, the intra-population diversity was also remarkable among the A\* strains. Here, temporal expression of 16 virulence and/or pathogenicity-related genes (Table 1) in an *Xcc*-A\* strain was monitored semi-quantitatively. The relative importance of the investigated genes at different time points during establishment of the pathogen in the host plant is discussed.

## Materials and method

### Plant materials

*Citrus aurantifolia* young seedlings were prepared from Hormozgan province, Iran. The seedlings were kept in greenhouse at 30 °C under natural light and constant humidity for three months.

### Bacterial strains and inoculation

NIGEB-088, an isolate of *Xanthomonas citri* subsp. *citri* (*Xcc*), was provided by the Bacterial Citrus Canker Collection of National Institute of Genetic Engineering and Biotechnology (NIGEB). Bacterium was cultured on YPGA (yeast extract 3 g.l<sup>-1</sup>, pepton 5 g.l<sup>-1</sup>, glucose 7 g.l<sup>-1</sup> and agar 15 g.l<sup>-1</sup>) at 28 °C for 48 h. A single colony was sub-cultured in YP medium (yeast extract 3 g.l<sup>-1</sup> and pepton 5 g.l<sup>-1</sup>) on a rotary shaker at 180 rpm at 28 °C till OD<sub>600</sub> reached up to 0.3, *i.e.*, 5 × 10<sup>8</sup> cfu/ml. To prepare the inoculum, the culture was centrifuged 10 min at 5000 rpm and the pellet

was resuspended and diluted with dH<sub>2</sub>O to reach a concentration of 1 × 10<sup>5</sup> cfu/ml and 50 µl of which was considered as the uninduced bacterium. For inoculation, host plant leaves (4 leaves per plant in a total of 10 plants) were surface-sterilized with ethanol and the inoculum (50 µl of 1 × 10<sup>5</sup> cfu/ml) was injected into the parenchymal space on the abaxial leaf side.

### Bacterial recovery from the inoculated leaves

According to Mehta and Rosato (2003), bacteria were recovered in different time points (1, 2, 3, 4, and 7 days post inoculation) for RNA isolation. The harvested leaves were sterilized with 70% (v/v) ethanol for 3 min and washed with dH<sub>2</sub>O. The leaves were cut into small pieces and placed in 20 ml dH<sub>2</sub>O for 20 min to allow release of bacterial cells into the solution. The resuspended bacterial solution was centrifuged at 13,000 rpm and the bacterial cell pellet was stored at -80 °C.

### RNA extraction and first strand cDNA synthesis

Bacterial total RNA was isolated using RNX-Plus kit (CinnaGen Inc., Tehran, Iran) according to the manufacturer's instructions. The RNA was treated with DNase I, RNase-free (Fermentas GmbH., Opeltstrasse, Germany) and its integrity and quantity was evaluated via 1.6% (w/v) agarose gel electrophoresis and spectrophotometry, respectively. To check for DNA contamination, melting curve was generated by real-time PCR using RNA as template. First strand cDNA was synthesized according to Gonzalez and Robb (2007) using 1 µg RNA. First strand cDNA was made with four separate hexamers (ATTTTT, GTTTTT, TGTTTT, and TTTTTT). The synthesized cDNA for each hexamer was mixed with others at the same ratio and called mixed-primer cDNA which was used in PCR as template.

### Polymerase chain reaction

The PCR reaction contained 1 × PCR buffer, 1.5 mM MgCl<sub>2</sub>, 0.2 mM dNTPs, 200 pmol mixed forward and reverse primers, 1.25 U *Taq* DNA polymerase (CinnaGen Inc., Tehran, Iran) and 2.5

μl cDNA as template. The primer pairs (100 nmoles each) used for each amplicon are listed in Table 1 and designed using Primer3 (Rozen and Skaletsky, 2000, available at <http://frodo.wi.mit.edu/>). PCR was performed with an initial denaturation for 5 min at 95 °C followed

by 30 cycles of [95 °C for 30 sec; 58 °C for 30 sec; 72 °C for 1 min], followed by 72 °C for 10 min. PCR products were separated on 1% (w/v) agarose gels containing 0.1 mg/ml Ethidium bromide.

**Table 1** General characteristics of gene specific primers with their names and TMs, designed for transcript analysis of candidate genes involved in pathogenicity of *Xanthomonas citri* subsp. *citri*. Candidate genes are categorized based on their *in vivo* expression into the host plant cells.

Candidate gene	Accession No.	Organism	Molecular Function	Specific primers (5'-3')	Expression pattern	Reference
<i>hk</i>	XAC3673	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	Signal transduction	hk-F: CCTACCTGGCCTTCATCCTG hk-R: GCGACAGCCAGTAGCCTGA	Early	Laia et al., 2009
<i>hrpB4</i>	XAC0410	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	PAI-virulence protein HrpB4	hrpB4-F: GGATGGTCGGGATTGTCTG hrpB4-R: CACTGTTCGAGCCAGTCCAG	Early	da Silva et al., 2002
<i>rpfB</i>	XAC1880	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	long-chain fatty acyl CoA ligase	rpfB-F: CGGCAAGACCATCACCTACC rpfB-R: GGGGTGTACAGCGGATTGAC	Early	da Silva et al., 2002
<i>btuB</i>	XAC1310	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	TonB-dependent receptor	btuB-F: CGTCTCCACCGTCAAGGTCT btuB-R: GCTGTAGCCGTCCACATCCT	Late	da Silva et al., 2002
<i>hrpF</i>	XAC0394	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	Translocator of effector proteins	hrpF-F: GGGACAGCAAGATCCTGGAC hrpF-R: GGCCAGCTTCAAGGCAAG	Late	da Silva et al., 2002; Rossier et al., 2000
<i>pthA*</i>	EF473086	<i>Xanthomonas citri</i> pv. <i>citri</i> Strain: Xc270	Avirulence protein	PthA*-F: CATAACAGAGGCTGCCACAGG PthA*-R: GAGCGTGCCTAGATCCACCT	No amplification	Al-Saadi et al., 2007
<i>avrBs2</i>	XAC0076	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	Avirulence protein	avrBs2-F: GAGTTCTCCGAGCAGCGTCT avrBs2-R: CCAGATACACCGGCTCCAG	Constant	da Silva et al., 2002
<i>hrcJ</i>	XAC0409	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	T3SS Membrane protein HrcJ	hrcJ-F: CAGCTGTCGACAGCCTTGTC hrcJ-R: *7*\$*7&&7&&\$&*7*7	Constant	da Silva et al., 2002
<i>clp</i>	XAC0483	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	cAMP-regulatory protein	clp-F: CGGCACCCTCTACTACGTGA clp-R: AGCCGCTCGTAGCTGATCTC	Constant	da Silva et al., 2002
<i>flgE</i>	XAC1983	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	Flagellar biosynthesis hook protein/motor activity	flgE-F: GACCTGCAGTTGCTGACCAC flgE-R: GCCCAGCGAGTCGTAGACAT	Constant	da Silva et al., 2002
<i>hrcC</i>	XAC0415	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	HrcC protein/transporter activity	hrcC-F: CCTGCCTGTACCACAGTCCA hrcC-R: GGTGACTTCTGGCGAGATCC	Constant	da Silva et al., 2002
<i>hrpB2</i>	XAC0408	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	HrpB2 protein	hrpB2-F: GCCGGCACCAGTATTACCTC hrpB2-R: CACGCCATCGTTCTGCAC	Constant	da Silva et al., 2002
<i>hrpE</i>	XAC0397	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	HrpE protein	hrpE-F: GGTGTGTCCGGTGGAATCTC hrpE-R: CAGCGCCACGTTGAAGTC	Constant	da Silva et al., 2002
<i>hrpG</i>	XAC1265	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	two-component response regulator activity	hrpG-F: CCTGGTCTCGCAGGTCAAC hrpG-R: GCCATGCCAGCAGTGAAC	Constant	da Silva et al., 2002
<i>hrpXCT</i>	XAC1266	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	transcription factor activity	hrpXct-F: CTGCTCTACAACCGCTACGC hrpXct-R: GTAGCTGTAGGCGCGTCGAT	Constant	da Silva et al., 2002
<i>tlyC</i>	XAC1709	<i>Xanthomonas axonopodis</i> pv. <i>citri</i> strain 306	hemolysin C	tlyC-F: GTGACCATCAGCGACCTG tlyC-R: GTGTGGTAGTCGCTTCGTC	Constant	da Silva et al., 2002
<i>16S rRNA</i>	DQ660898	<i>Xanthomonas citri</i> strain FB-1314	housekeeping gene	16S-F: GTGGTTCGAGTCCTCCAGA 16S-R: ACTCGAAGTCCCCGCTTAG	Constant	Shaad et al., 2006

## Results

### A number of pathogenicity-related candidate genes were expressed in non-inducible culture medium

Amongst sixteen reported pathogenicity-related genes that were analyzed here, three were constitutively expressed in non-inducible medium (Fig. 1A). Surprisingly, two of these genes (*hk* and *rpfB*) were those involved in quorum sensing and cell to cell communication pathway. The other *in-vitro* responding gene was *btuB*, a gene involved in iron uptake in bacteria. The rest of the genes appeared not to be expressed in non-inducible condition. The 16S rRNA gene was used as reference gene for data normalization.

### Disease-related candidate genes were expressed in a time course-dependent manner

Except for one gene, *pthA*\* whose expression was not detected in this experiment, all other genes appeared to be expressed in different time points after inoculation (Fig. 1B-D). Different patterns of gene expression were noticed. Three candidate genes, *hk*, *hrpB4* and *rpfB* showed an early-responsive pattern of expression. A relatively strong expression immediately after inoculation with no or very weak expression thereafter was demonstrated for these genes. The late-responsive genes to inoculation were *hrpF* and *btuB*. The rest of the candidate genes were constitutively expressed in the course of time including *avrBs2*, *clp*, *flgE*, *hrcJ*, *hrpB2*, *hrpE*, *hrpG*, *hrpXCT*, and *tlyC*. The 16S rRNA expression level was considered to serve as the reference gene.

## Discussion

A relatively precise comprehension of chronological events involved in establishment, pathogenicity, quorum sensing and the regulation of all the steps required for a pathogenic system to function properly, can be achieved via transcript analysis of bacterial key genes. This data may lead to targeted approaches of how, where and when to enforce

plant cells against the invasion of bacterial cells with devastating effects on yield and the quality of produce. A candidate gene approach including 16 *hrp*-related genes, previously reported to be involved in different steps of *Xanthomonas citri* subsp. *citri* (strain 306) pathogenicity (daSilva *et al.*, 2002), was undertaken for the pathotype A\*. The transcript analysis was performed for bacterial cells grown in culture media (uninduced) and cells isolated from leaves of Mexican lime (induced).

In *Xanthomonas citri* subsp. *citri*, *pthA* and its homologs are recognized as genes indispensable for pathogenicity and elicitation of specific symptoms of asiatic citrus canker disease (Swarup *et al.*, 1992; Al-Saadi *et al.*, 2007). However, using gene specific primers designed based on the A\*-type strain (Xc270) sequence failed to produce any amplicons with either cDNAs in both uninduced and induced conditions (Fig. 1A and 1C) or genomic DNA (data not shown). The *pthA* and its homologs, *pthA*\*, *pthA*<sup>w</sup>, *pthB* and *pthC* are known to be involved in pathogenicity response of *Xcc* (Al-Saadi *et al.*, 2007). These genes are members of *avrBs2* gene family responsible for host specificity and avirulence of plant bacterial pathogens (Gurlebeck *et al.*, 2006). It has been demonstrated that alteration in the nucleotide sequence of the members of this gene family causes failure in specific host recognition (Yang *et al.*, 2005). Thus, sequence conversions are probable between and within pathotypes as we have reported earlier (Soltaninejad, *et al.*, 2010).

Bacterial cells use quorum sensing, a way of communicating between cells for the regulation of diverse arrays of physiological activities, such as biofilm formation, competence, conjugation, antibiotic production, motility, sporulation, symbiosis, virulence, and pathogenicity (Miller and Bassler, 2001). Although for such cooperative actions a local threshold cell density is required (Diggle *et al.*, 2007). Similarly, *Xcc* utilizes the system for the regulation of xanthan (EPS) biosynthesis, gene expression, motility, adaptation, and bacterial virulence (Laia *et al.*, 2009). Amongst many genes involved in this biological process, two

genes were transcriptionally analyzed; *hk*, histidine kinase sensor which is a member of two-component signal transduction systems, and *rpfB*, involved in the production of a diffusible signal factor.

Prokaryotes and *Xanthomonas* use HK and a response regulator (RR) to sense and respond to the surrounding stimuli (Qian *et al.*, 2008; Laia *et al.*, 2009), suggesting a possible role in quorum sensing (Neiditch and Hughson, 2007). In our study, *hk* was expressed in uninduced medium and further analysis on Mexican lime demonstrated its expression in the first few days of inoculation and its disappearance in the following days (Fig. 1B). Our findings were in agreement with the previously reported role of HK to be a bacterial sensor, sensing the neighboring milieu during quorum sensing until bacterial cells reach a certain population density.

*rpfB*, a member of gene cluster known as *rpf* family, is part of the diffusible signal factor type quorum sensing with a critical role in regulating the production of virulence factor (Barber *et al.*, 1997; He *et al.*, 2007). Similar to *hk*, *rpfB* was expressed in the first day post inoculation, but stronger than *HK* and disappeared in the following days (Fig. 1B). The early expression pattern of *rpfB* gene in the first day post inoculation may reflect its pivotal role in pathogenicity process of *Xcc-A\** and its necessity for expression of the other genes involved in a successful compatible interaction of the bacterium with its proper host plant. Similarly, *rpfB* was expressed in uninduced condition (Fig. 1A). It can be concluded that this gene amongst many others is involved in bacterial colonization and host infection, if the environmental conditions are in favor of the bacterial cells.

Within the late expressing genes, *hrpF* showed an expression near to nil in first two days and a blast in expression in the following days (Fig. 1C). This data is in agreement with the reported function of *hrpF* as being protein terminator of the injectisome and translocator of effector proteins into the host plant cytoplasm (Rossier *et al.*, 2000).

In addition to the effector proteins that are directly involved in pathogenicity, other gene products are necessary to weaken host defense mechanisms. These may induce signaling molecules and specifically reactive oxygen species (ROS) (Fones and Preston, 2011). One of such genes is *btuB* that may indirectly suppress the ROS action in host cell. Bacterial cells use superoxide dismutases (SODs) for ROS detoxification. Some of which are iron dependent dismutases (FeSOD); failing to act in iron deficient environments (Kim *et al.*, 1999). This highlights the critical role of iron in pathogenicity (Pandey and Sonti, 2010), and the need of bacterial cells to have a rather sophisticated machinery for iron uptake. A gene with a pivotal role in this system is *btuB* that through insertional mutation analysis, an iron uptake deficiency in *Xanthomonas campestris* pv. *campestris* was demonstrated (Qian *et al.*, 2005). Here, analysis of *btuB* expression pattern revealed that its expression starts at day 3 post inoculation and increases between days 4 to 7 (Fig. 1C). The surge in later days may infer the time that the bacterial cell density has passed a certain limit that has turned on the ROS metabolic pathways in the host that is needed to be detoxified by iron-dependent dismutases such as BtuB. Furthermore, it has been noted that *btuB* is expressed in uninduced media (Fig. 1A). Nutritional deficiency is an inevitable consequence of bacterial growth in the medium. In these situations, iron uptake system is activated to overcome the iron deficiency in the media. The next step in the hierarchy of pathogenesis, would be to introduce the regulatory genes for both providing bacterial survival and enforcing proper infection by the bacteria. The controlling genes can be divided into transcription factors, genes involved in plant-bacteria interactions and quorum sensing.

Some of the structural genes, involved in this study, including *hrcJ*, *hrpB2* (Schulz and Buttner, 2011), and *hrpE* (Weber and Koebnik, 2005) demonstrated constant low expression, while *hrcC* expression was high during the first week of induction (Fig. 1D). It is apparent that the gene products that are being used as the

core components of injectisome (Schulz and Buttner, 2011) are expressed all the way through the establishment of the structure. Therefore, a low continuous expression suffices the bacterial need. In contrast, HrcC is an oligomeric protein complex, comprising the outer part of the core (Deng and Huang, 1999; Lin *et al.*, 2006), and due to its polymeric nature a constant high expression is required by bacterial cells (Fig. 1D). Interestingly in contrast, *hrpB4* was expressed only one day after inoculation (Fig. 1D). This different expression pattern may be due to its role in the formation of injectisome. HrpB4 is one of the main building blocks of the core that perhaps is needed early in the structural formation of the basal parts of the injectisome. Nevertheless, this hypothesis needs to be tested via *in situ* localization studies. Eventually, by finalizing the injectisome structure it is time for some of the effector proteins to be biosynthesized and be translocated into the host cytoplasm via the tube-like structure. It is worth stating that some of the effectors keep synthesizing all the way through, even before the injectisome construction is completed. The constant low expression of these effector proteins seems to be significant in signaling to the virulence apparatus that the bacterial cells have approached the host and further might have a vital role in keeping virulence machinery turned on. These hypotheses need to be further confirmed by mutant analyses of the genes followed by determining any behavioral changes in bacterial pathogenicity.


Clp (cAMP receptor-like protein) as a global transcription factor, controls the expression of a subset of genes involved in pathogenesis in xanthomonads (Leduc and Roberts, 2009) and its regulatory function is apparent from its constitutive expression (Fig. 1D).

The secretion of effector proteins need to be tightly regulated by regulatory elements known as *HrpG* and *HrpX* regulons (Yamazaki *et al.*, 2008; Guo *et al.*, 2011). A recent global analysis of transcripts revealed that these two regulons are vital for the pathogenicity of *Xcc* (Guo *et al.*, 2011).

*HrpG* and *HrpX* regulons control all 24 T3SS secretion system genes, 23 T3SS effector genes, and 29 type II secretion system substrate genes. Furthermore, these two regulons adjust the bacterial behavior in response to the host environment by altering a range of metabolic pathways including amino acid biosynthesis; oxidative phosphorylation; pentose-phosphate pathway; transport of sugar, iron, and potassium; and phenolic catabolism (Guo *et al.*, 2011). Our data about the constant low expression pattern of *hrpG* and *hrpXCT* conforms to their global regulatory and signaling roles.

Among the constitutively expressed genes, *avrBs2* and *tlyC* (Astua-Monge *et al.*, 2005) showed low expressions from day 1 inferring that these genes belong to the second class with a signaling role (Fig. 1D).

The last gene was *flgE* (hook) that is involved in flagellum motility and biofilm formation. In the absence of flagellin the pathogenicity of *Xanthomonas* is reduced (Malamud *et al.*, 2011). *flgE* had a constant expression pattern (Fig. 1D) indicating that the studied *Xcc*-A\* strain may require its flagellum to cause disease. In conclusion, in this study we took advantage of RT-PCR technology to find elements of regulatory interconnections between genes involved in pathogenicity of *Xanthomonas citri* subsp. *citri* type A\*. The dynamics of injectisome formation was also elucidated. This research demonstrated that pathogenicity system in the NIGEB-088 A\*-type strain of *Xcc*, is activated upon bacterial infiltration into the host plant leaf cells in the first day after inoculation. The expression of *hrpF* gene and translocon synthesis on the fourth day post inoculation shows eventually that injectisome is completed after four days and that the T3SS of the given strain is apparently ready to secrete protein effectors into the host plant cells to trigger pathogenicity reaction. By including more candidate genes and using a more sensitive method of gene expression analysis such as real time PCR, it would be possible to find more regulatory elements in the pathogenicity network.



**Figure 1** *In vitro* and *in vivo* transcript analysis of sixteen candidate genes involved in pathogenicity of *Xanthomonas citri* subsp. *citri*. Uninduced condition: A) Candidate genes that expressed in uninduced condition. Amongst 16 candidate genes, RT-PCR demonstrated the expression of three genes in vitro condition, namely *rpfB*, *btuB*, and *hk*. Induced condition: B) The expression pattern of early responsive genes C) The expression pattern of late responsive genes D) The expression pattern of candidate genes with constant expression from day 1-7 after inoculation. The PCR product of reverse transcription was separated on 1% agarose gel. 16S rRNA was used as standard. dai: days after inoculation.

### Acknowledgement

This work was performed as part of a National Institute for Genetic Engineering and Biotechnology (NIGEB) grant (NIGEB-369) for controlling and management of Asiatic Citrus Canker disease.

### References

Alfano, J. R. and Collmer, A. 1997. The type III (Hrp) secretion pathway of plant pathogenic bacteria: trafficking harpins, Avr proteins and death. *Journal of Bacteriology*, 179: 5655-5662.

Al-Saadi, A., Reddy, D., Duan, Y. P., Brunings, A. M., Yuan, Q. and Gabriel, D. W. 2007. All five host-range variants of *Xanthomonas citri* carry one *pthA* homolog with 17.5 repeats that determines pathogenicity on citrus, but none determine host-range variation. *Molecular Plant Microbe Interaction*, 20: 934-943.

Astua-Monge, G., Freitas-Astua, J., Bacocina, G., Roncoletta, J., Carvalho, S. A. and Machado M. A. 2005. Expression profiling of virulence and pathogenicity genes of *Xanthomonas axonopodis* pv. *citri*. *Journal of Bacteriology*, 187: 1201-1205.

- Barber, C. E., Tang, J. L., Feng, J. X., Pan, M. Q., Wilson, T. J., Slater, H., Dow, J. M., Williams, P. and Daniels, M. J. 1997. A novel regulatory system required for pathogenicity of *Xanthomonas campestris* is mediated by a small diffusible signal molecule. *Molecular Microbiology*, 24: 555-566.
- Bock, C. H., Graham, J. H., Gottwald, T. R., Cook, A. Z. and Parker, P. E. 2010. Wind speed effects on the quantity of *Xanthomonas citri* subsp. *citri* dispersed downwind from canopies of grapefruit trees infected with citrus canker. *Plant Disease*, 94: 725-736.
- Brunings, A. M. and Gabriel, D. W. 2003. *Xanthomonas citri*: breaking the surface. *Molecular Plant Pathology*, 4: 141-157.
- Bui Thi Ngoc, L., Verniere, C., Jouen, E., Ah-You, N., Lefeuvre, P., Chiroleu, F., Gagnevin, L. and Pruvost, O. 2010. Amplified fragment length polymorphism and multilocus sequence analysis-based genotypic relatedness among pathogenic variants of *Xanthomonas citri* subsp. *citri* and *Xanthomonas campestris* pv. *bilvae*. *International Journal of Systematic and Evolutionary Microbiology*, 60: 515-525.
- Buttner, D. and Bonas, U. 2002. Getting across-bacterial type III effector proteins on their way to the plant cell. *EMBO Journal*, 21: 5313-5322.
- Cornelis, G. R. 2006. The type III secretion injectisome. *Nature Review of Microbiology*, 4: 811-825.
- Cubero, J. and Graham, J. H. 2002. Genetic relationship among worldwide strains of *Xanthomonas* causing canker in citrus species and design of new primers for their identification by PCR. *Applied and Environmental Microbiology*, 68: 1257-1264.
- da Silva, A. C., Ferro, J. A., Reinach, F. C., Farah, C. S., Furlan, L. R., Quaggio, R. B., Monteiro-Vitorello, C. B., Van Sluys, M. A., Almeida, N. F., Alves, L. M., do Amaral, A. M., Bertolini, M. C., Camargo, L. E., Camarotte, G., Cannavan, F., Cardozo, J., Chamberg, F., Ciapina, L. P., Cicarelli, R. M., Coutinho, L. L., Cursino-Santos, J. R., El-Dorry, H., Faria, J. B., Ferreira, A. J., Ferreira, R. C., Ferro, M. I., Formighieri, E. F., Franco, M. C., Greggio, C. C., Gruber, A., Katsuyama, A. M., Kishi, L. T., Leite, R. P., Lemos, E. G., Lemos, M. V., Locali, E. C., Machado, M. A., Madeira, A. M., Martinez-Rossi, N. M., Martins, E. C., Meidanis, J., Menck, C. F., Miyaki, C. Y., Moon, D. H., Moreira, L. M., Novo, M. T., Okura, V. K., Oliveira, M. C., Oliveira, V. R., Pereira, H. A., Rossi, A., Sena, J. A., Silva, C., de Souza, R. F., Spinola, L. A., Takita, M. A., Tamura, R. E., Teixeira, E. C., Tezza, R. I., Trindade dos Santos, M., Truffi, D., Tsai, S. M., White, F. F., Setubal, J. C. and Kitajima, J. P. 2002. Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature*, 417: 459-463.
- Deng, W. L. and Huang, H. C. 1999. Cellular locations of *Pseudomonas syringae* pv. *syringae* HrcC and HrcJ proteins, required for harpin secretion via the Type III Pathway. *Journal of Bacteriology*, 181: 2298-2301.
- Diggle, S. P., Gardner, A., West, S. A. and Griffin, A. S. 2007. Evolutionary theory of bacterial quorum sensing: when is a signal not a signal? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362: 1241-1249.
- Fones, H. and Preston, G. M. 2011. Reactive oxygen and oxidative stress tolerance in plant pathogenic *Pseudomonas*. *FEMS Microbiology Letters*: doi: 10.1111/Journal of 1574-6968.2011.02449.x.
- Gonzalez, J. M. and Robb, F. T. 2007. Counterselection of prokaryotic ribosomal RNA during reverse transcription using non-random hexameric oligonucleotides. *Journal of Microbiological Methods*, 71: 288-291.
- Guo, Y., Figueiredo, F., Jones, J. and Wang, N. 2011. HrpG and HrpX play global roles in coordinating different virulence traits of *Xanthomonas axonopodis* pv. *citri*. *Molecular Plant Microbe Interaction*, 24: 649-661.



- He, Y. W., Ng, A. Y., Xu, M., Lin, K., Wang, L. H., Dong, Y. H. and Zhang, L. H. 2007. *Xanthomonas campestris* cell-cell communication involves a putative nucleotide receptor protein Clp and a hierarchical signaling network. *Molecular Microbiology*, 64: 281-292.
- Iwamoto, M. and Oku, T. 2000. Cloning and molecular characterization of *hrpX* from *Xanthomonas axonopodis* pv. *citri*. *DNA Sequence*, 11: 167-173.
- Kanamori, H. and Tsuyumu, S. 1998. Comparison of nucleotide sequence of canker-forming and non-canker-forming *pthA* homologues in *Xanthomonas campestris* pv. *citri*. *Annals of the Phytopathological Society of Japan*, 64: 462-470.
- Kay, S. and Bonas, U. 2009. How *Xanthomonas* type III effectors manipulate the host plant. *Current Opinion in Microbiology*, 12: 37-43.
- Kim, Y. C., Miller, C. D. and Anderson, A. 1999. Transcriptional regulation by iron and role during plant pathogenesis of genes encoding iron- and manganese-superoxide dismutases of *Pseudomonas syringae* pv. *syringae* B728a. *Physiological and Molecular Plant Pathology*, 55: 327-339.
- Laia, M. L., Moreira, L. M., Dezajacomo, J., Brigati, J. B., Ferreira, C. B., Ferro, M. I. T., Silva, A. C. R., Ferro, J. A. and Oliveira, J. C. F. 2009. New genes of *Xanthomonas citri* subsp. *citri* involved in pathogenesis and adaptation revealed by a transposon-based mutant library. *BMC Microbiology*, 9: 12.
- Leduc, L. and Roberts, G. P. 2009. Cyclic di-GMP allosterically inhibits the CRP-like protein (Clp) of *Xanthomonas axonopodis* pv. *citri*. *Journal of Bacteriology*, 191: 7121-7122.
- Lin, Y. C., Hsu, S. T. and Huang, H. C. 2006. Evidence for the interaction between HrpT and HrcC of *Pseudomonas syringae* pv. *syringae* 61. *Plant Pathology Bulletin*, 15: 171-185.
- Malamud, F., Torres, P. S., Roeschlin, R., Rigano, L. A., Enrique, R., Bonomi, H. R., Castagnaro, A. P., Marano, M. R. and Vojnov, A. A. 2011. The *Xanthomonas axonopodis* pv. *citri* flagellum is required for mature biofilm and canker development. *Microbiology*, 157: 819-829.
- Mehta, A. and Rosato, Y. B. 2003. A simple method for in vivo expression studies of *Xanthomonas axonopodis* pv. *citri*. *Current Microbiology*, 47: 400-403.
- Miller, M. B. and Bassler, B. L. 2001. Quorum sensing in bacteria. *Annual. Review of Microbiology*, 55: 165-199.
- Neiditch, M. B. and Hughson, F. M. 2007. The regulation of histidine sensor kinase complexes by quorum sensing signal molecules. *Methods of Enzymology*, 423: 250-263.
- Pandey, A. and Sonti, R. V. 2010. Role of the FeoB protein and siderophore in promoting virulence of *Xanthomonas oryzae* pv. *oryzae* on rice. *Journal of Bacteriology*, 192: 3187-3203.
- Qian, W., Han, Z. J., Tao, J. and He, C. 2008. Genome-scale mutagenesis and phenotypic characterization of two-component signal transduction systems in *Xanthomonas campestris* pv. *campestris* ATCC 33913. *Molecular Plant Microbe Interaction*, 21: 1128-1138.
- Qian, W., Jia, Y., Ren, S. X., He, Y. Q., Feng, J. X., Lu, L. F., Sun, Q., Ying, G., Tang, D. J., Tang, H., Wu, W., Hao, P., Wang, L., Jiang, B. L., Zeng, S., Gu, W. Y., Lu, G., Rong, L., Tian, Y., Yao, Z., Fu, G., Chen, B., Fang, R., Qiang, B., Chen, Z., Zhao, G. P., Tang, J. L. and He, C. 2005. Comparative and functional genomic analyses of the pathogenicity of phytopathogen *Xanthomonas campestris* pv. *campestris*. *Genome Research*, 15: 757-767.
- Rezaei, M. K., Shams-Bakhsh, M. and Alizadeh, A. 2012. Genetic diversity among *Xanthomonas citri* subsp. *citri* in Iran. *Journal of Plant Protection Research*, 51: 1-9.
- Rossier, O., Van den Ackerveken, G. and Bonas, U. 2000. HrpB2 and HrpF from *Xanthomonas* are type III-secreted proteins and essential for pathogenicity and

- recognition by the host plant. *Molecular Microbiology*, 38: 828-838.
- Rozen, S. and Skaletsky, H. J. 2000. Primer3 on the WWW for general users and for biologist programmers, In: Krawetz S, Misener S (Eds), *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa, NJ, pp 365-386.
- Ryan, R. P., Vorholter, F. J., Potnis, N., Jones, J. B., Van Sluys, M. A., Bogdanove, A. J. and Dow, J. M. 2011. Pathogenomics of *Xanthomonas*: understanding bacterium-plant interactions. *Nature Review of Microbiology*, 9: 344-355.
- Schaad, N. W., Postikova, E., Lacy, H., Sechler, A., Agarkova, I., Stromberg, E., Stromberg, K. and Vidaver, K. 2005. Reclassification of *Xanthomonas campestris* pv. *citri* (ex Hasse 1915) Dye 1978 forms A, B, C, D, and E as *X. smithii* subsp. *citri* (ex hasse) sp. Nov. nom. Rev. comb. nov., *X. fuscans* subsp. *Aurantifolli* (ex Gabriel 1989) sp. nov. nom. rev. comb. nov., and *X. alfalfae* subsp. *citrumelo* (ex Riker and Jones) Gabriel *et al.*, 1989 sp. nov. nom. rev. comb. nov.; *X. campestris* pv. *malvacearum* (ex Smith 1901) dye 1978 as *X. campestris* subsp. *smithii* nov. comb. nov. nom. nov.; *X. campestris* pv. *alfalfae* (ex Riker *et al.*, 1935) sp. Nov. nom. Rev.; and "var. *fuscans*" of *X. campestris* pv. *phaseoli* (ex Smith, 1987) Dye 1978 as *X. fuscans* subsp. *fuscans* sp. nov. Syst. Applied Microbiology, 28: 494-518.
- Schulz, S. and Buttner, D. 2011. Functional characterization of the type III secretion substrate specificity switch protein HpaC from *Xanthomonas campestris* pv. *vesicatoria*. *Infection and Immunity*, 79: 2998-3011.
- Shiotania, H., Uematsub, H., Tsukamotob, T., Shimizub, Y., Uedab, K., Mizunob, A. and Sato, S. 2009. Survival and dispersal of *Xanthomonas citri* subsp. *citri* from infected Satsuma mandarin fruit. *Crop Protection*, 28: 19-23.
- Soltaninejad, H., Parchami, N., Alavi, S. M. and Ghasemi, A. 2010. A narrow genetic variation is detectable among Iranian strains of *Xanthomonas citri* subsp. *citri*, causal agent of bacterial citrus canker. *Proceedings of 12<sup>th</sup> International Conference on Plant Pathogenic Bacteria in Reunion Island, France*, p. 56.
- Swarup, S., Yang, Y., Kingsley, M. T. and Gabriel, D. W. 1992. An *Xanthomonas citri* pathogenicity gene, *pthA*, pleiotropically encodes gratuitous avirulence on non-hosts. *Molecular Plant Microbe Interaction*, 5: 204-213.
- Traoré, Y. N., Ngoc, L. B. T., Vernière, C. and Pruvost, O. 2008. First report of *Xanthomonas citri* subsp. *citri* causing citrus canker in Mali. *Plant Disease*, 92: 977.
- Verniere, C., Hartung, J. S., Pruvost, O., Civerolo, E. L., Alvarez, A. M., Maestri, P. and Luisetti, J. 1998. Characterization of phenotypically distinct strains of *Xanthomonas axonopodis* pv. *citri* from Southwest Asia. *European Journal of Plant Pathology*, 104: 477-488.
- Vojnov, A. A., do Amaral, A. M., Dow, J. M., Castagnaro, A. P. and Marano, M. R. 2010. Bacteria causing important diseases of citrus utilize distinct modes of pathogenesis to attack a common host. *Applied Microbiology and Biotechnology*, 87: 467-477.
- Weber, E. and Koebnik, R. 2005. Domain structure of HrpE, the Hrp pilus subunit of *Xanthomonas campestris* pv. *vesicatoria*. *Journal of Bacteriology*, 187: 6175-6186.
- Yamazaki, A., Hirata, H. and Tsuyumu, S. 2008. Type III regulators HrpG and HrpXct control synthesis of  $\alpha$ -amylase, which is involved in in planta multiplication of *Xanthomonas axonopodis* pv. *citri*. *Journal of General Plant Pathology*, 74: 254-257.
- Yang, B., Sugio, A. and White, F. F. 2005. Avoidance of host recognition by alterations in the repetitive and C-terminal regions of AvrXa7, a type III effector of *Xanthomonas oryzae* pv. *oryzae*. *Molecular Plant Microbe Interaction*, 18: 142-149.

## آنالیز ترانسکریپت بعضی از عناصر دخیل در بیماری‌زایی یک جدایه ایرانی از تیپ بیماری‌زای A\* باکتری *Xanthomonas citri* subsp. *citri*

رعنا روحانی<sup>۱،۲</sup>، مسعود سلطانی نجف‌آبادی<sup>۳</sup>، سید مهدی علوی<sup>۴\*</sup>، ناصر فرخی<sup>۴</sup> و مسعود شمس‌بخش<sup>۲</sup>

- ۱- گروه بیوتکنولوژی گیاهی، پژوهشگاه ملی مهندسی ژنتیک و زیست فناوری، صندوق پستی ۱۴۱۵۵-۶۳۴۳، تهران، ایران
  - ۲- گروه بیماری‌شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران
  - ۳- بخش دانه‌های روغنی، مؤسسه تحقیقات اصلاح نهال و بذر، کرج، ایران
  - ۴- گروه زراعت و اصلاح نباتات، دانشکده مهندسی کشاورزی، دانشگاه صنعتی شاهرود، صندوق پستی ۳۶۱۹۹۹۵۱۶، شاهرود، ایران
- \* پست الکترونیکی نویسنده مسئول مکاتبه: mealavi@nigeb.ac.ir, shamsbakhsh@modares.ac.ir  
دریافت: ۲۳ اردیبهشت ۱۳۹۱؛ پذیرش: ۸ آذر ۱۳۹۱

**چکیده:** شانکر آسیایی مرکبات از جمله بیماری‌های مخربی است که زیان‌های سنگینی بر اقتصاد کشاورزی کلیه کشورهای تولیدکننده مرکبات در جهان وارد می‌سازد. در بین سه نوع مختلف بیماری (A، A\* و A\*)، انواع A\* از نظر ژنتیکی کمتر مطالعه شده‌اند. به‌منظور درک صحیح رفتار باکتری عامل شانکر آسیایی مرکبات سویه A\* (*Xanthomonas citri* subsp. *citri*) در مواجهه با گیاه میزبان، از روش RT-PCR استفاده شد. به همین منظور، بیان ۱۶ ژن کاندیدا دخیل در مراحل مختلف بیماری‌زایی این باکتری در محیط کشت (شرایط غیرالقایی) و در مواجهه با گیاه لیموترش (*Citrus aurantifolia* L.) (شرایط القایی) با یکدیگر مقایسه شدند. نتایج حاکی از بیش‌بانی برخی ژنها در شرایط القایی نسبت به حالت غیرالقایی بود که خود بیانگر برهمکنش باکتری و میزبان است. از این‌رو بیان ژن‌های مورد مطالعه در پاسخ به آلودگی حاصل از جدایه مذکور را می‌توان در سه کلاس دائمی، زود-پاسخ و دیر-پاسخ طبقه‌بندی نمود.

**واژگان کلیدی:** Xcc، RT-PCR، شانکر مرکبات، لیموی شیراز