

Some mesostigmatic mites from Iran with their world distribution

Fariba Kadkhodae Eliaderani¹, Alireza Nemati^{2*} and Farhan Kocheili¹

1. Department of Plant Protection, Faculty of Agriculture, Shahid Chamran University of Ahvaz, Iran.

2. Department of Plant Protection, Faculty of Agriculture, University of Shahrekord, Iran.

Abstract: Edaphic Mesostigmata are important, because these feed on arthropods and other invertebrates which may affect natural equilibrium of other living micro-organisms. In order to study the fauna of mesostigmatic mites, samples of soil and litter were collected from different parts of Esfahan region. Mites were extracted by means of Berlese funnel, cleared in Lactic acid and mounted in Hoyer's medium. In this research, the world and Iranian distribution of species with some information about their habitats are provided. Two genera and 3 species were recorded as new records for Iran mite fauna that are marked by an asterisk. The list of identified genera and species is as follows:

Ascidae: *Arctoseius cetratus* (Sellnick, 1940), *A. pristinus** Karg, 1962, *A. venustus* (Berlese, 1917), *Gamasellodes bicolor* (Berlese, 1918); **Blattisociidae:** *Lasioseius sugawarai* Ehara, 1964, *L. youcefi* Athias-Henriot, 1959; **Melicharidae:** *Proctolaelaps pygmaeus* (Muller, 1860); **Parholaspididae:** *Proparholaspulus angustatus** Ishikawa, 1987 **Trachytidae:** *Uroseius traegardhi** (Hirschmann & Zirngiebl-Nicol, 1969).

Keywords: Acari, Mesostigmata, Soil, World distribution, Iran

Introduction

The Mesostigmata are a group of mites (Acari) comprising a great diversity of vertebrate parasites-symbionts of insects, and for the most part, free-living predators. Soil, litter, plants, dung and decaying wood are all inhabited by a range of free-living mesostigmatic mites (Walter *et al.*, 1998; Halliday, 2000; Beaulieu and weeks, 2007). The order mesostigmata is divided into three suborders. The suborder Monogynaspida includes a multitude of free-living and parasitic species that occur in countless terrestrial, marine and freshwater habitats throughout the world. This suborder includes five cohorts. Gamasina cohort comprises most of the described species of

mesostigmata. The Gamasina is considered to comprise four subcohorts and 10 superfamilies (Lindquist *et al.*, 2009).

Mites of the family Ascidae are commonly found in soil and litter, among other habitats, where they may play an important role as predators of other mites, other arthropods and nematodes (Halliday *et al.*, 1998; Lindquist *et al.*, 2009; Walter, 1988; Walter *et al.*, 1993). They are also among the gamasine mites which are useful as biological indicators of soil conditions and changes in soil ecosystems (Karg and Freier, 1995; Minero *et al.*, 2009). Based on Krantz and Walter (2009) the family Ascidae includes two subfamilies; the Arctoseiinae and the Ascinae and the melicarines, blattisociines and platyseiiines are excluded from this family. The Melicharidae has been classified as a subfamily or a tribe in previous accounts, but now it is treated as a separate family.

Many species of melicharids are predatory, others have adopted to feeding on fungi, pollen,

Handling Editor: Dr. Mohammad Khanjani

*Corresponding author, e-mail: nemati.alireza@agr.sku.ac.ir

Received: 21 January 2013; Accepted: 27 February 2013

and nectar, and one is reported to be parasitic on cockroaches (Lindquist *et al.*, 2009).

The Blattisociidae is a diverse group that has adopted to a broad spectrum of terrestrial, arboreal, and subaquatic habitats. This family includes two subfamilies, the Platyseinae and Blattisociinae. The family Parholaspididae comprises 11 genera of moderately to heavily sclerotized predatory mites that often are found in forest litter, soil, decaying vegetable matter, moss, and tree holes (Lindquist *et al.*, 2009). There are sporadic and in some case incomplete and poor information about the fauna of these families in Iran and also world distribution of their members. In this paper data is provided for some species that have been collected in this survey.

Materials and Methods

Materials were collected from different microhabitats in some parts of Esfahan region during 2011-2012. Mites were extracted from the soil and litter samples by use of Berlese funnels. The specimens were collected in beakers of 75% ethanol, and then fixed on permanent slides in Hoyer's medium. The list of species is presented by different families and related genera, including information about diagnosis, world distribution, records in Iran with new records in this survey. Species' microhabitat and the details of the findings encompass date of collection, locality, microhabitats or hosts that were recorded.

Results

Family Ascidae

Genus *Arctoseius* Thor, 1930

Arctoseius cetratus (Sellnick, 1940)

Diagnosis. Setae *J2*, *Z1*, *Z2*, and *S1*, *S2* short, not reaching the bases of following setae (Gwiazdowicz, 2007). Posterior half of dorsal shield with longitudinal garland of puncta between *J* and *Z* setae, peritremes extending anteriorly to mid-level of coxa II (Halliday *et al.*, 1998).

World distribution. North America (Chant, 1963); Europe (Bregetova, 1977; Karg, 1993; Salmane and Heldt, 2001; Kaluz and Fenda, 2005); Western Europe (Bregetova, 1977); Russia: soil (Petrova, 1982; Makarova, 2000); Western Siberia (Davydova and Nikolsky, 1986); Middle Ural (Russia): fallow land (Khomjakov, 1988); Australia: dung-baited, pitfall trap, New South Wales: cow dung, Victoria: compost (Halliday *et al.*, 1998); Poland: nest of Greater spotted Eagle (*Aquila clanga*) (Gwiazdowicz *et al.*, 2000), soil (Seniczak *et al.*, 2000), rotten wood, leaf litter and other microhabitats (Gwiazdowicz and Klemt, 2004), sod, stuffing from nests of (*Larus ridibundas*) (Gwiazdowicz and Kmita, 2004), litter, rot wood, moss (Gwiazdowicz and Matysiak, 2004), litter, rotting wood, sod, meadow, bark beetles galleries (Gwiazdowicz, 2007), Nests of formicidae (*Formica polyctena*), (*Lasius niger*) (Gwiazdowicz, 2008a), associated with Scolytidae (Gwiazdowicz, 2008b), soil (Gabrys *et al.*, 2008); Sakhalin and Kuril Islands: manure (Marchenko and Mapyehko, 2002); Slovakia: various habitats (Kaluz and Fenda, 2005), nests of mound-building mouse (*Mus spicilegus*) (Masan and Stanko, 2005), birds nests (Fenda and Schniererova, 2010); Central Argentina: Soil (Bedano and Ruf, 2007); Northern Taiga (Makarova, 2009); Finland: pine forest, seashore, meadow, dry meadow, ant hills, compost, dung (Huhta *et al.*, 2010); Saudi Arabia: fruit and timber trees (Al-Atawi, 2011); Austria: roots and soil of plants (Wissuwa *et al.*, 2012).

Iran Records. Following Nemati *et al.* (2012a).

New Records. Soil of different parts of Esfahan: 31.iii.2012, Abshar street; 21.iii.2011, 12.iii.2012, Bagh-daryache street; 28.iii.2011, Bishe-habib park; 10.iii.2012, Khayam street; 25.iii.2011, Khomeinishahr; 03.iii.2011, Khoram street; 28.iii.2011, Marnan park; 04.iv.2012, Mirzataher street; 12.iii.2012, 11.iv.2012, Moshtagh street; 29.iii.2011, Nazhvan park; 25.iii.2012, Ordibehesht street; 06.iv.2012, Takhtpolad graveyard.

***Arctoseius pristinus* Karg, 1962**

Diagnosis. All of the dorsal setae short, tines of epistome with fine, often barely distinguishable branching, genital shield not widened posteriorly (Bregetova, 1977).

World distribution. Central Europe (Bregetova, 1977; Karg, 1993; Kaluz and Fenda, 2005); Slovakia (Kaluz and Fenda, 2005); Finland: alpine, compost, dung (Huhta et al., 2010).

New Records. Soil of different parts of Esfahan: 17.iii.2012, 4.iv.2012, Atash street; 12.iii.2012, Bagh-daryache street; 19.iii.2012, Bostan kodak park; 19.iii.2012, Khayam street; 31.iii.2012, Moshtagh street; 10.iii.2012, 05.viii.2011, 25.iii.2012, Ordibehesht street; 25.iii.2012, Saeb street; 06.iv.2012, Takhtpolad graveyard; 10.iii.2012, 14.iii.2012, Vahid street.

***Arctoseius venustus* (Berlese, 1917)**

Diagnosis. Epistome tri-ramous, margin of epistome pointed, setae Z5 same length as J5 or slightly longer, setae in row R simple, pair of presternal plates present (Gwiazdowicz, 2007).

World distribution. Switzerland (Airoldi et al., 1989); Europe (Karg, 1993; Kaluz and Fenda, 2005); England and Wales: associated with yew (Skorupski and Luxton, 1998); Slovakia (Kaluz and Fenda, 2005), nests of mound-building mouse (*Mus spicilegus*) (Masan and Stanko, 2005); Austria: roots and soil of plants (Wissuwa et al., 2012).

Iran Records. Following Nemati et al. (2012a).

New Records. Soil of different parts of Esfahan: 21.iii.2011, Atash street; 19.iii.2011, Bostan kodak park; 3.iii.2011, Khoram street; 27.iii.2011, Marnan bridge; 24.iv.2011, Saeb street.

Genus *Gamasellodes* Athias-Henriot, 1961***Gamasellodes bicolor* (Berlese, 1918)**

Diagnosis. Ventral idiosoma with a ventrianal shield with three circum-anal setae and at least one pair of ventrianal setae, peritreme long, extending to coxa I, humeral setae r3 of approximately same length as other setae on podonotal shield (Gwiazdowicz, 2007). Anterior margin of ventrianal shield broadly rounded, shield

widest along its middle one-third, at level near anus (Halliday et al., 1998).

World distribution. Tanzania (Hurlbutt, 1971); New York (United states): birds nests (Baker et al., 1976); Norway (Mehl, 1979); Europe (Karg, 1993; Salmane and Heldt, 2001); England and Wales, associated with yew (Skorupski and Luxton, 1998); Latvia: coastal meadow (Salmane, 1999), washed ashore material, dunes (Salmane, 2001a), primary dunes (Salmane, 2001b), associated with *Aphylophorales* fungi from dead or live tree trunks (Salmane, 2005), under the bark, in the frass, rotting wood, mosses and lichens on the tree trunks and stumps (Salmane, 2007); Poland: nests of white-tailed sea Eagle (*Haliaeetus albicila*) (Gwiazdowicz et al., 2000), soil (Seniczak et al., 2000), nests of red-backed shrike (*Lanius collurio*) (Tryjanowski et al., 2001), rotten wood, leaf litter and other microhabitats (Gwiazdowicz and Klemt, 2004), litter, rot wood, moss (Gwiazdowicz and Matysiak, 2004), litter, sod (Gwiazdowicz and Kmita, 2004), litter, fruiting bodies, bark beetles galleries, sod, moss, rotting wood (Gwiazdowicz, 2007), Nests of formicidae (*Formica polyctena*), (*Lasius niger*), (*Formica rufa*), (*Lasius flavus*), (*Lasius fuliginosus*) (Gwiazdowicz, 2008a), associated with Scolytidae (Gwiazdowicz, 2008b), soil (Skorupski et al., 2008), rotting wood (Gwiazdowicz, 2010); South Africa, North and South America, India, Israel (Bhattacharry and Sanyal, 2002); Hungary: soil, litter (Salmane and Kotschan, 2005a), litter (Salmane and Kotschan, 2005b); Central Argentina: soil (Bedano and Ruf, 2007); Alabama: citrus orchards (Fadamiro et al., 2009); Northern Taiga (Makarova, 2009); Finland: pine forest, spruce forest, deciduous forest, seashore, meadow, dry meadow, dead wood, ant hills, compost, dung (Huhta et al., 2010); Slovakia: birds nests (Fenda and Schmiererova, 2010); Turkey: *Canna indica*, *Crocus ancyrensis*, *Crocus biflorus*, *Iris* sp, wild mushroom (Cakmak et al., 2011); Austria: roots and soil of plants (Wissuwa et al., 2012).

Iran Records. Following Nemati *et al.* (2012a).
New Records. Esfahan, soil, 2011.iii.19, Bostan koodak park; 2011.iii.27, Marnan bridge, soil.

Family Blattisociidae

Genus *Lasioseius* Berlese, 1916

Lasioseius sugawarai Ehara, 1964

Diagnosis. Length of $Z3 = 2/3$ the distance between $Z3$ and $S5$, median point of epistome as long as the lateral branches, the three branches as long as wide, macrochaetae on leg $IV = 35$, $Z5 = 47$ (Christian and Karg 2006).

World distribution. Japan (Ehara, 1964), associated with Millipedes (Ishikawa, 1986), (Christian and Karg, 2006), soil, litter (Takaku and Sasaki, 2007).

Iran Records. Following Nemati *et al.* (2012b).

New Records. Esfahan, soil, 13.vii.2011, Marnan park.

Lasioseius youcefi Athias Henriot, 1959

Diagnosis. Holodorsal shield lacking two pairs of setae in row J , some setae on dorsal shield serrate (Gwiazdowicz, 2007). Many of the setae on the dorsal shield and the postanal setae with a short pubescence (Bregetova, 1977).

World distribution. Africa, Southern and Northern Africa (Athias-Henriot, 1959; Karg, 1993); Western Europe, Crimea (Ukraine), Gorkiy oblast, Trans-caucasia, Latvia, Transcarpathia (Bregetova, 1977); North America (Walter and Lindquist, 1989); Europe (Karg, 1993); Latvia: coastal meadow (Salmane, 1999), inland and coastal meadow, spruce stand, bogs, mosses, nests of birds (Salmane, 2001a), soil (Salmane, 2001b), associated with *Aphylophorales* fungi from dead or live tree trunks (Salmane, 2005), under bark and in rotting wood of (*Alnus incana*), (Salmane, 2007); South Africa: clover and Oxalis sp, grass, capeweed cultivated lawn, irrigated pasture (Halliday, 2005); Poland: bark beetles galleries, nests of birds, litter, moss (Gwiazdowicz, 2007); Italy (Plumari, 2009); Austria: roots and soil of plants (Wissuwa *et al.*, 2012).

Iran Records. Following Nemati *et al.* (2012b).

New Records. Soil of different parts of Esfahan: 06.ix.2011, Asgarie street; 19.iii.2011, Bostan koodak park; 21.iii.2011, Kohandej street; 13.vi.2011, Marnan park; 04.ix.2011, Nazhvan park; 04.ix.2011, Ordibehesht street; 06.iv.2012, Takhtpolad graveyard.

Family Melicharidae

Genus *Proctolaelaps* Berlese, 1923

Proctolaelaps pygmaeus (Müller, 1860)

Diagnosis. Anal shield pear-shaped, anus large, $j4$ approximately same length as $z4$, hypostomal $C1$ setae distinctly thicker than other setae on gnathosoma, anterior margin of epistome curved and irregularly denticulate, fixed-digit of chelicerae with 4-7 large proximal teeth and curved row of 3-5 small distal teeth (Gwiazdowicz, 2007; Halliday *et al.*, 1998).

World distribution. North America (Chant, 1963; Karg, 1971); Japan (Ehara, 1964); Europe, South Africa, Indonesia, Australia, New Zealand (Karg, 1971); New Zealand (Spain and Luxton, 1971); Norway: small mammals (Edler and Mehl, 1972), (Mehl, 1979), soil and tree hollows (Slomian *et al.*, 2005); Russia (Bregetova, 1977; Petrova, 1982); Poland (Haitlinger, 1983) (Haitlinger, 1989a,b), nests of greater spotted Eagle (*Aquila clanga*) (Gwiazdowicz *et al.*, 2000), nests of the red-backed shrike (*Lanius collurio*) (Tryjanowski *et al.*, 2001), litter, rotten wood, moss (Gwiazdowicz and Matysiak, 2004), soil, sand dune (Gwiazdowicz and Kmita, 2004), rotten wood, leaf litter and other microhabitat (Gwiazdowicz and Klemm, 2004), nests of white stork (*Ciconia ciconia*) (Bloszyk *et al.*, 2005), nests of birds, rotting wood, litter, fungal fruiting bodies, bark beetles galleries (Gwiazdowicz, 2007), soil (Gabrys *et al.*, 2008), bracket fungi (Gwiazdowicz, 2010); Western Siberia (Davydova and Nikolsky, 1986); Hungary: associated with small mammals (*M. arvalis*) (Ambrose, 1987); Middle Ural (Khomjakov, 1988); Egypt: in the house dust (Shoukry *et al.*, 1990); New South Wales: in bat guano, Queensland: on *Neogeoscapheus* sp. (Blaberidae), palm, orchid plants, on (*Rattus assimilis*), in artificial insect

diet, in cat faeces, metallactus colony, Western Australia: on pseudomys nanus (Halliday *et al.*, 1998); England and Wales: associated with yew (Skorupski and Luxton, 1998); Kuril Island: (*Microtus oeconomus*) (Rodentia, Cricetidae), walls of home aquarium above water level (Klimov, 1998); China (RuiYi *et al.*, 2000); New South Wales: Jenolan caves (Halliday, 2001); Latvia: fields, nests of wild birds (Salmane, 2001a), soil (Salmane 2001b), associated with *Aphylllophorales* fungi from dead or live trees and tree trunks (Salmane, 2005), in rotting (*Alnus incana*) wood, under bark of fallen (*Populus tremula*), in frass and under bark of (*Betula pendula*), in rotting wood from (*Quercus robur*) hole, jointly with (*Soronia grisea*) (Coleoptera, Nitidulidae) in yeasty sap of (*Betula pendula*), in frass and under bark of (*Betula pendula*) (Salmane, 2007); Sakhalin and Kuril Island: litter (Marchenko and Mapyehko, 2002); India (Chaudhury *et al.*, 2005); Slovakia: nests of mound-building mouse (*Mus spicilegus*) (Masan and Stanko, 2005), birds nests (Fenda and Schniererova, 2010); Northern Taiga (Makarova, 2009); Finland: decid forest, shore, compost, dung, alpine (Huhta *et al.*, 2010).

Iran Records. Iran (Sepasgozarian, 1971; Modares awal, 1994; Modares awal, 1997; Ostovan and Mosadegh; 1999); Fars, soil (Ostovan, 1993; Ostovan and Kamali, 1994; Amirazodi and Ostovan, 2012); Dezful, soil of citrus orchards (Malekzade *et al.*, 2000); Esfahan, soil of tree fruit orchards (Jalaeian *et al.*, 2004); Orumieh, potato fields (Mosavi *et al.*, 2004); Ahwaz, soil (Baharloo *et al.*, 2006); Bushehr, Chaharmahal-Bakhtiari and Khuzestan provinces, associated with insect (Nemati and babaeian, 2010); Gilan, soil (Hajizadeh *et al.*, 2010); Marvdasht, soil (Soleimani *et al.*, 2011); North-West of Kerman, soil (Masnavipour *et al.*, 2011); Shahreza, soil (Kavianpour *et al.*, 2011); Shiraz, soil (Montazeri *et al.*, 2011); Bam region, Ghale'e Asgar (Kerman province), soil and litter (Mehrzhad *et al.*, 2012); Darab region, soil and foliage of cotton fields (Khorsand *et al.*, 2012); Fars province, Doroodzan, soil

(Asadpoor *et al.*, 2012); Larestan region, cotton and wheat field (Khadempour *et al.*, 2012); Rafsanjan, Fruit trees, (Mohammadi *et al.*, 2012); Shiraz (Fars Province), associated with insectariums (samples taken from potato, zucchini, flour and the floor of insectariums) (Granpayeh *et al.*, 2012); Zanjan, soil (Zare *et al.*, 2012).

Records in this survey. Soil of different parts of Esfahan: 12.iii.2012, Bagh-daryache street; 29.iii.2012, Bostan Saadi street; 21.iii.2012, Kohandej street; 28.iii.2012, Marnan park; 05.viii.2011, 25.iii.2012, Ordibehesht street.

Family Parholaspididae

Genus *Proparholaspulus* Ishikawa, 1980

Proparholaspulus angustatus Ishikawa, 1987

Diagnosis. Dorsal shield entire, weakly reticulated, bearing 32 pairs of simple setae, *J1* shorter than *z1*, with two pairs of presternal shields, epyginal shield coalesced with ventri-anal shield, genitor-ventri-anal shield about twice longer than wide, epistome with elongate median extension and several short spines on either side, and without lateral elongate projections (Ishikawa, 1987).

World distribution. Philippine, litter (Ishikawa, 1987).

Record. Esfahan, soil, coll., A. Nemati, 2012.

Family Trachytidae

Genus *Uroseius* Berlese, 1888

Uroseius traegardhi Hirschmann and Zimgieble-Nicol, 1969

Diagnosis. Dorsal marginal setae (outside shield) inserted on a sclerotized band, metapodal shields short, oval, twice as long as wide, setae *c1*, *c2* short, subequal in length; *c1* slightly more dilated basally (Fain, 1998).

World distribution. Algeria, The Netherlands, France, Switzerland (Wisniewski and Hirshmann, 1993); North Africa, Europe (Fain, 1998).

Records. Esfahan, soil, 28.iii.2011, Marnan park.

Note: The uropodine mites of Iran with short information about their distribution and habitats have been reviewed by Kazemi and

Kontschan (2007). In their study they reported *Uroseius infirmus* (Berlese, 1887) without information about its distribution. Kazemi and Rajaei (2013) published an annotated checklist of Iranian Mesostigmata excluding the dubious species. Because they didn't mention this genus, so this is the first record of the genus *Uroseius* and *U. traegardhi* from Iran.

References

- Airolli, J. P., Solomon, L. and Duca, A. V. 1989. Les Gamasides (Acari) des nids de la forme fouisseuse du campagnol terrestre *Arvicola terrestris* L. Revue Suisse de Zoologie, 96 (1): 161-186.
- Al-Atawi, F. J. 2011. Six new records of predaceous mites associated with some trees from Riyadh, Saudi Arabia. Acarines, 5 (1): 37-39.
- Ambros, M. 1987. Mites (Acari: Mesostigmata) from small mammals in Hungary. Parasite Hungary, 20: 99-107.
- Amirzodi, R. and Ostovan, H. 2012. Reports of mites associated with conifers collected in Science and Research Branch. Proceedings of 20th Iranian Plant Protection Congress, Shiraz, Iran, p. 484.
- Asadpoor, N., Ostovan, H. and Haghani, M. 2012. A faunistic study on edaphic mesostigmatic mites in Doroodzan's region fields. Proceedings of 20th Iranian Plant Protection Congress, Shiraz, Iran, p. 515.
- Athias-Henriot, C. 1959. Phytoseiidae et Aceosejidae (Acarina, Gamasina) d'Algérie. III. Contribution aux Aceosejinae. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord, 50: 158-195.
- Baharlo, M., Shishehbor, P., Mossadegh, M.S., Khanjani, M. and Ueckermann, E. 2006. Investigation on the fauna of mesostigmatic mites of Ahvaz region. Proceedings of 17th Iranian Plant Protection Congress, Karaj, Iran, p. 191.
- Baker, E. W., Delfinado, M. D. and Abbatiello, M. J. 1976. Terrestrial mites of New York II. Mites in birds nest (Acarina). Journal of New York Entomological Society, 84 (1): 48-66.
- Beaulieu, F. and Weeks, A. R. 2007. Free-living mesostigmatic mites in Australia: their roles in biological control and bioindication. Australian Journal of Experimental Agriculture, 47: 460-478.
- Bedano, J. C. and Ruf, A. 2007. Soil predatory mite communities (Acari: Gamasina) in agroecosystems of Central Argentina. Applied Soil Ecology, 36 (1): 22-31.
- Bhattacharyya, A. K. and Sanyal, A. K. 2002. Three new species and some new records of the genus *Gamasellodes* Athias-Henriot (Acarina: Ascidae) from India. Acarologia, 42 (3): 229-238.
- Bloszyk, J., Gwiazdowicz, D. J., Bajerlein, D. and Halliday, R. B. 2005. Nests of the white stork *Ciconia ciconia* (L.) as a habitat for mesostigmatic mites (Acari, Mesostigmata). Acta parasitologica, 50 (2): 171-175.
- Bregetova, N. G. 1977. The family Aceosejidae Baker & Wharton, 1952. In: Ghilarov, M.S. (Ed.), A key to the Soil-Inhabiting Mites for Mesostigmata. Nauke, Leningard, USSR, pp. 169-226.
- Cakmak, I., Faraji, F. and Cobanoglu, S. 2011. A checklist and key to the Ascoidea and Phytoseioidea (except Phytoseiidae) species of Turkey with three new species records (Acari: Mesostigmata). Turkish journal of Entomology, 35 (4): 575-586.
- Chant, D. A. 1963. The subfamily Blattisocinae Garman (= Aceosejinae Evans) (Acarina: Blattisocidae Garman) (= Aceosejidae Baker and Wharton) in north America, with descriptions of new species. Canadian Journal of Zoology, 41: 243-305.
- Chaudhury, S., Roy, I., Podder, S., Gupta, S. K. and Saha, G. K. 2005. Diversity of synanthropic mites in Kolkata metropolis, India. Record of the Zoological Survey of India, 104 (3-4): 151-159.
- Christian, A. and Karg, W. 2006. The predatory mite genus *Lasioseius* Berlese, 1916 (Acarina, Gamasina). Abhandlungen und Berichte des Naturkunde Museums Görlitz, 77 (2): 99-250.

- Davydova, M. S. and Nikolskyi, V. V. 1986. The Gamasid Mites of Western Siberia. Novosibirsk. nauka publication, Russia.
- Edler, A. and Mehl, R. 1972. Mites (Acari, Gamasina) from small mammals in Norway. Norwegian Journal of Entomology, 19: 133-147.
- Ehara, S. 1964. Some mites of the families Phytoseiidae and Blattisocidae from Japan (Acarina, Mesostigmata). Journal of Faculty of Science, Hokkaido University, 15 (3): 378-394.
- Fadamiro, H. Y., Xiao, Y., Nesbitt, M. and Childers, C. C. 2009. Diversity and seasonal abundance of predacious mites in Alabamasatsuma citrus. Annals of the Entomological Society of America, 102 (4): 617-628.
- Fain, A. 1998. Description of mites (Acari) phoretic on Phoridae (Insecta: Diptera) with description of four new species of the genus *Uroseius* Berlese (Parasitiformes, Uropodina, Polyaspididae). International Journal of Acarology, 24 (3): 213-220.
- Fenda, P. and Schniererova, E. 2010. Mites (Acari: Mesostigmata) of the birds nests on the water in Southwestern Slovakia. Folia Faunistica Slovaca, 15 (8): 55-60.
- Gabrys, G., Makol, J., Bloszyk, J. and Gwiazdowicz, D. J. 2008. Mites (Acari) of the Karkonosze Mountains: a review. Biological letters, 45: 43-57.
- Gwiazdowicz, D. J. 2007. Ascid mites (Acari, Mesostigmata) from Selected Forest Ecosystems and Microhabitats in Poland. Wydawnictwo akademii rolniczej, Poznan.
- Gwiazdowicz, D. J. 2008a. Mesostigmatid Mites (Acari) Associated in Nests of Formicidae in Poland. In: Gwiazdowicz, D. J. (Eds.), Selected Problems of Acarological Research in Forests. Polskie Towarzystwo Lesne, Oddzial Wielkopolski Wydawnictwo Uniwersytetu Przyrodniczych w Poznaniu Poznan, Poland, pp. 97-113.
- Gwiazdowicz, D. J. 2008b. Mesostigmatid Mites (Acari) Associated with Scolytidae in Poland. In: Gwiazdowicz, D. J. (Eds.), Selected Problems of Acarological Research in Forests. Polskie Towarzystwo Lesne, Oddzial Wielkopolski Wydawnictwo Uniwersytetu Przyrodniczych w Poznaniu Poznan, Poland, pp. 59-97.
- Gwiazdowicz, D. J. 2010. Mites (Acari, Mesostigmata) of the Tatra National Park. Silvarum Colendararum Ratio et Industria Lignaria (Acta Scientiarum Polonorum), 9 (1): 5-18.
- Gwiazdowicz, D. J. and Klemt, J. 2004. Mites (Acari, Mesostigmata) in selected microhabitats of the Biebrza National Park (NE Poland). Biological letters, 41 (1): 11-19.
- Gwiazdowicz, D. J. and Kmita, M. 2004. Mites (Acari, Mesostigmata) from selected microhabitats of the Ujscie Warty National Park. Silvarum Colendararum Ratio et Industria Lignaria (Acta Scientiarum Polonorum), 3 (2): 49-55.
- Gwiazdowicz, D. J. and Matysiak, K. 2004. Mites (Acari, Mesostigmata) from selected microhabitats of the Bory Tucholskie National Park. Silvarum Colendararum Ratio et Industria Lignaria (Acta Scientiarum Polonorum), 3 (1): 17-24.
- Gwiazdowicz, D. J., Mizera, T. and Skorupski, M. 2000. Mites (Acari, Gamasida) from the nests of birds of prey in Poland. Buteo, 11: 97-100.
- Haitlinger, R. 1983. The mites (Acarina) of small mammals of the Pieniny Mts., Poland. Acta Zoologica Cracove, 26 (11): 355-386.
- Haitlinger, R. 1989a. Arthropods (Acari, Anoplura, Siphonaptera, Coleoptera) of small mammals of the Babia Gora Mts. Acta Zoologica Cracoviensia, 32 (2): 15-56.
- Haitlinger, R. 1989b. Arthropod communities occurring on small mammals from non-wooded areas of urban agglomeration of Wroclaw. Acta Parasitologica Polonica, 34 (1): 45-66.
- Hajizadeh, J., Faraji, F. and Rafati Fard, M. 2010. A new genus and four species records of the family Ascidae (Acari: Mesostigmata) for the Iran. Proceedings of the 19th Iranian Plant Protection Congress, Tehran, Iran, p. 356.

- Halliday, R. B. 2000. The Australian species of *Macrocheles* (Acarina: Macrochelidae). *Invertebrate Taxonomy*, 14 (2): 273-326.
- Halliday, R. B. 2001. Mesostigmatid mite fauna of Jenolan Caves, New South Wales (Acari: Mesostigmata). *Australian Journal of Entomology*, 40 (4): 299-311.
- Halliday, R. B. 2005. Predatory mites from crops and pastures in South Africa: potential natural enemies of redlegged earth mite *Halotydeus destructor* (Acari: Penthaleidae). *Zootaxa*, 1079: 11-64.
- Halliday, R. B., Walter D. E., Lindquist E. E. 1998. Revision of the Australian Ascidae (Acari: Mesostigmata). *Invertebrate Taxonomy*, 12: 1-54.
- Huhta, V., Siira-Pietikäinen, A., Penttinen, R. and Rätty, M. 2010. Soil fauna of Finland: Acarina, Collembola and Enchytraeidae. *Memoranda Societatis pro Fauna et Flora Fennica*, 86: 59-82.
- Hurlbutt, H. W. 1971. Ascinae and Podocinae (Acarina: Mesostigmata) from Tanzania. *Acarologia*, 13 (2): 280-300.
- Ishikawa, K. 1986. Gamasid mites (Acarina) associated with Japanese Millipedes. Report of Research Matsuyama Shinonome Junior College, 17: 165-177.
- Ishikawa, K. 1987. Occurrence of *Proparholaspulus* (Acari: Parholaspidae) in the Philippines. *Bulletin of National Science Museum*, 13 (2): 79-84.
- Jalaeian, M., Saboori, A. and Seydoleslami, H. 2004. Introduction of some genera and species of mesostigmatic mites to the fauna of Iran. *Proceeding of 16th Iranian Plant Protection Congress*, Tabriz, Iran, p. 254.
- Kalúz, S. and Fenda, P. 2005. Mites (Acari, Mesostigmata) of the Family Ascidae of Slovakia. *NOI Press*, Bratislava.
- Karg, W. 1971. Acari (Acarina), Milben. Unterordnung Anactinochaeta (Parasitiformes). *Die freilebenden Gamasina (Gamasides), Raubmilben. Die Tierwelt Deutschlands*, Veb Gustav Fisher Verlag Jena.
- Karg W. 1993. Acari (Acarina), Milben Parasitiformes (Anactinochaeta), Cohors Gamasina Leach, Raubmilben. *Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise*, 59: 1-523.
- Karg, W., Freier B. 1995. 'Parasitiforme Raubmilben als Indikatoren für den ökologischen Zustand von Ökosystemen. (Mitteilungen aus der Biologischen Bundesanstalt für Land und Forstwirtschaft, Berlin-Dahlem.
- Kavianpour, M., Nemati, A., Kocheili, F. and Gwiazdowicz, D. J. 2011. Some mites of Mesostigmata (Acari) from Shahreza region, Iran. *proceeding of the First Persian Congress of Acarology*, Kerman, Iran, p. 26.
- Kazemi, Sh. and Kotschan, J. 2007. A review of Uropodina mite fauna (Acari: Mesostigmata) of Iran and description of two new species. *Zoology in the Middle East*, 42: 79-82.
- Kazemi, Sh. and Rajaei, A. 2013. An annotated checklist of Iranian Mesostigmata (Acari), excluding the family Phytoseiidae. *Persian Journal of Acarology*, 2 (1): 63-158.
- Khadempour, F., Ostovan, H., Haghani, M. and Farzaneh, M. 2012. The faunistic survey of Mesostigmatic mites (Acari: Mesostigmata) in cotton and wheat field in Larestan region. *Proceedings of 20th Iranian Plant Protection Congress*, Shiraz, Iran, p. 427.
- Khomjakov, I. P. 1988. Peculiarities of the succession dynamics in Gamazina mites inhabiting fallow lands of different age. *Fauna i ekologiya paukoobraznykh, Russia*.
- Khorsand, A., Ostovan, H., Amin, Gh. A. and Khorsand, Z. 2012. Study on fauna of mesostigmata mites (Acari: Mesostigmata) associated with cotton fields in Darab region. *Proceedings of 20th Iranian Plant Protection Congress*, Shiraz, Iran, p. 513.
- Klimov, P. B. 1998. To the knowledge of mites and ticks (Acari) of Kuril Islands. *Far Eastern Entomologist*, 63: 1-36.
- Krantz, G. W. and Walter, D. E. 2009. *A Manual of Acarology*. 3th edition. Texas Tech University Press, USA.
- Lindquist, E. E., Krantz, G. W. and Walter, D. E. 2009. Order Mesostigmata. In: Krantz, G. W. and Walter, D. E. (Eds.), *A Manual of*

- Acarology. Texas Tech University Press, USA, pp. 124-232.
- Makarova, O. L. 2000. To studying mites of the genus *Arctoseius* Thor (Parasitiformes: Ascidae) on the far north. 1. Morphometric analysis of the taxonomical features of the arctic species and the description of *Arctoseius tschernovi* new species and *A. miranalisis* new species. *Zoologicheskii Zhurnal*, 79 (7): 800-817.
- Makarova, O. L. 2009. The fauna of free-living Gamasid mites (Parasitiformes, Mesostigmata) in the Northern Taiga: an analysis of the zonal specificity. *Entomological Review*, 89 (9): 1177-1193.
- Malekzadeh, M., Arbabi, M. and Shakerzadeh, A. 2000. Studies on fauna of citrus mites in Dezful area. *Proceeding of 14th Iranian Plant Protection Congress*, Esfahan, Iran, p. 265.
- Marchenko, I. and Mapyehko, N. 2002. Faunistic review of free-living Gamasina mites (Acari, Mesostigmata) from Sakhalin and Kuril Islands. *Euroasian Entomological Journal*, 1 (2): 31-48.
- Masnavipour, M., Kazemi, Sh., Latifi, M. and Ziaaddini, M. 2011. Edaphic mesostigmatic mites (Acari: Mesostigmata) fauna of north-western Kerman. *Proceeding of the First Persian Congress of Acarology*, Kerman, Iran, p. 41.
- Masan, P. and Stanko, M. 2005. Mesostigmatic mites (Acari) and fleas (Siphonaptera) associated with nests of mound-building mouse, *Mus spicilegus* Petényi, 1882 (Mammalia, Rodentia). *Acta Parasitologica*, 50 (3): 228-234.
- Mehl, R. 1979. Checklist of Norwegian ticks and mites (Acari). *Norwegian Journal of Entomology*, 26 (1): 31-45
- Mehrzad, N., Kazemi, Sh. and Masnavipour, M. 2012. Soil-inhabiting Mesostigmata (Acari) in Bam region, Kerman province. *Proceedings of 20th Iranian Plant Protection Congress*, Shiraz, Iran, p.471.
- Mineiro, J. L. DE.C., Lindquist, E. E. and De Moraes, G. J. 2009. Edaphic ascid mites (Acari: Mesostigmata: Ascidae) from the state of Sao Paulo, Brazil, with description of five new species. *Zootaxa*, 2024: 1-32.
- Modarres-Awal, M. 1994. *List of Agricultural Pests and Their Natural Enemies in Iran*. Ferdowsi University Publication, Mashhad.
- Modarres-Awal, M. 1997. *List of Agricultural Pests and Their Natural Enemies in Iran* (revised edition). Ferdowsi University Publication, Mashhad.
- Mohammadi, E., Izadi, H., Khanjani, M. and Samia, M. A. 2012. Faunistic study of Mesostigmatic mites (Acari: Mesostigmata) on fruit trees in Rafsanjan region of Iran. *Proceedings of 20th Iranian Plant Protection Congress*, Shiraz, Iran, p. 508.
- Montazeri, N., Nemati, A., Ostovan, H. and Gwiazdowicz, D. J. 2011. Fauna of some mesostigmatic mites (Acari: Mesostigmata) in Shiraz, Iran. *Proceeding of the First Persian Congress of Acarology*, Kerman, Iran, p. 44.
- Mosavi, Sh., Ostovan, H. and Adldoost, H. 2004. Study on the mite fauna of potato fields in Orimieh. *Proceeding of 16th Iranian Plant Protection Congress*, Tabriz, Iran, p. 263.
- Nemati, A. and Babaeian, E. 2010. Mites associated with insects in Chaharmahal-Bakhtiari, Khuzestan and Bushehr provinces. *Proceedings of 19th Iranian Plant Protection Congress*, Tehran, Iran, p. 364.
- Nemati, A., Gwiazdowicz, D. J., Riahi, E. and Mohseni, M. 2012a. Catalogue of the Iranian Mesostigmatid mites part 1: family Ascidae. *International Journal of Agriculture and Crop Sciences*, 4 (14): 1005-1011.
- Nemati, A., Riahi, E., Mohseni, M. and Gwiazdowicz, D. J. 2012b. Catalogue of the Iranian Mesostigmatid mites part 2: family Blattisociidae. *International Journal of Agriculture and Crop Sciences*, 4 (19): 1415-1420.
- Ostovan, H. 1993. Faunistic study of stores product mites in Kazerun and biology of important species, MSc thesis, Tarbiat Modarres University, Tehran. 172 pp.
- Ostovan, H. and Kamali, K. 1994. Mites of family Ascidae (Acari: Mesostigmata) associated with stored products in Kazerun,

- south of Iran. *Journal of Entomological Society*, 14: 9-18
- Ostovan, H. and Mosaddegh, M. S. 1999. Mites associated with honey bee (*Apis mellifera* L.) hives in Iran. *Proceeding of 35th International Congress, Apiculture (Apimondia)*, Canada, p. 254.
- Petrova, A. D. 1982. On soil Gamasid Mites Fauna (Parasitiformes, Mesostigmata) of Moscow Oblast. *Nauka publ, Russia*.
- Plumari, P. 2009. New records of Ascidae and Blattisociidae species from Italy (Acarina, Mesostigmata). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 150 (1): 77-90.
- RuiYu, Y., WieDong, L., Bing, S. 2000. Six new records of gamasid mite in Xinjiang and the deutonymph morphology of *Hirstionyssus meridianus* Zemskaia, 1955 (Acari). *Endemic Diseases Bulletin*, 15 (4): 48-50.
- Salmane, I. 1999. Soil free-living predatory Gamasina mites (Acari, Mesostigmata) from the coastal meadows of Riga Gulf, Latvia. *Latvijas Entomologs*, 37: 104-114.
- Salmane, I. 2001a. A check-list of Latvian Gamasina mites (Acari, Mesostigmata) with short notes to their ecology. *Latvijas Entomologs*, 38: 50-61.
- Salmane, I. 2001b. Investigations of Gamasina mites in natural and man-affected soils in Latvia (Acari: Mesostigmata). *Proceedings 13th international colloquium European Invertebrate Survey, Leiden*, pp: 129-137.
- Salmane, I. 2005. List of Mesostigmata mites (Acari, Parasitiformes) Associated with Aphyllophorales Fungi (Basidiomycetes) in Latvia. *Latvijas Entomologs*, 42: 57-71.
- Salmane, I. 2007. Mesostigmata Mite (Acari, Parasitiformes) Fauna of Wood-Related Microhabitats in Latvia. *Latvijas Entomologs*, 44: 69-86.
- Salmane, I. and Heldt, S. 2001. Predatory soil mites (Acari, Mesostigmata, Gamasina) from the Western Baltic Coast of Latvia. *Acarologia*, 41, 295-301.
- Salmane, I. and Kontschan, J. 2005a. Soil Gamasina mites (Acari, Parasitiformes, Mesostigmata) from Hungary. I. *Latvijas Entomologs*, 42: 48-56.
- Salmane, I. and Kontschan, J. 2005b. Soil Mesostigmata mites (Acari, Parasitiformes) from Hungary. II. *Latvijas Entomologs*, 43: 14-17.
- Seniczak, S., Klimek, A. and Kaczmarek, S. 2000. Soil Mites (Acari) of the Ecotone Between the Scots Pine Forest and Meadow in the Forest Landscape in Tuchola Forest, Poland. In: Rychling, A., Lechnio, J. and Malinowska, E. (Eds.), *Landscape Ecology (IALE). The Problems of Landscape Ecology*. In Pultusk school of Humanities IALE, PALE, Warszawa, pp. 247-260.
- Sepasgozarian, H. 1971. Mites of Economic Importance in Iran, Tehran University Publication, Faculty of Agriculture, Karaj.
- Shoukry, I. F., Kandeel, M. H. and El-Bashier, Z. 1990. Incidence of acarofauna in the house dust in Shaiqiya Governorate. *Egypt. Bulletin de la societe Entomologique de Egypte*, 69: 183-195.
- Skorupski, M., Belter, W., Kamczye, J. and Wierzbicka, A. 2008. Soil mites (Acari, Mesostigmata) of the Torfowiska Doliny Izery reserve in the Sudety Mountains. *Soil Organism*, 80 (2): 261-270.
- Skorupski, M. and Luxton, M. 1998. Mesostigmatid mites (Acari: Parasitiformes) associated with yew (*Taxus baccata*) in England and Wales. *Journal of Natural History*, 32 (3): 419-439.
- Slomain, S., Gulvik, E., Madej, G. and Austad, I. 2005. Gamasina and Microgyniina (Acari, Gamasida) from soil and tree hollows at two traditional farms in Sogn og Fjordane, Norway. *Norwegian journal of Entomology*, 52 (1): 39-48.
- Soleimani, M., Ostovan, H. and Joharchi, O. 2011. Mesostigmatic mites (Acari: Mesostigmata) in Marvdasht, Fars province, Iran. *Proceedings of the First Persian Congress of Acarology, Kerman, Iran*, p. 17.
- Spain, A. V. and Luxton, M. 1971. Catalog and bibliography of the Acari of the New Zealand subregion. *Pacific Insect Monograph*, 25: 179-226.

- Takaku, G. and Sasaki, T. 2007. Arboreal and forest floor mites (Acari: Gamasida, Oribatida) found in the Tomakomai experimental forest of Hokkaido university, Hokkaido, Northern Japan. *Journal of Hokkaido university of education*, 58 (1): 23-36.
- Tryjanowski, P., Baraniak, E., Bajaczyk, R., Gwiazdowicz, D. J., Konwerski, S., Olszanowski, Z. and Szymkowiak, P. 2001. Arthropods in nests of the red-backed shrike (*Lanius collurio*) in Poland. *Belgian Journal of Zoology*, 131 (1): 69-74.
- Walter, D. E. 1988. Nematophagy by soil arthropods from the shortgrass steppe Chihuahuan desert and Rocky Mountains of the Central United State. *Agriculture, Ecosystems and Environment*, 24: 307-316.
- Walter, D. E. Halliday, R. B and Lindquist E. E. 1993. A review of the genus *Asca* (Acarina: Ascidae) in Australia, with descriptions of three new leaf-inhabiting species. *Invertebrate Taxonomy*, 7: 1327-1347.
- Walter, D. E. and Lindquist, E. E. 1989. Life history and behavior of mites in the genus *Lasioseius* (Acari: Mesostigmata: Ascidae) from grassland soils in Colorado, with taxonomic notes and description of a new species. *Canadian Journal of Zoology*, 67 (11), 2797-2813.
- Walter, D. E., Seeman, O., Rodgers, D. and Kitching, R. L. 1998. Mites in the mist: how unique is a rainforest canopy-knockdown fauna. *Australian Journal of Ecology*, 23: 501-508.
- Wisniewski, J. and Hirschmann, W. 1993. *Gangsystematik der Parasitiformes, Teil 584, Katalog der Gunggtungen, Unterguttungen, Gruppen und Arten der Uropodiden der Erde (Taxonomie, Literatur, Grosse, Verbreitung, Vorkommen)*. In: Hirschmann, W. and Wisniewski, J. (Eds.), *Acari Parasitiformes Supercohors Atrichopygidiina Hirschmann 1975 Die Uropodiden der Erde*. Hirschmann Verlag, Nurnberg, pp. 1-220.
- Wissuwa, J., Salamon, J. A. and Frank, T. 2012. Effects of habitat age and plant species on predatory mites (Acari, Mesostigmata) in grassy arable fallows in Eastern Austria. *Soil Biology and Biochemistry*, 50: 96-107.
- Zare, M., Rahmani, H., Faraji, F. and Akrami, M. A. 2012. *Ascoidea (Acari: Mesostigmata) of Zanjan County of Iran with record of two new species*. *Proceedings of the 20th Iranian Plant Protection Congress, Shiraz, Iran*, p.459.

برخی از کنه‌های راسته میان‌استیگمایان از ایران و پراکنش جهانی آنها

فریبا کدخدایی الیادرائی^۱، علیرضا نعمتی^{۲*} و فرحان کچیلی^۱

۱- گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه شهید چمران اهواز، ایران

۲- گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه شهرکرد، ایران

* پست الکترونیکی نویسنده مسئول مکاتبه: nemati.alireza@agr.sku.ac.ir

دریافت: ۲ بهمن ۱۳۹۱؛ پذیرش ۹ اسفند ۱۳۹۱

چکیده: کنه‌های میان‌استیگمایان خاکزی از بندپایان و سایر بی‌مهرگانی تغذیه می‌کنند که ممکن است باعث تأثیر روی تعادل طبیعی موجودات زنده شده و از این جهت دارای اهمیت زیادی می‌باشند. به‌منظور مطالعه فون برخی از خانواده‌های راسته میان‌استیگمایان، نمونه‌هایی از خاک و گیاه‌خاک از مناطق مختلف شهرستان اصفهان جمع‌آوری شد. کنه‌ها با استفاده از قیف برلز استخراج و پس از شفاف شدن با اسید لاکتیک، از نمونه‌ها به کمک محلول هویر اسلاید دائمی تهیه شد. در این تحقیق پراکنش گونه‌ها در جهان و ایران همراه با اطلاعاتی در مورد زیستگاه آن‌ها ارائه شده است. ۲ جنس و ۳ گونه برای فون کنه‌های ایران جدید بوده و با علامت ستاره مشخص شده‌اند. جنس‌ها و گونه‌های شناسایی شده به شرح زیر می‌باشند:

Ascidae: *Arctoseius cetratus* (Sellnick, 1940), *A. pristinus** Karg, 1962, *A. venustus* (Berlese, 1917), *Gamasellodes bicolor* (Berlese, 1918); **Blattisociidae:** *Lasioseius sugawarai* Ehara, 1964, *L. youcefi* Athias-Henriot, 1959; **Melicharidae:** *Proctolaelaps pygmaeus* (Muller, 1860); **Parholaspididae:** *Proparholaspulus angustatus** Ishikawa, 1987; **Trachytidae:** *Uroseius traegardhi** (Hirschmann & Zirngiebl-Nicol, 1969).

واژگان کلیدی: کنه‌ها، میان‌استیگمایان، خاک، پراکنش جهانی، ایران