

Short Paper

## New record of the genus and species *Callitula ferrierei* (Bouček) (Hymenoptera: Pteromalidae) from Iran

Zainab-Alssadat Bayegan<sup>1\*</sup>, Hosseinali Lotfalizadeh<sup>2</sup>, Mohammad-Reza Zargaran<sup>3</sup> and Rana Pooraiouby<sup>4</sup>

1. M. Sc. Student of Agricultural Entomology, Department of Plant Protection, Islamic Azad University, Tabriz Branch, Tabriz, Iran.
2. Department of Plant Protection, Azarbaijan-e-Sharghi Research Center for Agriculture and Natural Resources, Tabriz, Iran.
3. Department of Forestry, Natural Resource Faculty, University of Urmia, Urmia, Iran.
4. Department of Plant Protection, Islamic Azad University, Tabriz Branch, Tabriz, Iran.

**Abstract:** Among the recently collected chalcidoid wasps in a rice field of eastern Guilan province (coast of Caspian Sea), in July 2012, there was a specimen of Pteromalidae belonging to the genus *Callitula*. It was identified as *Callitula ferrierei* (Bouček, 1964). Both genus and species are new record for Iranian fauna. Morphological characters and geographical distribution of the newly recorded species are given.

**Keywords:** *Callitula*, new record, Iran, fauna, distribution

### Introduction

The family Pteromalidae (Hymenoptera: Chalcidoidea) with more than 3,500 described species and 390 genera is one of the largest families among parasitic Hymenoptera. It includes important natural enemies of many plant pests, widely distributed in major insect orders such as Coleoptera, Diptera, Lepidoptera, Hymenoptera and Hemiptera (Dzhanokmenk, 1989, 1990).

Based on recently published data about 100 pteromalids species have been reported in Iran so far (Lotfalizadeh and Gharali, 2008; Alipanah *et al.*, 2013; Mitroiu *et al.*, 2011; Hasani *et al.*, 2011; Hasani and Madjdzadeh, 2012).

### Materials and Methods

Among the recently collected chalcidoid wasps from Langroud (37°08'050"N and 50°16'750"E),

in eastern Guilan province (coast of Caspian sea), we found a female specimen of *Callitula* (Hym.: Pteromalidae) in a Malaise trap set up in a rice field on July, 2012. Identification was made using Bouček (1964) and Xiao *et al.*, (2005). The morphological terminology follows Bouček and Rasplus (1991).

### Results and Discussion

The genus *Callitula* has been reported here for the first time from Iran. This genus includes 38 species in the world, with a cosmopolitan distribution (Noyes, 2013). Bouček (1964), Bouček and Rasplus (1991) and Xiao *et al.* (2005) listed following characters for this genus: Metallic body dark green; head without occipital carina; antennal formula 1, 1, 3, 5, 3, calva bearing distinct spicula, antennae inserted in the middle of face; pronotum distinctly narrower than mesonotum and with frontal carina dorsal, notauli incomplete, popodeum with distinct reticulate and subglobose nucha; first metasomal tergite evidently enlarged.

Handling Editor: Dr. Ali Asghar Talebi

\*Corresponding author, e-mail: za.bayegan@gmail.com  
Received: 22 October 2013, Accepted: 25 November 2013  
Published online: 27 November 2013

This genus is nearly allied to three genera in the West-Palaearctic region: *Norbanus* Walker, *Homoporus* Thomson and *Merisus* Walker with sharing characteristic spicula on antennal calva (Fig. 1D), but the genus *Homoporus* can be separated by having 3-segmented anelli.


***Callitula ferrierei* (Bouček, 1964) (Fig. 1)**

**Material examined:** Iran, Guilan, Langroud, 37°08'050"N & 50°16'750"E, Malaise trap, July 2012, Z. Bayegan leg., 1♀ (deposited in Insect collection of Department of Plant Protection, Azarbaijan-e-Sharghi Research Center for Agriculture and Natural Resources, Tabriz).

All the features of the specimen we collected in Iran match well with those reported by Bouček (1964). The typical morphological features of *C. ferrierei* are as follow: Body length in female about 2 mm,

body metallic dark green with brownish-yellow scape and legs, metasoma brown with dorsal dark bands (Fig. 1B); scape not reaching frons level, first funicular segment quadrate (Fig. 1D); mesosoma slightly arched in profile (Fig. 1A); marginal vein longer than stigmal and postmarginal veins (Fig. 1C); metasoma acuminate (Fig. 1E), not longer than the thorax (Fig. 1A), sunken dorsally (in dried specimens), first tergite about 0.25 length of metasoma, second tergite shorter than first tergite but not much longer than the third (Fig. 1E).

*Callitula ferrierei* had previously been reported from China (Xiao *et al.*, 2005), Bulgaria, Czech Republic, Germany, Moldova, Netherlands, Romania, Slovakia, Sweden, United Kingdom, England and former Yugoslavia (Noyes, 2013).



**Figure 1** *Callitula ferrierei*: (A): female in lateral view, (B): female in dorsal view, (C): forewing, (D): female antenna with calva bearing a distinct spicula, (E): gaster in dorsal view (T1-3 = first to third tergite).

*Callitula* species generally attack insects living in leaves or stems of their host plants; although hosts range from some Diptera (Chloropidae, Agromyzidae, Cecidomyiidae) (Bouček, 1964), to Hymenoptera (Tenthredinidae). *Callitula ferrierei* is mainly primary parasitoid, rarely as hyperparasites of the above dipteran hosts through Procrupoidea (Xiao et al., 2005) and in the studied area it may be parasitoid of dipterous pests of rice fields.

### Acknowledgments

We are grateful to Islamic Azad University, Tabriz Branch, and Azarbaijan-e-Sharghi Research Center for Agriculture and Natural Resources for supporting this research.

### References

- Alipanah, V., Nazemi Rafie, J. and Lotfalizadeh, H. 2013. First report of three pteromalid wasps from Iran. *Journal of Plant Protection*, 26 (4): 60.
- Bouček, Z. 1964. Die Europäischen arten der Gattung *Callitula* Spinola (Hymenoptera: Pteromalidae). *Entomophaga*, 9 (1): 9-15.
- Bouček, Z. and Rasplus, J. Y. 1991. Illustrated key to West-Palearctic genera of Pteromalidae (Hymenoptera: Chalcidoidea). Institut National de la Recherche Agronomique, 140 pp.
- Dzhanokmenk, A. 1989. Trophic links of the pteromalid wasps with the Diptera. *Entomological Review*, 68 (3): 92-98.
- Dzhanokmenk, A. 1990. Trophic association of parasitic Hymenoptera of the family Pteromalidae. *Entomologicheskoe Obozrenie*, 119 (4): 764-781.
- Hasani, A. and Madjdzadeh, S. M. 2012. Contribution to the knowledge of the Pteromalidae (Hymenoptera: Chalcidoidea) from Khorasan Razavi province, northeastern Iran. *Iranian Journal of Animal Biosystematics*, 8 (1): 57-69.
- Hasani, A., Mitroiu, M. D. and Madjdzadeh, S. M. 2011. New records of Pteromalidae (Hymenoptera: Chalcidoidea) from northeastern Iran. *Acta Zoologica Bulgarica*, 63 (3): 323-325.
- Lotfalizadeh, H. and Gharali, B. 2008. Pteromalidae (Hymenoptera: Chalcidoidea) of Iran: New record and preliminary checklist. *Entomofauna*, 29 (6): 93-120.
- Mitroiu, M. D., Abolhassanzadeh, F. and Madjdzadeh, S. M. 2011. New records of Pteromalidae (Hymenoptera: Chalcidoidea) from Iran, with description of a new species. *North-Western Journal of Zoology*, 7 (2): 243-249.
- Noyes, J. S. 2013. Universal Chalcidoid Database. The Natural History Museum. Available on: <http://www.nhm.ac.uk/entomology/chalcidoid>. (accessed 19 October, 2013).
- Xiao, H., Polaszek, A. and Huang, D. 2005. A study of the genus *Callitula* Spinola (Hymenoptera: Pteromalidae) from China. *Oriental Insects*, 39: 233-240.

گزارش جدید جنس و گونه *Callitula ferrierei* (Bouček) (Hymenoptera: Pteromalidae) از ایرانزینب‌السادات بایگان<sup>۱\*</sup>، حسینعلی لطفعلی‌زاده<sup>۲</sup>، محمدرضا زرگران<sup>۳</sup> و رعنا پورایوبی<sup>۴</sup>

- ۱- گروه گیاهپزشکی، دانشگاه آزاد اسلامی، واحد تبریز، تبریز، ایران.
  - ۲- بخش تحقیقات گیاهپزشکی، مرکز تحقیقات کشاورزی و منابع طبیعی استان آذربایجان شرقی، تبریز، ایران.
  - ۳- گروه جنگلداری، دانشکده منابع طبیعی، دانشگاه ارومیه، ارومیه، ایران.
  - ۴- گروه گیاهپزشکی، دانشگاه آزاد اسلامی واحد تبریز، تبریز، ایران.
- \* پست الکترونیکی نویسنده مسئول مکاتبه: za.bayegan@gmail.com  
دریافت: ۳۰ مهر ۱۳۹۲؛ پذیرش: ۴ آذر ۱۳۹۲

**چکیده:** در میان زنبورهای جمع‌آوری شده از بالاخانواده‌ی Chalcidoidea در یک مزرعه برنج در شرق استان گیلان (شهرستان لنگرود) طی تابستان ۱۳۹۱ نمونه‌ای از خانواده‌ی Pteromalidae متعلق به جنس *Callitula* مشاهده شد. این نمونه تحت عنوان *Callitula ferrierei* (Bouček, 1964) شناسایی شد. جنس و گونه‌ی آن برای فون ایران جدید هستند. ویژگی‌های مرفولوژیک افتراقی جنس و گونه و پراکنش جغرافیایی این گونه ارائه گردید.

**واژگان کلیدی:** *Callitula*. گزارش جدید، ایران، فون، پراکنش