

Research Article

Two species of the genus *Elachertus* Spinola (Hym.: Eulophidae) new larval ectoparasitoids of *Tuta absoluta* (Meyreck) (Lep.: Gelechiidae)

Fatemeh Yarahmadi^{1*}, Zohreh Salehi¹ and Hossein Lotfalizadeh²

1. Ramin Agriculture and Natural Resources University, Mollasani, Ahvaz, Iran.
2. East-Azerbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran.

Abstract: This is the first report of two ectoparasitoid wasps, *Elachertus inunctus* (Nees, 1834) in Iran and *Elachertus pulcher* (Erdös, 1961) (Hym.: Eulophidae) in the world, that parasitize larvae of the tomato leaf miner, *Tuta absoluta* (Meyrick, 1917) (Lep.: Gelechiidae). The specimens were collected from tomato fields and greenhouses in Ahvaz, Khuzestan province (south west of Iran). Both species are new records for fauna of Iran. The knowledge about these parasitoids is still scanty. The potential of these parasitoids for biological control of *T. absoluta* in tomato fields and greenhouses should be investigated.

Keywords: tomato leaf miner, parasitoids, identification, biological control

Introduction

The Eulophidae is one of the largest families of Chalcidoidea. The chalcid parasitoid wasps attack insects from many orders and also mites. Many eulophid wasps parasitize several pests on different crops. They can regulate their host's populations in natural conditions (Yefremova and Myartseva, 2004). Eulophidae are composed of four subfamilies, Entedoninae (Förster, 1856), Euderinae (Lacordaire, 1866), Eulophinae (Westwood, 1829) and Tetrastichinae (Förster, 1856) (Schauff *et al.*, 1997). The family has a total of 297 genera and 4472 described species (Noyes, 2015). The members of the subfamily Euolophinae are generally ectoparasitoid of many holometabolous insects (Noyes, 2015). Many species of the family have proven to be highly successful biological control agents. Although eulophid wasps are generally parasitoids of

holometabolous insects, the overall range of hosts and biologies in eulophid wasps is remarkably diverse (Gauthier *et al.*, 2000).

Species of the genus *Elachertus* Spinola, 1811 (Hym.: Eulophidae) are primary parasitoids of a variety of lepidopteran larvae. Some species are polyphagous that parasitize hosts belonging to different insect families. The larvae of these wasps are often gregarious and their pupae can be observed on the surface of plant leaves or the body of their host. They seem to prefer host larvae that occupy sheltered situations such as leaf rolls, leaf mines or that bore inside twigs or growing tips. Several of these host species are economically important (Schauff, 1985).

The tomato leaf miner, *Tuta absoluta* Meyrick, is considered as a key pest of tomato both in the field and under protected conditions (Yankova, 2012) in Asia (Desneux *et al.*, 2011). Larvae preferentially feed on all above-ground parts of tomato, create mines on the leaves and penetrate into young stems and fruits. Both yield and fruit quality can be significantly reduced by the direct feeding of the pest and the secondary pathogens which may then enter through the

Handling Editor: Ali Asghar Talebi

* **Corresponding author**, e-mail: yarahmadi@ramin.ac.ir
 Received: 20 February 2016, Accepted: 14 June 2016
 Published online: 26 June 2016

wounds made by the pest. Severely attacked tomato fruits lose their commercial value (Cristina *et al.*, 2008).

The following species belong to *Elachertus* genus was previously recorded from Iran: *E. gallicus* (Erdős, 1958) and *E. lateralis* (Spinola, 1808) (Yefremova *et al.*, 2007; Talebi *et al.*, 2011).

Materials and Methods

During bio-ecological studies on tomato leaf miner, *Tuta absoluta* (Meyrick) (Lep.: Gelechiidae), in tomato greenhouses and fields in Ahwaz (Khouzestan province, south west Iran), some larval parasitoid specimens were directly collected from tomato leaf mines by fine needle and reared at 25 ± 1 °C, RH 60% and photoperiod 16:8 L:D in incubator. The adult wasps emerged from last instar larvae.

The emerged parasitoids were carefully collected using an aspirator and placed into 75% ethanol for further examination. External morphology was illustrated using an Olympus™ SZH, equipped with a Canon™ A720 digital camera. The specimens were identified according to the reliable keys and descriptions (Askew and Boucek, 1968; Trjapitzin, 1978; Zhu and Huang, 2001; Yefremova and Myartseva, 2004; Lotfalizadeh, 2013).

The specimens were deposited in collection of the Department of Plant Protection, East-Azerbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran.

Results

In this research two *Elachertus* (Eulophidae) species were reared for the first time as larval ectoparasitoids of *Tuta absoluta*. These species are as follow:

***Elachertus inunctus* (Nees, 1834)** (Fig. 1 A-B)

Material examined: Iran, Khouzestan province, Ahwaz, greenhouse and fields of Veis region (N 3130 E 490), from *Tuta absoluta* (Meyrick) larvae on tomato, 17 March to 13 July 2015, 2♀ and 2♂.

Diagnosis of female. First funicular segment 3 times longer than wide, the rest about 2.5 times longer than wide; scutellum smooth and shining, mid lobe of mesoscutum with only 3 pairs of strong setae; metasoma of female 2.0 times as long as wide, legs completely yellow; head and thorax dark-coloured, scape brownish.

Geographical distribution. It is widely distributed in Europe (Austria, Czech Republic, Germany, Hungary, Italy, Lithuania, Moldova, Norway, Poland, Slovakia, Sweden, Switzerland, United Kingdom and former Yugoslavia) and in Asia (Japan, South Korea and Turkmenistan) (Noyes, 2015); new record for Iran.

Hosts. We reared *E. inunctus* for the first time from larval stage of tomato leaf miner, *T. absoluta* (Meyrick) (Lep.: Gelechiidae) in Iran. This species has been reported from different lepidopterous pests of the families Elachistidae, Epermeniidae, Gracillariidae, Lyonetiidae, Momphidae, Nepticulidae, Oecophoridae and Tortricidae (Noyes, 2015). The parasitoid wasps have been previously reported on *T. absoluta* (Desneux *et al.*, 2010; Marja *et al.*, 2011).

***Elachertus pulcher* (Erdős, 1961)** (Fig. 1 C-E)

Material examined: Iran, Khouzestan province, Ahwaz, greenhouse and fields of Veis region (N 3130 E 490), from *Tuta absoluta* larvae on tomato, 17 March to 13 July 2015, 5♀.

Diagnosis of female. First funicular segment 1.6 times longer than pedicel and about 3 times longer than wide, the rest more than 2.5 times longer than wide; mid lobe of mesoscutum with additional setae scattered on disc in addition to some pairs of strong setae. Body mainly yellow, head and thorax with small dark spots, antennae dark brown, legs yellow.

Geographical distribution. It has been reported from Europe (Austria, Croatia, Czech Republic, France, Germany, Hungary, Moldova, Russia, Slovakia and Spain) and Asia (Azerbaijan, Taiwan, Turkey, Turkmenistan and Yemen) (Noyes, 2015); new record for Iran.




Figure 1 Two *Elachertus* species parasitizing *Tuta absoluta* in the south of Iran: A, B- *Elachertus inunctus* (A- female in lateral view, B- male in lateral view); C-E- *Elachertus pulcher* (C- female in lateral view, D- female antenna, E- fore wing).

Hosts. The parasitoid wasp was collected on *T. absoluta* for the first time. There is no biological information about this species on *T. absoluta* larvae.

Discussion

Elachertus is a large and widespread genus of the Eulophinae, with over 111 species worldwide (Noyes, 1998). The genus includes more than 65 Palearctic species (Yefremova, 1998). This

species belongs to the *charondas* species group (Zhu and Huang, 2001). This is the first record of *E. inunctus* in Iran and *E. pulcher* on *Tuta absoluta* in the world. However, two unknown species of this genus have been reported by Zappala *et al.* (2012a) from Italy. These two unknown species cause 8.6% parasitism in larval population of *T. absoluta* (Zappala *et al.*, 2012 b).

Only two species of the genus *Elachertus* have been previously reported from Iran and we add two new records (Table 1) and this report extends

geographical distribution of *E. inunctus* and *E. pulcher*. Five species belong to genus *Elachertus* (including *E. charondas* Walker, *E. fenestratus* Nee, *E. isadas* Walker, *E. lateralis* Spinola and *E. pulcher*) were reported from Yemen (Yefremova, 2007) and five species including *E. charondas* Walker, *E. fenestratus* Nees, *E. inunctus*, *E. kopetdagensis* Yefremova & Myartseva and *E. pulcher* were reported from Turkmenistan.

Phyllocnistidae (*Phyllocnistis blancardella* F., *Ph. coryli* Nic.), Tortricidae, Leucopteridae, Oecophoridae, Elachistidae were previously mentioned as hosts of *E. inunctus* (Trjapitzin, 1978). Also, *E. pulcher* were collected from more than 70 arthropod species, 20% predators and 80 % parasitoids, were recorded attacking various life

stages of *T. absoluta* so far. Some parasitoid wasps belonging to Eulophidae and Braconidae have promising potential to be considered as effective tools in integrated pest management program of the pest in the newly invaded areas. Some identified and unidentified eulophid species belonging to genera *Baryscapus*, *Cirrospilus*, *Chrysocharis*, *Closterocerus*, *Diglyphus*, *Closterocerus*, *Elasmus*, *Hemiptarsenus*, *Necremnus*, *Necremnus*, *Neochrysochari* and *Elachertus* were reported as larval parasitoids of *T. absoluta* in the Palaearctic countries (Zappala *et al.*, 2013). Of the genus *Elachertus*, *E. inunctus* (during spring) and two unidentified species (during summer and autumn) were collected from the pest larvae in Italy (Zappala *et al.*, 2012a).

Table 1 Recorded species of the genus *Elachertus* from Iran.

<i>Elachertus</i> species	Host	Distribution	References
<i>E. fenestratus</i> Nees, 1834	<i>Cephus pygmaeus</i> L. (Hym.: Cephidae)	Widely distributed in Iran	Talebi <i>et al.</i> (2011)
<i>E. gallicus</i> Erdős, 1958	<i>Pyllocnistis citrella</i> Stainton (Lep.: Gracillariidae)	Fars province, Jahrom	Yefremova <i>et al.</i> (2007)
<i>E. inunctus</i> Nees, 1834	<i>Tuta absoluta</i> (Lep.: Gelechiidae)	Khouzestan province	this study
<i>E. pulcher</i> (Erdős, 1961)	<i>Tuta absoluta</i> (Lep.: Gelechiidae)	Khouzestan province	this study

Acknowledgements

The research was financially supported by research deputy of Ramin Agriculture and Natural Resources University of Khouzestan and East-Azerbaijan Research Center for Agriculture and Natural Resources, Tabriz. Many thanks to anonymous reviewers of this article for their suggestions.

References

- Askew, R. R. and Boucek, Z. 1968. Index of palaeactic Eulophidae (excl. Tetrastichinae). In: Delucchi, V. and Remaudiere, G. (Eds.), Index of Entomophagous Insects. Paris. 223 pp.
- Cristina, A. F., Jorge, B. T., Adriano, M. V. F. and Angela, M. I. F. 2008. Parasitism of *Tuta absoluta* in tomato plants by *Trichogramma pretiosum* Riley in response to host density and plant structures. *Ciencia Rural*, 38: 1504-1509.
- Desneux N., Luna M. G., Guillemaud T. and Urbaneja A. 2011. The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: The new threat to tomato world production. *Journal of Pest Science*, 84: 403-408.
- Desneux, N., Wajnberg, E., Wyckhuys, K. A. G., Burgio, G., Arpaia, S., Narváez-Vasquez, C. A., González-Cabrera, J., Ruescas, D. C., Tabone, E. and Frandon, J. 2010. Biological invasion of European tomato crops by *Tuta absoluta*: ecology, geographic expansion and prospects for biological control. *Journal of Pest Science*, 83 (3): 197-215.
- Gauthier, N., LaSalle, J., Quicke, D. L. J. and Godfray, H. C. J. 2000. Phylogeny of

- Eulophidae (Hymenoptera: Chalcidoidea), with a reclassification of Eulophinae and the recognition that Elasmidae are derived eulophids. *Systematic Entomology*, 25: 521-539.
- Lotfalizadeh, H. 2013. Introduction to Hymenopterous Parasitoids of Important Agricultural Pests. Islamic Azad University Press, Tabriz, 86 pp.
- Noyes, J. S. 1998. Catalogue of the Chalcidoidea of the world, Electronic publication (CD-ROM), ETI, Amsterdam.
- Noyes, J. S. 2015. Universal Chalcidoidea Database. World Wide Web electronic publication. Available on: <http://www.nhm.ac.uk/entomology/chalcidoids/index.html> (accessed September 05, 2015).
- Schauff, M. E. 1985. Taxonomic study of the Nearctic species of *Elachertus* Spinola (Hymenoptera: Eulophidae). *Proceeding of Entomological Society of Washington*, 87 (4): 843-858.
- Schauff, M. E., LaSalle, J. and Coote, L. D. 1997. Eulophidae. In: Gibson, G. A. P., Huber, J. T. and Woolley, J. B. (Eds.), *Annotated Keys to the Genera of Nearctic Chalcidoidea* (Hymenoptera). National Research Council of Canada, pp: 327-429.
- Talebi, A. A., Mohammadi Khoramabadi, A. and Rakhshani, E. 2011. Checklist of eulophid wasps (Insecta: Hymenoptera: Eulophidae) of Iran. *Check List*, 7 (6): 709-719.
- Trjapitzin, V. A. 1978. Hymenoptera II. Chalcidoidea 13. Eulophidae (excl. Tetrastichinae). *Opredelitel Nasekomykh Evropeyskoy Chasti SSSR*, 381-430.
- van der Straten, M. J., Potting, R. P. and van der Linden, A. 2011. Introduction of the tomato leafminer *Tuta absoluta* into Europe. *Proceedings of Netherlands Entomological Society Meeting*, 22: 23-30.
- Yankova, V. 2012. Damage caused by tomato leaf miner (*Tuta absoluta* Meyrick) in tomato varieties grown in greenhouse. *Plant Science*, 49: 92-97.
- Yefremova, Z. A. 1998. The subfamily Eulophinae (Hymenoptera: Eulophidae) of Palaearctic (Morphology, biology, systematic, evaluation and phylogeny). *Autoreferat Dissertatii Doktora Biologicheskikh Nauk*, St. Petersburg. 47 p.
- Yefremova, Z. A. 2007. The subfamilies Eulophinae, Euderinae and Entedoninae (Hymenoptera: Eulophidae) in Yemen. *Fauna of Arabia*, 23: 335-368
- Yefremova, Z. and Myartseva, S. N. 2004. Eulophidae (Hymenoptera: Chalcidoidea) of Turkmenistan, with emphasis on the genera *Elachertus* Spinola, 1811 and *Hyssopus* Girault, 1916. *Entomologist's Monthly Magazine*, 140: 113-122.
- Yefremova, Z., Ebrahimi, E. and Yegorenkova, E. 2007. The subfamilies Eulophinae, Entedoninae and Tetrastichinae in Iran, with description of new species (Hymenoptera, Eulophidae). *Entomofauna*, 30: 405-440.
- Zappala, L., Bernardo, U., Biodl, A., Alma, A., Al-Jboory, E., Arno, J., Bayram, A., Chailleux, A., El-Arnauty, A., Gerling, D., Guenaoui, Y., Shaltiel-Harpaz, L., Siscaro, G., Stavrinides, M., Tavella, L., Anzar, R.V., Urbaneja, A., Deneux, N. 2013. Natural enemies of the South American moth, *Tuta absoluta*, in Europe, North Africa and Middle East, and their potential use in pest control strategies. *Journal of Pest Science*, 86: 635-647.
- Zappala, L., Bernardo, U., Biodl, A., Cocco, A., Deliperi, S., Delipero, G., Giorgini, M., Pedata, P., Rapisarda, C., Tropea Garzia G. and Siscaro, G. 2012a. Recruitment of native parasitoids by the exotic pest *Tuta absoluta* in Southern Italy. *Bulletin of Insectology*, 65 (1): 51-61.
- Zappala, L., Biondi, A., Siscaro, G., Garzia, G. T., van Achterberg, K. and Desneux, N. 2012b. Adaptation of indigenous parasitoids to *Tuta absoluta* in Italy: the parasitic wasp *Bracon nigricans*. *Atti Accademia Nazionale Italiana di Entomologia*, Anno LX, pp. 85-93.
- Zhu, C. D. and Huang, D. W. 2001. A study of Chinese *Elachertus* Spinola (Hymenoptera: Eulophidae). *Zoological Studies*, 40 (4): 317-354.

دو گونه از جنس *Elachertus* Spinola (Hym.: Eulophidae) به عنوان پارازیتوئید خارجی
(Lep.: Gelechiidae) *Tuta absoluta* (Meyreck)

فاطمه یاراحمدی^{۱*}، زهره صالحی^۱ و حسین لطفعلی زاده^۲

۱- گروه گیاه پزشکی، دانشکده کشاورزی، دانشگاه کشاورزی و منابع طبیعی رامین خوزستان، ملاتانی، اهواز، ایران.

۲- مرکز تحقیقات کشاورزی و منابع طبیعی آذربایجان شرقی، تبریز، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: yarahmadi@ramin.ac.ir

دریافت: ۱ اسفند ۱۳۹۴؛ پذیرش: ۲۵ خرداد ۱۳۹۵

چکیده: این اولین گزارش دو گونه زنبور پارازیتوئید خارجی (*Elachertus inunctus* (Nees, 1834) در ایران و (*Elachertus pulcher* (Erdös, 1961) از خانواده Eulophidae در جهان از روی لارو مینوز برگ گوجه فرنگی، (*Tuta absoluta* (Meyrick, 1917) (Lep.: Gelechiidae) می باشد. این گونه ها از مزارع و گلخانه های گوجه فرنگی در استان خوزستان، اهواز (جنوب غربی ایران) جمع آوری شدند. هر دو گونه برای فون ایران جدید می باشند. اطلاعات کافی در مورد این دو پارازیتوئید وجود ندارد. کارایی بالقوه این پارازیتوئیدها برای کنترل بیولوژیک *T. absoluta* در مزارع و گلخانه های گوجه فرنگی می بایست مورد بررسی قرار گیرد.

واژگان کلیدی: مینوز برگ گوجه فرنگی، پارازیتوئیدها، شناسایی، کنترل بیولوژیک