

Research Article

Isolation and identification of entomopathogenic nematodes and their symbiotic bacteria from Kurdistan province in Iran

Arman Abdolmaleki¹, Zahra Tanha Maafi^{2*}, Hooshang Rafiee Dastjerdi¹, Bahram Naseri¹ and Abolghasem Ghasemi²

1. Department of Plant Protection, Faculty of Agricultural Sciences, University of Mohaghegh Ardabili, Ardabil, Iran.

2. Iranian Research Institute of Plant Protection, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran.

Abstract: Entomopathogenic nematodes (EPNs) are commonly used as biological agents for control of insect pests. This study was carried out to identify EPNs, determine dominant and frequent species in Kurdistan province, Iran and characterize their symbiotic bacteria. Identification of EPNs was performed based on morphological and morphometrical characters and also rRNA-ITS gene sequences. Two EPNs, *Heterorhabditis bacteriophora* (Rhabditida: Heterorhabditidae) and *Steinernema feltiae* (Rhabditida: Steinernematidae) were identified. Out of totally 150 soil samples collected mainly from mid-southern parts of the province, 60% were positive for EPNs. *Heterorhabditis bacteriophora* showed the highest frequency in this region, remarkably 59.3% of soil samples contained this species. Incidences of *H. bacteriophora* in grasslands, woodlands and alfalfa fields were 57.3, 14.6 and 28.0 percent, respectively. *Steinernema feltiae* was found only in alfalfa fields with 0.66% frequency. Bacterial symbionts of *H. bacteriophora* and *S. feltiae* were also identified based on biochemical characters and *recA* gene sequencing. In this research, two species of *Photorhabdus* were isolated from *H. bacteriophora* namely *P. luminescens* subsp. *kayaii* and *P. temperata* subsp. *thracensis*. *Xenorhabdus bovienii* was identified from infective juveniles of *S. feltiae*.

Keywords: *Heterorhabditis bacteriophora*, *Photorhabdus luminescens*, *Steinernema feltiae*, *Xenorhabdus bovienii*, Phylogeny

Introduction

Entomopathogenic nematodes (EPNs) belonging to the families Heterorhabditidae and Steinernematidae are obligate pathogens of insects. These nematodes have symbiotic relationship with gram negative bacteria *Photorhabdus* and *Xenorhabdus* (Enterobacteriaceae). The infective juveniles (IJs) penetrate into the haemocoel by invading through natural openings or enter the haemocoel directly through the insect's

intersegment integument (Wang and Gaugler, 1998). After penetration into the host haemocoel, IJs release their bacterial symbionts of the genera *Photorhabdus* and *Xenorhabdus* associated with *Heterorhabditis* and *Steinernema*, respectively (Martens *et al.*, 2003; Ciche and Ensign, 2003). The bacteria and nematodes cooperate to overcome the host's immune response and result in killing the insect (Boemare and Akhurst, 2006). The nematodes produce at least two generations in the host body, and emerge from the cadavers as IJs that search for new hosts (Poinar, 1990; Kaya and Gaugler, 1993).

Efficacy of EPNs for controlling insect pests caused a huge attempt to find new efficient isolates. Earlier, morphological and morphometric methods

Handling Editor: Ebrahim Pourjam

*Corresponding author, e-mail: tanhamaafi@iripp.ir

Received: 11 July 2015, Accepted: 20 February 2016

Published online: 22 April 2016

were used for EPNs identification, however, recently DNA sequences variation in the internal transcribed spacer (ITS) regions of ribosomal DNA is extensively used as a complementary approach to morphological descriptions and also to estimate phylogenetic relationships among species in EPNs (Stock, 2002).

Phenotypic characteristics, biochemical and molecular methods are three stages for identification of unknown bacterial colonies. There are series of important biochemical characteristics, however, bioluminescens and catalase test which are physiologically very significant characters, are positive for *Photorhabdus* and negative for *Xenorhabdus* (Forst *et al.*, 1997). Due to substantial differences in biochemical reactions of these bacteria (Grimont *et al.*, 1984; Akhurst and Boemare, 1988), results of these experiments could not approach to precise identification. Thus, molecular consideration of isolated bacteria was necessary. Recombinase A (*recA*) can serve as molecular marker for the identification of *Xenorhabdus* and *Photorhabdus* (Ludwig and Schleifer, 1999). RecA protein and *recA* gene sequence comparisons have been used to consider phylogenetic relationships among different genera and species of prokaryotes (Eisen, 1995).

Five species of entomopathogenic nematodes have been reported from Iran including *H. bacteriophora* Poinar, 1976, *S. bicornutum* Tallosi, Peters and Ehlers, 1995, *S. carpocapsae* Weiser, 1955, *S. feltiae* (Filipjev, 1934) Wouts, Mráček, Gerdin and Bedding, 1982, and *S. glaseri* (Steiner, 1929) Wouts, Mráček, Gerdin and Bedding, 1982. The species are mainly described from East and West Azerbaijan, Ardabil, Tehran, and North Khorasan provinces (Parvizi, 2001; Tanha Maafi *et al.*, 2006; Karimi and Kharazipakdel, 2007; Eivazian Kary *et al.*, 2009; Nikdel *et al.*, 2010; Kamali *et al.*, 2013). The bacterial strains associated with *S. glaseri* and *S. carpocapsae* in Iran were characterised as *X. nematophila* and *X. poinarii* (Karimi *et al.*, 2011). Taxonomic studies using biochemical and molecular analyses showed symbiotic relationship between *Photorhabdus* and *Heterorhbditis*, and *Xenorhabdus* and *Steinernema* (Euzeby and Boemare, 2000). Isolation and identification of

symbiotic bacteria of different EPNs were performed through several studies that identified different species including *P. luminescens* subsp. *lumondii*, *P. temperate*, *X. bovienii*, *X. budapestensis*, *X. nematophila*, *X. poinarii* (Iraki *et al.*, 2000; Hazir *et al.*, 2004; Bussaman *et al.*, 2009; Agazadeh, *et al.*, 2010; Jang *et al.*, 2011; Karimi *et al.*, 2011). The aims of this study were to: *a*: identify the entomopathogenic nematodes by morphological, morphometric and molecular data, *b*: find and introduce frequent native EPN species in Kurdistan province, *c*: characterize the symbiotic bacteria associated with EPNs based on biochemical and molecular approaches.

Materials and Methods

Soil Sampling and Isolation of EPNs

Soil samples were collected from oak forests, grasslands and alfalfa fields in mid-southern regions of Kurdistan province, Iran including: Qorveh, Dehgolan, Sanandaj, Sarv Abad and Marivan. Elevation of sampling locations was between 1106 and 2359 meters above sea level. The samples were taken during March and April 2012 at depth of 0-20 cm from an area of 100-500 m², each sample consisted of 5-10 subsamples of 50 g soil. Totally 150 soil samples were collected. Entomopathogenic nematodes were isolated from soil samples by using *Galleria* bait technique (Bedding and Akhurst, 1975). *Galleria mellonella* (wax moth larva) was reared on an artificial diet suggested by Bronskill (1961) at room conditions. Dead *Galleria* larvae were moved to a White trap for obtaining IJs from the cadavers (White, 1927), while a few cadavers were directly dissected to obtain adult nematodes of the first and second generations.

Morphological analysis

Infective juveniles and adult nematodes were fixed in TAF (triethanolamine 2 ml, formaldehyde 7 ml, distilled water 91 ml) and transferred to dehydrated glycerin according to De Grisse (1969). Morphological characters that are commonly used in EPNs diagnosis included: post anal swelling, vulval lips, presence of mucro and epiptygma and testis reflexion. Also,

morphometric features *i.e.* body length, anterior end to excretory pore, tail length, pharynx length, maximum body width, D% and E% ratios, spicule length, gubernaculum length and SW ratio were measured using a camera Lucida installed on an Olympus BH-2 microscope. The morphological and morphometric characters were analyzed according to Stock and Hunt (2005).

DNA extraction, PCR and sequencing

Freshly harvested infective juveniles were used for DNA extraction. Protocols of DNA extraction, PCR and sequencing were described by Tanha Maafi *et al.* (2003). The ITS regions of rRNA gene were amplified using universal primers, forward TW81 5'-GTTTCCGTAGGTGAACCTGC-3' and reverse AB28 5'-ATATGCTTAAGTTCAGCGG GT-3' (Joyce *et al.*, 1994). The PCR products were directly sequenced in both directions at Bioneer Company (South Korea). The obtained sequences along with their chromatograms were checked visually and aligned manually by using BioEdit program. The sequences were deposited in GenBank under accession numbers KR152294, KT070192 and KT070191. The sequences of ITS-rRNA of different isolates of *H. bacteriophora* and *S. feltiae* were used for phylogenetic analyses (Table 1). The newly obtained sequences and the sequences from GenBank database were aligned using Clustal X 1.83 (Thomson *et al.*, 1997) with default parameters. Sequence alignments were manually edited using BioEdit (Hall *et al.*, 1999). Outgroup taxa for each data set were chosen according to previous published data (Spiridonov *et al.*, 2004). Phylogenetic analysis of the sequence data sets was performed with Bayesian inference (BI) using MrBayes 3.2 (Huelsenbeck and Ronquist, 2001) under the HKY + G model, for each gene was initiated with a random starting tree and was run with four chains for 1.0×10^6 generations. The Markov chains were sampled at intervals of 100 generations. Posterior probabilities (PP) are given on appropriate clades.

Isolation of bacteria

Bacteria associated with EPNs were directly isolated from IJs. About 100 IJs were surface sterilized with sodium hypochlorite 1% for five

minutes, crushed and homogenized for one minute in 100 µl of distilled water in micro-tube. Then, one milliliter of suspension was added and smeared onto nutrient bromo-thymol blue tri-phenyl-tetrazolium chloride agar (NBTA) medium (Lacey, 1997). After 48-72 h, all morphologically distinct colonies were purified through serially sub-culturing until uniform colonies in sense of size and morphology were obtained. The colonies were tested for pathogenicity on the last instar larvae of *G. mellonella*. To approach this, the isolates were inoculated in liquid LB medium and incubated for 24 hours at 28 ± 2 °C. 10 µl of 10^4 cells of each isolate was injected into the last instar larvae of wax moth through proleg by a Hamilton syringe. The pathogenic isolates that caused mortality on wax moth larvae were identified by biochemical and molecular methods.

Identification of bacteria

Pathogenic strains on larvae of *G. mellonella* were characterized by gram reaction, anaerobic growth, catalase test, gelatin and aesculin hydrolysis, DNase, arginine dehydrolysis, lecithinase and nitrate reduction activity, growth on salicin and Simmon citrate, indole production and H₂S production from cysteine according to Akhurst and Boemare (1983).

According to phenotypic results some strains were selected to confirm by molecular identification. Total genomic DNA of the bacterial strains was extracted by alkaline lysis method (Rademaker and De Bruijn, 1997). For amplification of *recA* gene, specific Rec A primers synthesized by Eurofins MWG Operon Corporation, Germany were used based on the sequence of *recombinase A (rec A)* in *Erwinia* species (Waleron *et al.*, 2002). The RecA-F and RecA-R sequences were 5'-GGTAAAGGGT CTATCATGCG-3' and 5'-CCTTCACCATACAT AATTTGGA-3', respectively. DNA amplification was done according to a conventional method described by Waleron *et al.* (2002).

The bacterial PCR products were sequenced directly in both directions at Macrogen, (South Korea). The obtained sequences were checked as mentioned in previous section and the checked sequences were submitted to GenBank data base

under accessions KT070193- KT070198, KT070200-201, KR491942. Five sequences from GenBank data base along with ten sequences from the present study were used for construction of phylogenetic tree. Multiple sequence alignment of selected data was performed using MEGA5 software (Tamura *et al.*, 2011). The clustering stability of the NJ tree was evaluated by bootstrap analysis of 500 data sets (Felsenstein, 1985).

Results

EPNs Identification

Heterorhabditis bacteriophora and *S. feltiae* were identified based on morphological and morphometrical characters of third-stage infective juveniles and adults (Table 2) which were generally in agreement with those published for these species (Nguyen and Smart, 1996; Stock and Hunt, 2005).

Table 1 Species and isolates of *Heterorhabditis bacteriophora*, *Steinernema feltiae* and species and subspecies of *Photorhabdus temperata*, *Photorhabdus luminescens* and *Xenorhabdus bovienii* used in phylogenetic tree construction in this study.

| Species | Locality | Accession number | Reference |
|---|-----------------------------------|------------------|-------------------------------------|
| <i>H. bacteriophora</i> | Washington, USA | KT443980.1 | Fu <i>et al.</i> (2015) |
| <i>H. bacteriophora</i> | Switzerland | KJ938576.1 | Fesselet <i>et al.</i> (2014) |
| <i>H. bacteriophora</i> | Iran | KC675180.1 | Kamali and Karimi (2013) |
| <i>H. bacteriophora</i> | Bulgaria | JX993984.1 | Petrova <i>et al.</i> (2012) |
| <i>H. bacteriophora</i> | Florida, USA | JX403718.1 | Akyazi <i>et al.</i> , 2012) |
| <i>H. bacteriophora</i> | Mashhad, Khorasan, Iran | JX164230.1 | Hassani-Kakhki <i>et al.</i> (2012) |
| <i>H. bacteriophora</i> | Marivan, Kurdistan province, Iran | KR152294 | Present Study |
| <i>H. bacteriophora</i> | Qorveh, Kurdistan province, Iran | KT070192 | Present Study |
| <i>H. bacteriophora</i> | West Azarbaijan, Iran | EU598228 | Eivazian Kary <i>et al.</i> (2009) |
| <i>H. bacteriophora</i> | Ardabil, Iran | Eu598231 | Eivazian Kary <i>et al.</i> (2009) |
| <i>H. bacteriophora</i> | East Azarbaijan, Iran | Eu598232 | Eivazian Kary <i>et al.</i> (2009) |
| <i>H. bacteriophora</i> | East Azarbaijan, Iran | EU516355 | Eivazian Kary <i>et al.</i> (2009) |
| <i>H. bacteriophora</i> | East Azarbaijan, Iran | EU598222 | Eivazian Kary <i>et al.</i> (2009) |
| <i>H. bacteriophora</i> | Tehran, Iran | EU163272 | Karimi <i>et al.</i> (2007) |
| <i>S. feltiae</i> | Mashhad, Khorasan Razavi, Iran | JN098449.1 | Safari <i>et al.</i> (2011) |
| <i>S. feltiae</i> | Mashhad, Khorasan Razavi, Iran | JN098450.1 | Safari <i>et al.</i> (2011) |
| <i>S. feltiae</i> | Kyiv, Ukraine | KF939329.1 | Yakovlev <i>et al.</i> (2014) |
| <i>S. feltiae</i> | Hokkaido, Erimo, Japan | AB243439.1 | Kuwata and Yoshiga, 2007) |
| <i>S. feltiae</i> | Qorveh, Kurdistan province, Iran | KT070191 | Present Study |
| <i>S. feltiae</i> | East Azerbaijan, Iran | EU598240 | Eivazian Kary <i>et al.</i> (2008) |
| <i>S. feltiae</i> | Armenia | AY171256 | Spiridonov <i>et al.</i> (2013) |
| <i>S. feltiae</i> | Karaj, Iran | FJ657532 | Karimi <i>et al.</i> (2009) |
| <i>S. feltiae</i> | Tehran, Iran | FJ770381 | Karimi <i>et al.</i> (2009) |
| <i>P. temperate</i> subsp <i>thracensis</i> | France | FJ862016.1 | Tailliez <i>et al.</i> (2010) |
| <i>P. temperate</i> subsp <i>thracensis</i> | Kurdistan province, Iran | KT070201 | Present Study |
| <i>P. temperate</i> subsp <i>thracensis</i> | Kurdistan province, Iran | KT070200 | Present Study |
| <i>P. temperate</i> subsp <i>thracensis</i> | Kurdistan province, Iran | KT070194 | Present Study |
| <i>P. temperate</i> subsp <i>thracensis</i> | Kurdistan province, Iran | KT070199 | Present Study |
| <i>P. luminescens</i> subsp <i>kayayii</i> | France | FJ861996.1 | Tailliez <i>et al.</i> (2010) |
| <i>P. luminescens</i> subsp <i>kayayii</i> | Kurdistan province, Iran | KT070193 | Present Study |
| <i>P. luminescens</i> subsp <i>kayayii</i> | Kurdistan province, Iran | KR491942 | Present Study |
| <i>X. bovienii</i> | Australia | U87924.1 | Pinyon <i>et al.</i> (2000) |
| <i>X. bovienii</i> | Lebanon | LN835358.1 | Noujeim <i>et al.</i> (2015) |
| <i>X. bovienii</i> | Australia | FJ823426 | Tailliez <i>et al.</i> (2010) |
| <i>X. bovienii</i> | Kurdistan province, Iran | KT070195 | Present Study |
| <i>X. bovienii</i> | Kurdistan province, Iran | KT070196 | Present Study |
| <i>X. bovienii</i> | Kurdistan province, Iran | KT070197 | Present Study |
| <i>X. bovienii</i> | Kurdistan province, Iran | KT070198 | Present Study |

In terms of EPNs recovery 60% of collected soil samples were positive. *H. bacteriophora* was recovered from 59.33% of soil samples while only 0.66 % of samples contained *S. feltiae*. Grasslands habitat showed higher recovery frequency than other habitats. Incidence of *H. bacteriophora* in grasslands, oak forests and alfalfa fields were 57.30, 14.61 and 28.08%, respectively. *S. feltiae* was only found in an alfalfa field in Qorveh (Fig. 1).

The amplification of rRNA-ITS regions plus 5.8S gene yielded a single fragment of approximately 800 bp and 750 bp for *H. bacteriophora* and *S. feltiae*, respectively based on gel electrophoresis. BLAST search from the NCBI showed 100% homology in the ITS sequences of the isolates of *H. bacteriophora*, from Marivan (KR152294) and Qorveh (KT070192) with isolates of Mashhad, Khorasan

(JX164230, KC675180), USA, Switzerland, Bulgaria (KT443980, KJ938576 JX993984, JX403718). *Steinernema feltiae* isolated from Qorveh, Kurdistan (KT070191) showed 99% similarity with those sequences of *S. feltiae* deposited in GenBank i.e. JN098449, JN098450 from Mashhad, Iran, AB243439 from Japan and KF939329 from Ukraine. In the phylogenetic tree (Fig. 2) all the isolates of *H. bacteriophora* used for tree construction generated one clade, all of them with good posterior probability values, the newly sequenced isolates of *H. bacteriophora* from Kurdistan showed high homology with the other isolates. In the phylogenetic tree constructed from the ITS-rRNA gene sequences of the isolates of *S. feltiae* one clade was formed. The isolate from Qorveh, Kurdistan formed a sister group with the isolates from Khorasan, Iran, and the rest of *S. feltiae*.

Table 2 Morphometric and morphologic characters of entomopathogenic nematodes isolated from Kurdistan Province.

| Characters | <i>Steinernema feltiae</i> ¹ | | | <i>Heterorhabditis bacteriophora</i> ¹ | | |
|----------------------------------|---|----------------------------------|------------|---|---------------------------------|---------------------------------|
| | IJ | Male | Female | IJ | Male | Female |
| N | 15 | 5 | 5 | 15 | 5 | 5 |
| Total body Length | 862.14 ± 47.42 (814.72 - 909.56) | | | 619.37 ± 29.66 (589.71 - 649.03) | | |
| Excretory pore (EP) | 60.35 ± 4.25 (56.10 - 64.60) | 75 ± 6 (69 - 81) | | 102.69 (99.71 - 105.67) | | 90 ± 2.00 (88 - 92) |
| Tail length (TL) | 79.14 ± 6.90 (72.24 - 86.04) | | | 94.12 ± 5.70 (88.42 - 99.82) | | |
| Oesophagus Length | 123.50 ± 8.64 (114.86 - 132.14) | 152 ± 13.73 (138.27 - 165.73) | | 125.94 ± 5.49 (125.50 - 131.43) | | 117 ± 2.16 (115.84 - 119.16) |
| Maximum Body width | 26 ± 2.25 (23.75 - 28.25) | | | 25±1.41 (23.59 - 26.41) | | |
| D% (EP/OesophaguS length×100) | 48.973 (45.79 - 52.15) | 49.61 ±5.88 (43.73 - 55.49) | | 81.65 ± 3.51 (78.14 - 85.16) | | 77.19 ± 1.75 (75.44 - 78.94) |
| E% (EP/TL × 100) | 76.52 ± 5.27 (71.25 - 81.79) | | | 109.42 ± 6.35 (103.07 - 115.77) | | |
| Spicule length (Sp L) | | 71.30 ± 2.06 (69.24 - 73.36) | | | 39.6 ± 1.14 (38.46 - 40.74) | |
| Gubernaculum Length | | 41.50 ± 1.29 (40.29 - 42.79) | | | 22.40 ± 2.07 (20.33 - 24.47) | |
| Testis reflexion | | | | | 76.60 ± 2.79 (73.81 - 79.39) | |
| SW (SPL/cloacal Body width) | | 1.30 ± 026 (1.04 - 1.56) | | | | Present |
| Post - anal Swelling | | | | | | Present |
| Vulval lips | | | Protruding | | | |
| Mucro | | | Present | | | |
| Epiptygma | | | Present | | | |

¹. Measurements in µm.




Figure 1 Distribution of entomopathogenic nematodes in Kurdistan province.




Figure 2 Bayesian phylogenetic tree inferred from ITS-rRNA gene sequences of some *Heterorhabditis bacteriophora* and *Steinernema feltiae*. Posterior probabilities more than 70% are given for appropriate clades. Name and accession number of sequences of current study are in bold.

Bacterial Identification

All bacterial strains showed the common phenotypic characters of the *Xenorhabdus* and *Photorhabdus* genera including granular and translucent forms. Results of biochemical tests are showed in table 3, indicating 22 strains having characteristics close to *Photorhabdus*, whereas, 14 strains from *S. feltiae* are related to *Xenorhabdus*. The catalase test was negative for *Xenorhabdus*, however, it was positive for *Photorhabdus* as it was expected. Eight strains of two genera were subjected to molecular identification to confirm the biochemical diagnosis.

Table 3 Results of biochemical experiments of isolated symbiosis bacteria.

| Test | <i>Xenorhabdus</i> (14 strains) | <i>Photorhabdus</i> (22 strains) |
|--------------------------------|------------------------------------|-------------------------------------|
| Gram | - | - |
| Anaerobic growth | + | + |
| Catalase | - | + |
| Gelatin hydrolysis | + | + |
| Aesculin hydrolysis | - | ± |
| Salicin | ± | ± |
| Lecitinase activity | - | - |
| Nitrate reduction | - | - |
| DNA's activity | + | - |
| Growth on simmon citrate | + | ± |
| Indol production | - | - |
| H ₂ S from cysteine | + | - |
| Arginine dehydrolysis | - | - |

+: 80% or more strains positive, -: 80% or more strains negative, ±: variable.

The amplification of *recA* gene produced a fragment of 730 bp. Genetic sequence analysis in a nucleotide blast (BLASTn) search of strains KT070193 (PLIRI7) and KR491942 (Kurd1) revealed 100% similarity to *Photorhabdus luminescens* subsp. *kayaii* (FJ861996.1) and strains KT070201 (PTIRI9), KT070200 (PTIRI8), KT070194 (PTIRI6) and KT070199 (PTIRI10) to *Photorhabdus temperata* subsp. *thracensis* (FJ862016.1). The *recA* sequences of three isolates of *Xenorhabdus* (KT070195 (XBIRI5), KT070196 (XBIRI4), KT070197 (XBIRI3) and KT070198 (XBIRI2)) showed 99% homology with those of *Xenorhabdus bovienii* presented in the database (U87924.1, LN835358, FJ823426).

Figure 3 shows the phylogentic tree constructed using sequences of previously isolated strains and from the new isolates of this survey. Groups 1, 2 and 3 include all *P. temperate* subsp. *thracensis*, *P. luminescens* subsp. *Kayaii* and *X. bovienii* strains, respectively. Groups 1 and 2 representing sister group that their differences may be enough to separate them as two subspecies. Group 3 shows an out-group to groups 1 and 2 that illustrate high genetically differences to two other groups.




Figure 3 Neighbor joining Phylogenetic tree (Saitou, N. and Nei, 1987) showing affiliation of the symbiotic bacteria of entomopathogenic nematodes to reference data on the basis of *recA* gene sequence alignment. Bootstrap value (500 replicates) above 50% are shown next to the branches. Bolded strains show name and accession number of sequences of current study.

Discussion

Two species of entomopathogenic nematodes, *H. bacteriophora* and *S. feltiae* were recovered from soil samples collected from Kurdistan province. The morphological identification was confirmed by molecular data. *H. bacteriophora* was the dominant species and showed remarkably high frequency in this region. In this study, more than half of the collected samples (60%) were positive for EPNs, with a predominance of *H. bacteriophora* over *S. feltiae*. The 60% prevalence of entomopathogenic nematodes in Kurdistan province is a new record for the natural occurrence of these nematodes in Iran. The majority of the positive samples were isolated from grasslands habitat, in contrast to forests where oak species constitute most of the woodlands. This rate is much higher than those reported 3% in Arasbaran (Nikdel *et al.*, 2010), 3.2% in North West (Eivazian Kary *et al.*, 2009), 1% and 2% in North Khorasan (Kamali *et al.*, 2013). A study conducted in California showed the recovered isolates from woodlands coniferous and oak forests had the majority of the positive samples, 67.60 percent of samples were positive for presence of EPNs, in some regions all samples were positive (Stock *et al.*, 1999).

This study showed entomopathogenic nematodes particularly *H. bacteriophora* are widespread in Kurdistan Province. The soil samples were principally taken from natural and virgin areas with enough humidity and vegetation, which provides more stable conditions for growth and establishment of EPN populations (Campos-Herrera *et al.*, 2008).

High frequency of occurrence of EPNs in Kurdistan province is likely due to favorable environmental conditions in this region. Humidity and temperature have a significant effect on survival rates of EPNs. Kurdistan region has a temperate to cold climate with mean annual minimum and maximum temperatures of 11 to 14 °C. Meteorology information demonstrates that absolute maximum temperature and absolute minimum

temperature in sampling locations were 35 to 42 and -10 to -24, respectively in recent years. The mean annual precipitation of this region ranges from 347 mm in Qorveh to 728 mm in Marivan (Meteorology organization of Kurdistan, 2015). These conditions could reflect the abundance of EPNs and adaptability of particularly *H. bacteriophora* to the natural environmental conditions and habitats of this Province. Temperatures above 40 °C and below 8 °C are lethal for most EPNs (Griffin, 1993; Grewal *et al.*, 1994). Extremely high soil temperatures are rare in regions like Kurdistan province, but temperatures below 8 °C are common in this area and can be considered as a limiting factor.

The EPN species share mutual relationships with a single bacterial species; however, some species especially *Xenorhabdus* spp. are associated with more than one EPN species. In this study, two subspecies, *P. temperata* subsp. *thracensis* and *P. luminescens* subsp. *kayaii* were isolated from *H. bacteriophora*. *Xenorhabdus bovienii* was isolated from *S. feltiae*. Two isolates of *Xenorhabdus*, *X. nematophila* and *X. poinarii* were isolated from *S. glaseri* and *S. carpocapsae* collected from Tehran province, Iran. The bacterial isolates were characterized by their phenetic characters and sequences of 16S ribosomal RNA gene (Karimi *et al.*, 2011). Four isolates of *P. luminescens* subsp. *lumondii*, *X. bovienii*, *X. nematophilus* and *X. budapestensis* were reported and identified by 16S-rRNA gene sequences (Agazadeh *et al.*, 2010). Two species of *Photorhabdus* were identified and recorded based on *recA* gene for the first time from Iran. *P. temperata* subsp. *thracensis* and *P. luminescens* subsp. *kayaii* were reported from *H. bacteriophora* from Turkey and Lebanon (Hazir *et al.* 2004; Noujeim *et al.*, 2011). The isolation of these symbiotic bacteria for completing their phylogeny and clarifying the ambiguous aspects of their characteristics is inevitable. Due to variation of subspecies, the obtained data from this study viz. chemical characteristics and sequences can be very helpful in the future

research programs. Studies on signals between bacteria and nematodes and probability of substitution of robust bacteria for improving the lethal efficacy can be considered to develop the ambiguous aspects of their relationships. Surveying on factors that cause the specificity in nematode-bacteria interactions will provide information on how to manipulate this symbiotic relationship to accelerate their pathogenicity and broaden the insect host range.

The nematode isolates recovered from this survey can contribute to the biological control programs using native isolates in Kurdistan and adjacent provinces with similar climatic conditions.

Acknowledgment

This research was funded by Mohaghegh Ardabili University, Ardabil, Iran and Iranian Research Institute of Plant Protection. Authors acknowledge Dr. Javad Karimi for helpful advice in isolation of bacteria.

References

- Agazadeh, M., Mohammadi, D. and Eivazian Kary, N. 2010. Molecular identification of Iranian isolates of the genus *Photorhabdus* and *Xenorhabdus* (Enterobacteriaceae) based on 16S rRNA. *Munis Entomology and Zoology*, 5 (2): 772-779.
- Akhurst, R. J. and Boemare, N. E. 1983. Genus XL. *Xenorhabdus*. In: Brenner, D. J., Krieg, N. R., Staley, J. T. and Garrity, G. M. (Eds.), *Bergey's Manual of Systematic Bacteriology*. Vol. 2. Springer, New York, pp. 831-838.
- Akhurst, R. J. and Boemare, N. E. 1988. A numerical taxonomic study of the genus *Xenorhabdus* (Enterobacteriaceae) and proposed elevation of the subspecies of *X. nematophilus* to species. *Journal of General Microbiology*, 134: 1835-1845.
- Bedding, R. A. and Akhurst, R. J. 1975. A simple technique for the detection of insect parasitic rhabditid nematodes in soil. *Nematologica*, 21: 109-110.
- Boemare, N. and Akhurst, P. 2006. The genera *Photorhabdus* and *Xenorhabdus*. *Prokaryotes*, 6: 451-494.
- Bronskill, J. 1961. A cage to simplify the rearing of the greater wax moth, *Galleria mellonella* (Pyralidae). *Journal of the Lepidopterists Society*, 15: 102-104.
- Bussaman, P., Sobanboa, S., Grewal, P. S. and Chandrapatya, A. 2009. Pathogenicity of additional strains of *Photorhabdus* and *Xenorhabdus* (Enterobacteriaceae) to the mushroom mite *Luciaphorus perniciosus* (Acari: Pygmephoridae). *Applied Entomology and Zoology*, 44 (2): 293-299.
- Campos-Herrera, R., Gomez-Ros, J. M., Escuer, M., Cuadra, L., Barriosc, L. and Gutierrez, C. 2008. Diversity, occurrence, and life characteristics of natural entomopathogenic nematode populations from La Rioja (Northern Spain) under different agricultural management and their relationships with soil factors. *Soil Biology and Biochemistry*, 40: 1474-1484.
- Ciche, T. A. and Ensign, J. C. 2003. For the insect pathogen *Photorhabdus luminescens*, which end of a nematode is out? *Applied and Environmental Microbiology*, 69: 1890-1897.
- De Grisse, A. T. 1969. Redescription ou modifications de quelques techniques utilisées dans l'étude des nématodes phytoparasitaires. *Mededelingen van de Rijksfaculteit Landbouwwetenschappen Gent*, 34: 351-369.
- Eisen, J. A. 1995. The RecA protein as a model molecule for molecular systematic studies of bacteria: comparison of trees of RecAs and 16S rRNAs from the same species. *Journal of Molecular Evolution*, 41: 1105-1123.
- Eivazian Kary, N., Mohammadi, S. A., Moghaddam, M., Nikdel, M. and Niknam, G. 2009. A survey of entomopathogenic nematodes of the families Steinernematidae and Heterorhabditidae (Nematoda: Rhabditida) from north-west of Iran. *Nematology*, 11: 107-116.

- Euzeby, J. P. and Boemare, N. E. 2000. The modern Latin word *rhabdus* belongs to the feminine gender, inducing necessary corrections according to Rules 65 (2), 12c (1), and 13b of the Bacteriological Code (1990 Revision). *International Journal of Systematic and evolutionary Microbiology*, 50: 1691-1692.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39: 783-791.
- Forst, S., Dowds, B., Boemare, N. and Stackebrandt, E. 1997. *Xenorhabdus* spp. and *Photorhabdus* spp.: bugs that kill bugs. *Annual Review of Microbiology*, 51: 47-72.
- Grewal, P. S., Selvan, S. and Gaugler, R. 1994. Thermal adaptation of entomopathogenic nematodes: niche breadth for infection, establishment, and reproduction. *Journal of Thermal Biology*, 19: 245-253.
- Griffin, C. T. 1993. Temperature responses of entomopathogenic nematodes: Implications for the success of biological control programmes, In: Bedding, R. and Kaya, H. (Eds.), *Nematodes and the biological Control of Insect Pests*. Csiro Publishing, Melbourne, pp. 115-126.
- Grimont, P. A. D., Steigerwalt, A. G., Boemare, N., Hickman-Brenner, F. W., Deval, C., Grimont, F. and Brenner, D. J. 1984. Deoxyribonucleic acid relatedness and phenotypic study of the genus *Xenorhabdus*. *International Journal of Systematic and Evolutionary Microbiology*, 34: 378-388.
- HALL, T. A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucleic Acids Symposium Series*, 41: 95-98.
- Hazir, S., Stackebrandt, E., Lang, E., Schumann, P., Ehlers, R. U. and Keskin, N. 2004. Two new Subspecies of *Photorhabdus luminescens*, Isolated from *Heterorhabditis bacteriophora* (Nematoda: Heterorhabditidae): *Photorhabdus luminescens* subsp. *kayaii* subsp. nov. and *Photorhabdus luminescens* subsp. *thracensis* subsp. nov. *System. Journal of Applied Microbiology*, 27: 36-42.
- Huelsenbeck, J. P. and Ronquist, F. 2001. MrBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17: 754-755.
- Iraki, N., Salah, N., Sansour, M. A., Segal, D., Glazer, I., Johnigk, S. A., Hussein, M. A. and Ehlers, R. U. 2000. Isolation and characterization of two entomopathogenic nematode strains, *Heterorhabditis indica* (Nematoda, Rhabditida), from the West Bank, Palestinian Territories. *Journal of Applied Entomology*, 124: 375-380.
- Jang, E. K., Ullah, I., Kim, M. S., Lee, K. Y. and Shin, J. H. 2011. Isolation and characterization of the entomopathogenic bacterium, *Photorhabdus temperate* producing a heat stable insecticidal toxin. *Journal of Plant Diseases and Protection*, 118 (5): 178-184.
- Joyce, S. A., Reid, A., Driver, F. and Curran, J. 1994. Application of polymerase chain reaction (PCR) methods to identification of entomopathogenic nematodes. *Proceedings of Symposium and Workshop*, St. Patrick's College, Maynooth, Co. Kildare, Ireland, Luxembourg, European Commission, DG XII, p. 178.
- Kamali, S., Karimi, J., Hosseini, M., Campos-Herrera, R. and Duncan, L.W. 2013. Biocontrol potential of the entomopathogenic nematodes *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* on cucurbit fly, *Dacus ciliatus* (Diptera: Tephritidae). *Biocontrol Science and Technology*, 23 (11): 1307-1323.
- Karimi J. and Kharazi-pakdel, A. 2007. Incidence of natural infection of the white grub *Polyphylla olivieri* (Coleoptera: Scarabaeidae) with entomopathogenic nematodes in Iran. *Bulletin OILB-SROP*, 30 (1): 35-39.
- Karimi, J., Kharazi-Pakdel, A., Yoshiga, T., Koohi Habibi, M. and Hassani-Kakhki, M. 2011. Characterization of *Xenorhabdus* (γ -Proteobacteria) strains associated bacteria with the *Steinernema* (Nematoda: Steinernematidae) isolates from Iran. *Journal of Entomological Society of Iran*, 31 (1): 57-69.

- Kaya, H. K., Gaugler, R. 1993. Entomopathogenic nematodes. Annual Review of Entomology, 38: 181-206.
- Lacey, L. A. 1997. Manual of Techniques in Insect Pathology. Academic Press, London.
- Ludwig, W. and Schleifer, K. 1999. Phylogeny of bacteria beyond the 16S rRNA standard. American Society for Microbiology News, 65: 752-757.
- Martens, E. C., Heungens, K. and Goodrich-Blair, H. 2003. Early colonization events in the mutualistic association between *Steinernema carpocapsae* nematodes and *Xenorhabdus nematophila* bacteria. Journal of Bacteriology, 185: 3147-3154.
- Meteorology organization of Kurdistan. (2015). Available on: <http://www.kurdistanmet.ir/> (accessed July 8, 2015).
- Nguyen, K. B. and Smart, J. R. 1996. Identification of entomopathogenic nematodes in the Steinernematidae and Heterorhabditidae (Nemata: Rhabditida). Journal of Nematology, 28: 286-300.
- Nikdel, M., Niknam, G., Griffin, C. T. and Eivazian Kary N. 2010. Diversity of entomopathogenic nematodes (Nematoda: Steinernematidae, Heterorhabditidae) from Arasbaran forests and rangelands in north-west Iran. Nematology, 12: 767-773.
- Noujeim, E., Khater, C., Pages, S., Ogier, J. C., Tailliez, P., Hamze, M. and Thaler, O. 2011. The first record of entomopathogenic nematodes (Rhabditia: Steinernematidae and Heterorhabditidae) in natural ecosystems in Lebanon: A biogeographic approach in the Mediterranean region. Journal of Invertebrate Pathology, 107: 82-85.
- Parvizi, R. 2001. Study infectivity of *H. bacteriophora* and *Steinernema* sp. against white grub, *Polyphylla olivieri*. Journal of Entomological Society of Iran, 21: 63-72.
- Poinar G. O. Jr. 1990. Biology and taxonomy of Steinernematidae and Heterorhabditidae, In: Gaugler, R. and Kaya, H. K. (Eds.), Entomopathogenic Nematodes in Biological Control. CRC Press, Boca Raton, FL, USA, pp. 23-62.
- Rademaker, J. L. W. and De Bruijn, F. J. 1997. Characterization and classification of microbes by rep-PCR genomic fingerprinting and computer assisted pattern analysis, In: Caetano-Anollés, G. and Gresshoff, P. M. (Eds.), DNA Markers: Protocols, Applications and Overviews. J. Wiley and Sons, Inc., USA, pp. 151-171.
- Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for constructing phylogenetic tree. Molecular Biology and Evolution, 4: 406-425.
- Spiridonov, S. E., Reid, A. P., Podrucka, K. Subbotin, S. A. and Moens, M. 2004. Phylogenetic relationships within the genus *Steinernema* (Nematoda: Rhabditida) as inferred from analyses of sequences of the ITS1-5.8S-ITS2 region of rDNA and morphological features. Nematology, 6: 547-566.
- Stock, S. P. and Hunt, D. J. 2005. Morphology and systematics of Nematodes used in Biocontrol, In: Grewal, P. S., Ehlers, R. U. and Shapiro-Ilan, D. I. (Eds.), Nematodes as Biocontrol Agents. CABI Publishing, Wallingford, pp. 3-46.
- Stock, S. P. 2002. New trends in entomopathogenic nematode systematics: impact of molecular biology and phylogenetic reconstruction. Proceedings of the International Conference on Parasitology. Vancouver, Canada, p. 1.
- Stock, S. P., Pryor, P. M. and Kaya, H. K. 1999. Distribution of entomopathogenic nematodes (Steinernematidae and Heterorhabditidae) in natural habitats in California, USA. Biodiversity and Conservation, 8: 535-549.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei M. and Kumar, S. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Molecular Biology and Evolution, 28: 2731-2739.
- Tanha Maafi, Z., Ebrahimi, N., Abootorabi, E. and Spiridonov, S. E. 2006. Record of two Steinernematid species from Iran. 17th Iranian Plant Protection Congress, Karaj, Iran, p. 482.

- Tanha Maafi, Z., Subbotin, S. A. and Moens, M. 2003. Molecular identification of cyst-forming nematodes (Heteroderidae) from Iran and a phylogeny based on ITS-rDNA sequences. *Nematology*, 5: 99-111.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G. 1997. The CLUSTALX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, 25: 4876-4882.
- Waleron, M., Waleron, K., Podhajska, A. J. and Lojkowska, E. 2002. Genotyping of bacteria belonging to the former *Erwinia* genus by PCR-RFLP analysis of a *recA* gene fragment. *Microbiology*, 148: 583-595.
- Wang, Y. and Gaugler, R. 1998. Host and penetration site location by entomopathogenic nematodes against Japanese beetle larvae. *Journal of Invertebrate Pathology*, 72: 313-318.
- White, G. F. 1927. A method for obtaining infective nematode larvae from cultures. *Science*, 66: 302-303.

جداسازی و شناسایی نماتدهای بیمارگر حشرات و باکتری‌های هم‌زیست آنها از استان کردستان ایران

آرمان عبدالملکی^۱، زهرا تنهامعافی^{۲*}، هوشنگ رفیعی دستجردی^۱، بهرام ناصری^۱ و ابوالقاسم قاسمی^۲

۱- گروه گیاه‌پزشکی، دانشکده علوم کشاورزی، دانشگاه محقق اردبیلی، اردبیل، ایران.

۲- مؤسسه تحقیقات گیاه‌پزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: tanhamaafi@iripp.ir

دریافت: ۲۰ تیر ۱۳۹۴؛ پذیرش: ۱ اسفند ۱۳۹۴

چکیده: نماتدهای بیمارگر حشرات به‌عنوان عوامل بیولوژیکی در کنترل آفات کشاورزی مورد استفاده قرار می‌گیرند. این مطالعه به‌منظور شناسایی و تعیین فراوانی نماتدهای بیمارگر حشرات در استان کردستان و شناسایی و تعیین خصوصیات باکتری‌های هم‌زیست آنها انجام شد. شناسایی نماتدها براساس ویژگی‌های ریخت‌شناسی، ریخت‌سنجی و توالی نواحی ITS ژن rRNA انجام گرفت. دو گونه‌ی *Heterorhabditis bacteriophora* (Rhabditida: Heterorhabditidae) و *Steinernema feltiae* (Steinernematidae) شناسایی شدند. از تعداد ۱۵۰ نمونه‌ی خاک جمع‌آوری شده از نیمه‌ی جنوبی استان کردستان ۶۰ درصد آنها از نظر حضور نماتدهای بیمارگر حشرات مثبت بودند. گونه‌ی *H. bacteriophora* بیش‌ترین فراوانی را در این منطقه داشت به‌طوری‌که ۵۹/۳ درصد نمونه‌های خاک دارای این گونه بودند. حضور گونه‌ی *H. bacteriophora* در علف‌زارها، مناطق جنگلی و مزارع یونجه به‌ترتیب ۵۷/۳، ۱۴/۶ و ۲۸ درصد بود. گونه‌ی *S. feltiae* فقط در مزرعه یونجه با فراوانی ۰/۶۶ درصد یافت شد. شناسایی باکتری‌های هم‌زیست گونه‌های *H. bacteriophora* و *S. feltiae* بر پایه‌ی خصوصیات بیوشیمیایی و توالی‌یابی ژن *recA* انجام شد. در این مطالعه دو گونه از جنس *Photorhabdus* با نام‌های *P. luminescens* subsp. *kayaii* و *P. temperata* subsp. *thracensis* از *H. bacteriophora* جداسازی و شناسایی گردید. باکتری *Xenorhabdus bovienii* نیز از لاروهای بیماری‌زای *S. feltiae* تشخیص داده شد.

واژگان کلیدی: *Steinernema feltiae*، *Photorhabdus luminescens*، *Heterorhabditis bacteriophora*

Xenorhabdus bovienii، فیلوژنی