

Research Article

## Resistance of tomato cultivars to *Tuta absoluta* (Lepidoptera: Gelechiidae) under field condition

Fatemeh Azadi<sup>1</sup>, Ali Rajabpour<sup>1\*</sup>, Amin Lotfi Jalal Abadi<sup>2</sup> and Mehdi Mahjoub<sup>3</sup>

1. Department of Plant Protection, Faculty of Agriculture, Khuzestan Agricultural Sciences and Natural Resources University, Ramin, Iran.

2. Department of Agronomy and Plant Breeding, Faculty of Agriculture, Khuzestan Agricultural Sciences and Natural Resources University, Ramin, Iran.

3. Department of Plant Protection, Agricultural and Natural Resources Research Center of Kermanshah, Kermanshah, Ramin, Iran.

**Abstract:** Tomato leaf miner, *Tuta absoluta*, is the most destructive pest of tomato in Iran. Resistance levels of six tomato cultivars to the pest damage were evaluated during two tomato growing seasons (2014-2015). Samplings were done weekly. Leaf and fruit damages as well as the total yield of the cultivars were compared. Furthermore, leaf trichome density was evaluated. The highest to lowest leaf infestation rates were recorded for the Primo early, CaljN3, Petomek, Rio grande, Early urbana and Super 2270 cultivars respectively. Fruit infestation rate in Early urbana was significantly lower than the other cultivars in both growing seasons. Total yield of tomato (from the highest to the lowest) belonged to Super 2270, Early urbana, Rio grande, Petomek, Calj N3 and Primo early cultivars.

**Keywords:** Tomato leaf miner, Host plant resistance, Leaf and fruit damage, Ramin, Khuzestan

### Introduction

The tomato leaf miner (TLM), *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae), is considered as one of the most dangerous pests in greenhouse and field grown tomatoes. The pest is an invasive pest native to South America (Yankova, 2012) and it is newly distributed in Iran (Baniameri and Cheraghian, 2011). The female usually lays eggs on leaves, stems, and to a lesser extent on fruits. The young larvae mine the leaves or stems producing large galleries and burrow into the fruit. On leaves, the larvae feed only on mesophyll cells, leave the epidermis intact and make irregular leaf

mines, which may later become necrotic (OEPP/EPP, 2005) and affect photosynthesis in the plant (Desneux *et al.*, 2010). Damage from this pest throughout the entire growing cycle of tomatoes can significantly reduce both yield and fruit quality by the direct feeding of *T. absoluta* and secondary pathogens that may enter through the wounds made by the insect. In the absence of control strategies, larval feeding damage can reach up to 100% (Yankova, 2012). Large amounts of chemical insecticides have been applied against the pest in both field and greenhouse (Lietti *et al.*, 2005). The applications cause many problems such as increase the insecticide costs, destroy natural enemy populations, leave pesticide residues in fruits and make the pest population resistant to chemical compounds (Braham and Hajji, 2012). Also, the general endophytic behaviour of the larval instars makes it difficult to conduct

Handling Editor: Yaghsoub Fathipour

\* **Corresponding author**, e-mail: rajabpour@ramin.ac.ir  
Received: 6 February 2018, Accepted: 15 April 2018  
Published online: 29 April 2018

effective control practices against this pest (Lietti *et al.*, 2005; Guedes and Picanço, 2012).

Host plant resistance is economic and ecologically friendly strategy in integrated pest management (IPM) programs (Pedigo, 2002; Sharma and Ortiz, 2002). Use of cultivars resistant to the pest cause a major reduction in chemical insecticide application that can lead to increase in beneficial organism activity and decrease of pesticide residues in the environment and human food (Sharma and Ortiz, 2002). In many cases, use of resistant host plant is nearby compatible with other control strategies in IPM programs (Pedigo, 2002). Finding cultivars resistant to a pest is the primary step in the use of host plant resistance in IPM programs (Panda and Khush, 1995). Host plant defense in resistant cultivars may be related to morphological, biochemical and molecular traits to counter/offset herbivore attack (War *et al.*, 2012). Trichomes are hair-like appendages that develop from cells of the aerial epidermis and are produced by most plant species and contribute to host plant resistance against herbivore insects (Dalin *et al.*, 2008).

Resistance levels of some tomato cultivars to *T. absoluta* have been previously investigated by some researchers (Gharekhani and Salek-Ebrahimi, 2014a, b; Proffit *et al.*, 2011; Oliveira *et al.*, 2009; Sohrabi *et al.*, 2016; Ghaderi *et al.*, 2017). In this study, resistance level of six tomato cultivars to *T. absoluta* were studied in field condition. Also, the effect of leaf trichome on the resistance levels was evaluated.

## Materials and Methods

### Experimental design

Experiments were done in an experimental field, 5000m<sup>2</sup>, in Chogha Kabod region, Kermanshah, West Iran, during two tomato growing seasons 2015-2016. Six tomato cultivars including Rio gande, Super 2270, Calj N3, Early Urbana, Primo early and Petomek were cultivated (18000 plants per hectare). Between each plot (150m<sup>2</sup>) a ridge (2m) was made. Experiments were arranged in a randomized complete block design with four

replications (plots). Cultural practices were conducted according to instructions of the Kermanshah province agricultural organization.

### Sampling

Sampling was performed weekly from primary phenological step, 5-7 leaves, to end of the growing season. At each sampling date, ten randomly selected plants were checked by traveling in an X-shaped pattern through each plot. From each selected plant, three leaves and fruit were randomly chosen and the numbers of the pest larval mines in each leaf and fruit were separately recorded. At the end of growing season, total fruit yield in each plot was separately weighed.

### Effect of leaf trichome

For evaluation of leaf trichome effect on host plant resistance level, three cultivars with high, moderate and the low resistance level were chosen. Seeds of the cultivars were sown in plastic pots (12cm diameter and 22cm height). Ten pots were provided for each selected cultivar. The pots were kept in germinator at  $27 \pm 2$  °C, photoperiod 16: 8 (light: dark) and RH  $60 \pm 5$ %. At the eight leaf stage, six leaves were randomly detached from the potted plants. A paper quadrat, 5 × 5 mm, was randomly placed under each leaf and number of trichomes in each quadrat was recorded under stereomicroscope.

### Data analysis

The data were statistically analyzed by one way analysis of variance (one-way ANOVA) and the group means were compared by Duncan's multiple range test using SAS program (SAS Institute, 2003).

## Result

### Leaf and fruit damage

Leaf and fruit Infestation rates in various experimental cultivars during first (2015) and second (2016) growing seasons are presented in Table 1. Significant differences were observed between leaf infestation rates in various experimental varieties. In both growing

seasons, infestation levels in Primo early were significantly higher than the other cultivars (22.7 and 26.6% in the first and second growing season, respectively). While, infestation of Super 2270 was significantly lower than other cultivars (9.5 and 10.3% in the two growing seasons, respectively). Totally, order of infestation rates to TLM (from the highest to the lowest) was Primo early, CaljN3, Petomek, Rio grande, Early urbana and Super 2270.

Fruit infestation rate between various experimental cultivars was significantly different. Fruit infestation rates in Early Urbana were significantly lower than other cultivars in both growing seasons (0.5 and 6.5% during the first and second growing season, respectively). The highest fruit infestation rate was observed in Calj N3 (4.82%) and Petomek (12.42%) during the first and second growing season, respectively.

**Table 1** Percentages of damaged leaf, damaged fruit and total yield in tomato cultivars in response to *Tuta absoluta*.

Cultivars	Damaged leaf (%)		Damaged fruit (%)		Total yield (kg/m <sup>2</sup> )	
	2015	2016	2015	2016	2015	2016
Primo early	22.7 ± 3.3a*	26.6 ± 6.4a	2.2 ± 0.4b	9.9 ± 1.0b	127.4 ± 8.5c	131.8 ± 8.8c
Calj N3	18.7 ± 0.5b	24.1 ± 2.4a	4.8 ± 0.2a	7.6 ± 1.2c	133.6 ± 1.0c	141.9 ± 9.8bc
Early Urbana	8.0 ± 0.4d	12.4 ± 3.2b	0.5 ± 0.5e	6.5 ± 0.9c	112.6 ± 4.3d	157.0 ± 21.3b
Peto mek	18.0 ± 2.8b	22.7 ± 2.6a	1.6 ± 0.2c	12.4 ± 1.2a	178.9 ± 4.6a	156.0 ± 14.7b
Supper 2270	9.5 ± 1.9dc	10.4 ± 1.7b	1.0 ± 1.9d	7.8 ± 1.5c	175.4 ± 6.6a	207.3 ± 5.8a
Rio grande	12.0 ± 0.8c	14.7 ± 3.8b	1.5 ± 0c	8.4 ± 1.4bc	163.4 ± 2.8b	159.3 ± 14.8b
F (df = 5, 23)	33.29	13.55	106.8	8.5	93.6	12.22
P-value	< 0.001	< 0.001	< 0.001	0.0002	< 0.001	< 0.001

Means followed by the same letters in each column indicate non-significant differences (Duncan's Multiple Range Test,  $P \leq 0.05$ ).

### Total Yield

Total fruit yield of different cultivars in the first (2015) and second (2016) growing seasons is presented in Table 1.

Results indicated that there were significant differences between total yields of various cultivars. In both growing season, the highest fruit yield was obtained in Super 2270 (207.35 and 175.4 kg per plot and the lowest yield was that of Primo early (131.85 and 127.4 kg per plot). In the first growing season, total yield in decreasing order was that of Petomek, Super 2270, Rio grande, Calj n3, Primo early and Early Urbana. Whereas in in the second growing season it was that of Super 2270, Early Urbana, Rio grande, Petomek, Calj N3 and Primo early.

### Trichome density

According to the results of leaf infestation rates with TLM larvae, Super 2270, Early urbana and Primo early were selected as experimental cultivars with the highest, moderate and lowest resistance level to the pest, respectively. Trichome densities in these

cultivars indicate significant difference among them (Table 2). The highest and lowest trichome density were recorded in Super 2270 (the most resistant cultivar) and Primo early (the most susceptible cultivar), respectively (Table 2). Moreover, significant positive correlation was observed between trichome density and resistance level to TLM larvae ( $r = 0.740$ ).

**Table 2** Density of leaf trichome in tomato cultivars.

Cultivars	Trichome density ± SE (cm <sup>2</sup> )	Level
Primo early	7.94 ± 1.57a*	Low
Early Urbana	23.38 ± 2.34b	Medium
Super 2270	73.11 ± 4.56c	High
F (df = 2, 5)	120.92	
P-value	< 0.0001	

Means followed by the same letters in each column indicate non-significant differences (Duncan's Multiple Range Test,  $P \leq 0.05$ ).

### Discussion

The study revealed that there are significant differences between leaf damage, fruit damage

and total yield caused by TLM in the tested tomato cultivars. The relative resistance of the cultivars according to leaf damage and total yield data were in the decreasing order of: Super 2270, Early Urbana, Rio grande, Petomek, Calj N3 and Primo early.

Our findings are in line with the results of Gharekhani and Salek-Ebrahimi (2014a, b) who demonstrated that different resistance levels can be observed in various tomato cultivars. Resistance levels of ten tomato cultivars to *T. absoluta* were investigated by Sohrabi *et al.* (2016) based on number of mines per leaf, holes on the stem and holes per fruit in field condition. Among the tested cultivars, those with high density of leaf trichome, were more resistant to the pest. The researchers concluded that leaf trichome density is possibly the reason of resistance to TLM larvae. Similarly, Mulusew *et al.* (2013) showed that resistance level of different tomato cultivars was related to leaf trichome density. Navarro *et al.* (2015) reported that oviposition rates of TLM females were different in various tomato cultivars (antixenosis). But there was no antibiosis based resistance among the tested cultivars (Navarro *et al.*, 2015). Life table parameters of TLM on seven tomato cultivars, Primo early, Rio grande, Calj N3, Petomek, Early Urbanam, Super 2270 and Super strain B showed that the longest and the shortest developmental times of immature stages of TLM were on Early Urbana and Calj N3, respectively (Ghaderi *et al.*, 2017). The lowest and the highest values of the intrinsic rate of increase ( $r$ ) and finite rate of increase ( $\lambda$ ) were on Early Urbana Y and Cal JN3, respectively. This laboratory study showed that Calj N3 and Primo early were the most susceptible and most resistant cultivars, respectively (Ghaderi *et al.*, 2017). Difference in susceptibility of the cultivars may be due to different experimental conditions in laboratory and/ or field. Shahbaz *et al.* (2017) demonstrated that performance of TLM larvae fed on tomato cultivars with different resistance level was different. The larvae feeding on resistant cultivars reached the lowest final weight, whereas those reared on the susceptible cultivars reached the highest body

weight. Also, host plant resistance significantly affects the TLM physiology. The highest proteolytic and amylolytic activity were detected in larvae feeding on resistant cultivars, whereas larvae reared on the susceptible cultivar had the lowest enzymatic activity for both third and fourth instar larvae.

In conclusion, there are different resistance levels in tomato cultivars to TLM damage. The cultivars with dense leaf trichome are more resistant to the pest. These cultivars can be considered as candidates for use in integrated management programs of the TLM.

### Acknowledgements

This research was supported by Khuzestan Agricultural Sciences and Natural Resources University [grant number: 9428401].

### References

- Baniameri, V. and Cheraghian, A. 2011. The current status of *Tuta absoluta* in Iran and initial control strategies. EPPO/IOBC/FAO/NEPPO Joint International Symposium on Management of *Tuta absoluta* (tomato borer). Agadir, Morocco. P: 20.
- Braham, M. and Hajji, L. 2012. Management of *Tuta absoluta* (Lepidoptera, Gelechiidae) with insecticides on tomatoes. In: Perveen, F. (Ed.), Insecticides Pest Engineering. Intech Open Acces Publisher, pp. 333-354.
- Dalin, P., Ågren, J., Björkman, C., Huttunen, P. and Kärkkäinen, K. 2008. Leaf trichome formation and plant resistance to herbivory. In: Schaller, A. (Ed.), Induced plant resistance to herbivory. Springer, Dordrecht. pp. 89-105.
- Desneux, N., Wajnberg, E., Wyckhuys, K. A. G., Burgio, G., Arpaia, S., Narvez-Vasquez, C. A., Gonzalez-Cabrera, J., Catalan Ruescas, D., Tabone, E., Frandon, J., Pizzol, J., Poncet, C., Cabello, T., Ubaneja, A. 2010. Biological invasion of European tomato crops by *Tuta absoluta*: Ecology, geographic expansion and prospects for biological control. Journal of Pest Science, 83 (3): 197-215.

- Ghaderi, S., Fathipour Y. and Asgari, S. 2017. Susceptibility of Seven Selected Tomato Cultivars to *Tuta absoluta* (Lepidoptera: Gelechiidae): Implications for Its Management. *Journal of Economic Entomology*, 110 (2): 421-429.
- Gharekhani, G. H. and Salek-Ebrahimi, H. 2014a. Evaluating the damage of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) on some cultivars of tomato under greenhouse condition. *Archives of Phytopathology and Plant Protection*, 47 (4): 429-436
- Gharekhani, G. H. and Salek-Ebrahimi, H. 2014b. Life table parameters of *Tuta absoluta* (Lepidoptera: Gelechiidae) on different varieties of tomato. *Journal of Economic Entomology*, 107 (5): 1765-1770.
- Guedes, R. N. C. and Picanço, M. C. 2012. The tomato borer *Tuta absoluta* in South America: pest status, management and insecticide resistance. *EPPO Bulletin* 42 (2): 211-216.
- Liatti, M., Botto, E. and Alzogaray R. A. 2005. Insecticide resistance in Argentine populations of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Neotropical Entomology*, 34: 113-119.
- Mulusew, B. K. 2013. Resistance mechanism of tomato against *Tuta absoluta*; the role of trichomes in the resistance. M. Sc. thesis of breeding, Wageningen University, Netherland. 41 p.
- Navarro, P., Manzano, S., Jamilena, M., Fernández-Maldonado, F. J., Gallego, J. R. and OEPP/EPPO. 2005. *Tuta absoluta*. European and Mediterranean Plant Protection Organization, Bulletin OEPP/EPPO Bulletin, 35: 434-435. Available online in: [https://www.eppo.int/QUARANTINE/data\\_sheets/insects/DS\\_Tuta\\_absoluta.pdf](https://www.eppo.int/QUARANTINE/data_sheets/insects/DS_Tuta_absoluta.pdf).
- Oliveira, C. M. D., Andrade Júnior, V. C. D., Maluf, W. R., Neiva, I. P., and Maciel, G. M. 2012. Resistance of tomato strains to the moth *Tuta absoluta* imparted by allelochemicals and trichome density. *Ciência e Agrotecnologia*, 36 (1): 45-52.
- Oliveira, F. A., da Silva, D. J. H., Leite, G. L. D., Jham, G. N., and Picanço, M. 2009. Resistance of 57 greenhouse-grown accessions of *Lycopersicon esculentum* and three cultivars to *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Scientia Horticulturae*, 119 (2): 182-187.
- Panda, N., Khush, G. A. 1995. Host plant resistance to insects. CAB International, Wallingford, UK.
- Pedigo, L. P. 2002: Entomology and pest management. Iowa University press.
- Proffitt, M., Birgersson, G., Bengtsson, M., Reis, R., Witzgall, P., and Lima, E. 2011. Attraction and oviposition of *Tuta absoluta* females in response to tomato leaf volatiles. *Journal of Chemical Ecology*, 37 (6), 565-574.
- SAS. 2003. Statistical Analysis System. SAS Release 9.1 for windows, SAS Institute Inc. Cary, NC, USA.
- Shahbaz, M., Noorighanbalani, G. and Naseri, B. 2017. Comparative damage and digestive enzyme activity of *Tutaa bsoluta* (Meyrick) (Lepidoptera: Gelechiidae) on 12 tomato cultivars. *Entomological Research*. doi: 10.1111/1748-5967.12271.
- Sharma, H.C. and Ortiz, R. 2002. Host plant resistance to insects: An eco-friendly approach for pest management and environment conservation. *Journal of Environmental Biology*, 23 (2): 111-135.
- Sohrabi, F., Nooryazdan, H. R., Gharati, B. and Saedi, Z. 2016. Evaluation of ten tomato cultivars for resistance against tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under field infestation conditions. *Entomologia Generalis*, 36 (2): 163-175.
- War, A. R., Paulraj, M. G., Ahmad, T., Buhroo, A. A., Hussain, B., Ignacimuthu, S. and Sharma, H. C. 2012. Mechanisms of plant defense against insect herbivores. *Plant Signaling & Behavior*, 7 (10), 1306-1320.
- Yankova, V. 2012. Damage caused by tomato leaf miner *Tuta absoluta* (Meyrick) in tomato varieties grown in greenhouse. *Plant Science*, (Bulgaria): Available on: <http://agris.fao.org/agris-search/search.do?recordID = BG2012000401>.

## مقاومت ارقام گوجه‌فرنگی به شب‌پره مینوز برگ گوجه‌فرنگی *Tuta absoluta* (Lep: Gelechiidae) در شرایط مزرعه‌ای

فاطمه آزادی<sup>۱</sup>، علی رجب‌پور<sup>۱\*</sup>، امین لطفی جلال‌آبادی<sup>۲</sup> و مهدی محبوب<sup>۳</sup>

- ۱- گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه علوم کشاورزی و منابع طبیعی خوزستان، ملائانی، اهواز.
  - ۲- گروه زراعت و اصلاح نباتات، دانشکده کشاورزی، دانشگاه علوم کشاورزی و منابع طبیعی خوزستان، ملائانی، اهواز.
  - ۳- مرکز تحقیقات کشاورزی و منابع طبیعی استان کرمانشاه، کرمانشاه.
- \* پست الکترونیکی نویسنده مسئول مکاتبه: [rajabpour@ramin.ac.ir](mailto:rajabpour@ramin.ac.ir)  
دریافت: ۱۷ بهمن ۱۳۹۶؛ پذیرش: ۲۶ فروردین ۱۳۹۷

**چکیده:** شب‌پره مینوز برگ گوجه‌فرنگی (*Tuta absoluta*) خطرناک‌ترین آفت گوجه‌فرنگی در ایران است. سطوح مقاومت شش رقم گوجه‌فرنگی به خسارت این آفت طی دو فصل زراعی (۱۳۹۴-۱۳۹۵) مورد ارزیابی قرار گرفت. نمونه‌برداری به صورت هفتگی انجام شد. خسارت‌های برگ‌گی و میوه به همراه عملکرد کل میوه در این ارقام مورد مقایسه قرار گرفت. علاوه بر این، تراکم تریوموم‌های برگ‌گی نیز مشخص شد. ترتیب بیش‌تری به کم‌ترین نرخ آلودگی برگ‌گی ارقام پرموارلی، کالج ان ۳، پتومک، ریوگرند، ارلی اربانا و سوپر 2270 بود. در هر دو فصل زراعی، نرخ آلودگی در ارلی اربانا به صورت معنی‌داری کم‌تر از سایر ارقام بود. ترتیب عملکرد کل گوجه‌فرنگی (از بیش‌ترین به کم‌ترین) به ارقام سوپر ۲۲۷۰، ارلی اربانا، ریوگرند، پتومک، کالج ان ۳ و پرموارلی تعلق داشت.

**واژگان کلیدی:** شب‌پره مینوز برگ گوجه‌فرنگی، مقاومت گیاه میزبان، خسارت برگ‌گی و میوه، عملکرد کل