

Research Article

Effect of pederin on activity of digestive and detoxifying enzymes of *Ephestia kuehniella* (Lep: Pyralidae) and *Tribolium confusum* (Col: Tenebrionidae)

Fereshteh Amirmohammadi¹, Alireza Bandani^{1*}, Bagher Amirheidari² and Ghodratollah Sabahi¹

1. Department of Plant Protection, College of Agriculture and Natural Resources, University of Tehran, Iran.

2. Pharmaceutics Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran.

Abstract: During the last decade, plant and microbial-derived metabolites have received growing attention as potential tools for pest management in agriculture. Pederin (C₂₅H₄₅NO₉) is a vesicant toxin produced by *Pseudomonas*-like bacterial symbionts of rove beetles within the genus *Paederus* (Col: Staphylinidae). In this study, the toxicity of pederin to two stored product pests, *Ephestia kuehniella* Zeller (Lep: Pyralidae) and *Tribolium confusum* Jacquelin du Val (Col: Tenebrionidae) was evaluated using laboratory bioassays. Probit analysis estimated the median lethal concentrations of pederin as 1311.96 and 596.36ppm for *E. kuehniella* fourth larval instar and *T. confusum* adults, respectively. We also measured the activity of two major digestive enzymes (amylases and proteases) as well as three major detoxifying enzymes (P_{450s} monooxygenases, glutathione S-transferases, and carboxyl esterases) in insects treated orally with pederin. Feeding on pederin resulted in significant decrease in the activity of amylolytic, proteolytic, and carboxyl esterase enzymes, but significant increase in the activity of P_{450s} and glutathione S-transferases. Results of this study may highlight pederin as a novel source of pesticides with unique mode of action for use in pest management programs.

Keywords: Digestive Enzyme, Detoxifying Enzyme, Natural Products, *Paederus fuscipes*, pederin, toxin

Introduction

The postharvest losses of stored products caused by insects has been estimated to range from 9% in developed countries to 20% or more in developing countries (Subramanyam, 1995, Phillips and Throne, 2010). For decades, the use of synthetic fumigants has been the major commercial mean of stored product pest control (Fields and White, 2002, Cox *et al.*, 1984, Bell

and Savvidou, 1999). For example, sulfuryl fluoride, methyl bromide and phosphine have long been used for management of stored product pests throughout the world because of their rapid action and activity against a broad spectrum of pests (Fields and White, 2002). However, the harmful effects of these compounds on human health, non-target organisms and the environment as well as low efficiency against the target pest due to cryptic activity and rapid development of resistance have persuaded efforts for development of new classes of pesticides with more specificity against the target pest and less toxic effects to non-target organisms (Gill and Garg, 2014,

Handling Editor: Saeid Moharramipour

*Corresponding author, e-mail: abandani@ut.ac.ir

Received: 22 May 2018, Accepted: 6 December 2018

Published online: 2 February 2019

Desneux *et al.*, 2007, Aktar *et al.*, 2009). Natural products, derived from plants, animals, bacteria, and fungi, are a source of novel compounds with insecticidal, fungicidal, herbicidal, and therapeutic properties besides being environmentally safer molecules than many of the currently used pesticides (Duke *et al.*, 2010, Cantrell *et al.*, 2012). Almost 70% of all newly registered active pesticides have their origins in natural products (Dayan and Duke, 2014), indicating growing interest and exigency for development of pesticides with novel modes of action. Successful examples include the insecticide spinosad, the herbicide glufosinate, and the fungicide strobilurin (Duke *et al.*, 2010).

Polyketides are a class of secondary metabolites produced by certain organisms and serve survival functions for the organisms producing them (Demain and Fang, 2000, Chan *et al.*, 2009). The pederin family consists of numerous members of polyketides with potent cytotoxicity and antitumor and antiviral activity, which are mainly based on inhibition of eukaryotic protein biosynthesis (Piel, 2002, Kellner, 2002). Despite the fact that all other members of this family have been known only from marine sponges, pederin (C₂₅H₄₅NO₉) itself occurs exclusively in several species of rove beetles (Col: Staphylinidae) within the two genera *Paederus* and *Paederidus* (Piel *et al.*, 2004, Piel, 2002). In *Paederus* beetles, pederin is biosynthesized by a symbiotic bacterium in the γ subdivision of the Proteobacteria, which is closely related to the ubiquitous species, *Pseudomonas aeruginosa* (Piel *et al.*, 2004, Piel, 2002). The bacteria, and therefore the ability of pederin biosynthesis, are largely confined to adult females. However, larvae and adult males can also store pederin acquired maternally or by ingestion (Kellner, 1998). Although, the toxic properties of pederin to humans have been recognized for nearly one century (Pirajá Da Silva, 1912), its insecticidal activity has been very poorly studied (Tabadkani and Nozari, 2014, Kellner and Dettner, 1996). Kellner and Dettner (1996) showed that *Paederus* larvae are deterrent to

wolf spiders (Kellner and Dettner, 1996), but not to a range of predatory insects. In another study, Tabadkani and Nozari (2014) showed that the body fluid of *Paederus fuscipes* Curtis is highly toxic to the predatory ground beetle *Chlaenius pallipes* Gebler (Col: Carabidae).

Given the widespread use of natural products in crop protection and medicine, pederin may provide a source of new classes of insecticides with specific mode of action for pest management. In this study, the toxicity of pederin to two stored product pests, the Mediterranean flour moth, *Ephestia kuehniella* (Zeller) (Lep: Pyralidae) and the confused flour beetle, *Tribolium confusum* Jacquelin du Val (Col: Tenebrionidae) was studied under laboratory condition. Both species are common and important pests of stored cereals worldwide. In addition to direct damage caused through external feeding of the larvae, the quality of the stored product is severely affected by the presence of living and dead insects, frass, feces, silk, larval exuviae, and chemical excretions and odors (Stejskal and Hubert, 2006, Panizzi and Parra, 2012). We also investigated the activity of common digestive and detoxifying enzymes in response to infection with pederin.

Materials and Methods

Insects

Adult *P. fuscipes* were collected (between 11-15 o'clock) from rice fields at Rasht (37.2 N 49.60E) (Guilan province, north of Iran) using hand, sweep net, and light traps. After species identification, adult females were used for pederin extraction. Adult females are readily identified by absence of a distinct cleft at posterior edge of their last abdominal ventrite, as opposed to males.

E. kuehniella eggs were obtained from a commercial insectariums. A stock colony was established in the laboratory by distributing 0.5g eggs in plastic containers (10 × 7 × 5cm) containing 1000g wheat flour plus 100g yeast. The colony was maintained at controlled condition (25 ± 1 °C, 65 ± 5% RH, and 16L: 8D

photoperiod). The fourth instar larvae were used for all experiments.

Adult *T. confusum* were hand collected from an infected wheat warehouse and transferred to laboratory for establishment of stock colony. After species identification, by existing keys the beetles were released inside plastic containers (5.5 × 15.5 × 25cm) containing 500g wheat flour plus 15g yeast. The containers were maintained at growth chambers (28 ± 1 °C, 70 ± 5% RH, and absolute darkness). Newly emerged Adult beetles (< 24-h old) were used for all experiments.

Pederin extraction

Pederin extraction was performed following the method of Mohseni *et al.* (2014) (Mohseni *et al.*, 2014). Briefly, *P. fuscipes* adult females were finely crushed in Eppendorf micro-tubes containing absolute ethanol (Merck). The mixture was then centrifuged (Heraeus Fresco 21 centrifuge, Fisher Scientific, Germany) at 3,000 ×g for 10 min to remove tissue particles and the resulting supernatant was maintained at 37 °C for ethanol evaporation. After removal of lipids using hexane-methanol (1:1 v/v), the methanol phase was separated using decanter and evaporated by maintaining at 37 °C. The raw extract was dissolved in ethyl acetate (98%, Merck) and stored at 4 °C for further analysis. 10µl of the ethyl acetate extract and authentic pederin were applied to high-performance thin-layer chromatography (TLC) plates (0.2mm silica gel 60, 10 × 10cm, Merck), which were developed in ethyl acetate. After evaporation of the solvent, they were stained in a dipping solution of anisaldehyde: sulfuric acid: acetic acid (1: 2: 100v/v/v) (99%, Merck) and heated at 90 °C for 2min. Using this procedure, lilac spots of pederin (> 0.05µg) are provided at $R_f = 0.22$ (Kellner and Dettner, 1995).

Concentration-mortality response

The concentration-mortality response of the fourth larval instar of *E. kuehniella* and adult stage of *T. confusum* to the pederin extract was

studied to estimate the median lethal concentration (LC50) for each species. Based on concentration-setting pre-tests, the appropriate concentrations of pederin extract, causing the mortality range of 10-90%, were determined as 300, 448, 670, 1003, and 1500ppm for *T. confusum* and as 700, 962, 1322, 1818, and 2500 ppm for *E. kuehniella*. One mL of each concentration was mixed with 5g wheat flour and the resulting toxic diet was distributed at the floor of a Petri-dish (9cm diameter). Twenty *E. kuehniella* fourth larval instar or *T. confusum* adults, which had been starved for 24 hours, were released inside each Petri-dish. The Petri-dishes were maintained at controlled condition and mortality was recorded 24 hours post treatment. Both bioassays were conducted in a completely randomized design (CRD) with three replications. The percentage of mortality was corrected using Abbott's formula as follow:

$$\text{Corrected mortality} = \left(\frac{T - C}{100 - C} \right) \times 100$$

Where T and C are the numbers of dead insects in treatment and control, respectively (Abbott, 1925).

Enzyme assays

The effect of feeding on *P. fuscipes* body fluid on activity of digestive enzymes (amylases and proteases) and detoxifying enzymes (glutathione S-transferase (GST), cytochrome P₄₅₀, and carboxyl esterase) was studied in last larval instar of *E. kuehniella* and adult *T. confusum*. One mL of the LC₃₀ of pederin extract was mixed with 5g wheat flour and the resulting toxic diet was distributed at the base of a Petri-dish (9cm diameter). The last larval instar of *E. kuehniella* and adult *T. confusum* were then released inside Petri-dishes and allowed to feed on the toxic diet for 24 hours. The same volume of diet, without pederin contamination, was provided for these species as control. Treated insects were used for enzyme assays after 24 hours of feeding on toxic diet and control insects on ethyl acetate diet were investigated too.

Protein quantification

Total protein concentration of samples was determined according to Bradford method (Bradford, 1976) at 630nm, with bovine serum albumin as standard.

Digestive enzyme assays

Sample preparation

After feeding on toxic diet for 24 hours, 10 fourth larval instars of *E. kuehniella* were dissected under a stereomicroscope in saline solution (NaCl, 10mM) to extract guts. In case of adult *T. confusum*, the whole body (n = 20) was used for enzyme sample preparation. The samples were homogenized in pre-cooled homogenizer in distilled water (w/v). The homogenates were then centrifuged at $16,000 \times g$ for 10min at 4 °C and the resultant supernatants were stored at -20 °C for subsequent analysis of digestive enzyme activities.

Amylase activity

The total amylase activity of last larval instar of *E. kuehniella* and adult *T. confusum* was measured using a colorimetric method with 3,5-dinitrosalicylic acid (DNS) reagent and 1% (w/v) soluble starch as substrate according to the method of Bernfeld (1955) with slight modifications. The reaction mixture, containing 70μl of Tris-HCl buffer (20 mM, pH 7), 10μl of the enzyme solution, and 20μl of 1% starch, was incubated for 10 min at 35 °C. The reaction was stopped by the addition of 50μl DNS reagent and heating in boiling water for 10 min. The absorbance of the resultant complex was read at 540nm using a microplate reader (ELx808). One unit of α-amylase activity was defined as the amount of enzyme required to produce 1 mg maltose in 30 min at 37 °C. The negative control contained all reaction mixtures except enzyme source to prove the enzyme presence in the samples. This assay was carried out in triplicate.

Protease activity

The total proteolytic activity was quantified using azocasein as the substrate according to Elpidina *et al.*, (2001) with slight modifications.

The reaction mixture, consisting of 40μl of Tris-HCl buffer (20mM, pH 7), 5μl of 2% (w/v) azocasein (pH 5.0), and 10μl of the enzyme solution, was incubated for 60 min at 37 °C. The reaction was stopped by adding 100μl 30% TCA (trichloroacetic acid). The samples were incubated at 4 °C for 30min to precipitate non-hydrolyzed substrate, then centrifuged at $15,000 \times g$ for 15 min at 4 °C. One hundred μl of the resultant supernatant was mixed with equal volume of KOH 2M and the absorbance was read at 405nm. One unit of enzyme activity (U) was defined as a change in absorbance at 405nm of one mg protein after one minute. This assay was carried out in triplicate.

Detoxifying enzyme assays

Sample preparation

Bodies of 10 last larval instar of *E. kuehniella* and 20 adult *T. confusum* were homogenized in 400μl phosphate buffer (0.04 M, pH 7.0) containing Triton X-100 (0.1% v/v) and centrifuged at $10,000 \times g$ for 20 min at 4 °C. The supernatant was transferred to new tubes and preserved at -20 °C for analysis of the activity of carboxyl esterase, glutathione transferase and P450 and all steps were performed for controls too.

Glutathione S-transferase

Glutathione S-transferase (GST) activity was measured using 2,4-Dinitrochlorobenzene (CDNB) according to the method of Habig *et al.* (1974). The total reaction volume per well of a 96-well microplate was 200μl, consisting of 10μl of enzyme solution, 2μl of 200mM CDNB (containing 0.1% (v/v) ethanol), and 188μl of 10mM GSH in Tris-HCl buffer (0.05 M; pH 8). GST activity was determined by the change in absorbance as measured every 30 s for 5 min at 340nm. Control wells consisted of 2μl of CDNB, 188μl of GSH, and 10μl of Tris-HCl buffer (0.05 M; pH 8) (Habig *et al.*, 1974) and all steps were performed for controls too.

Cytochrome P₄₅₀

Cytochrome P₄₅₀ activity was measured in terms of general oxidase level using heme

peroxidation method with 3,3',5,5'-tetramethylbenzidine (TMBZ) (Sigma Aldrich) as the substrate (Brogdon *et al.*, 1997) (William and Janet, 1997). The total reaction volume per well of a 96-well microplate was 325µl, including 20µl of enzyme solution, 80µl of 0.625 M potassium phosphate buffer (pH 7.2), 200µl of TMBZ solution, and 25µl of hydrogen peroxide (H₂O₂) (3%). Plates were incubated at room temperature for 2h, after which the absorbance was read at 450nm. Control wells consisted of 20µl of distilled water, 80µl of 0.625M potassium phosphate buffer, 200µl of TMBZ solution, and 25µl of H₂O₂ (3%). A standard curve for heme peroxidase activity was prepared using different concentrations of cytochrome C (0.15, 0.3, 0.6, 0.8, 1, and 1.2) from horse heart (Sigma Aldrich). The activity of cytochrome P₄₅₀, obtained from plate reading, was expressed as equivalent units (EU) of cytochrome P₄₅₀ per milligram of protein using the standard curve of cytochrome C (William and Janet, 1997) and all steps were performed for controls too.

Carboxyl esterase

The general esterase activity of *E. kuehniella* and *T. confusum* were quantified using α-naphthyl-acetate and β-naphthyl acetate as substrates according to the method of Van Asperen with slight modifications (Van Asperen, 1962). The total reaction volume per well of a 96-well microplate was 270µl, consisting of 20µl of enzyme solution, 90µl of substrate, 70µl of phosphate buffer (0.2 M, pH 7.0), and 90µl of fast blue RR salt. Plates were incubated at 37 °C for 30min, after which, the absorbance was read at 450nm for α-naphthyl-acetate and at 540nm for β-naphthyl acetate every 2min for 10min (Van Asperen, 1962) 37.2 N 49.60 E and all steps were performed for controls too.

Data analysis

All experiments were carried out in completely randomized design (CRD) with

two treatments (control and treated insects) and three replications. The corrected mortality data of *E. kuehniella* larvae and *T. confusum* adults were used to estimate the median lethal concentration (LC₅₀) using Probit analysis in POLO-PC 2002 software (California, USA). The average enzyme activity was compared between control and treated larvae using independent *t*-tests of SPSS software (version 19).

Results and Discussion

The TLC yielded lilac spots of pederin at $R_f = 0.22$, indicating that pederin was successfully extracted using our procedure (Fig. 1). Our results revealed high toxicity of *P. fuscipes* body extract for both *E. kuehniella* and *T. confusum* via ingestion. The median lethal concentration (LC₅₀) of pederin for last larval instar of *E. kuehniella* and adult *T. confusum* was estimated as 1311.96 and 596.36ppm, respectively (Table 1).




Figure 1 Part of TLC plate showing successful extraction of pederin from *Paederus fuscipes*. Extracts of adult females were applied together with known amounts of authentic pederin to the starting line (bottom), developed in ethyl acetate and stained with anisaldehyde/sulfuric acid/acetic acid.

Table 1 Results of probit analysis (mean \pm 95% CI) for estimation of median lethal concentration (LC₅₀) of pederin against last larval instar of *Ephestia kuehniella* and adult *Tribolium confusum*.

Insect	n	Slope \pm SE	LC ₅₀ (95% lower-upper) (ppm)	χ^2	df	Heterogeneity
<i>E. kuehniella</i>	300	4.01 \pm 0.46	1311.96 (1187.17-1444.43)	8.11	13	0.624
<i>T. confusum</i>	300	3.17 \pm 0.38	596.36 (514.21-679.79)	6.69	13	0.515

In agreement with these results, Tabadkani and Nozari (2014) showed that *P. fuscipes* body fluid is highly toxic to the predatory ground beetle, *Ch. pallipes*. When the beetles fed a diet containing *P. fuscipes* body fluid, they became torpid within 24-36 hours and started to die after 48-60 hours of oral treatment (Tabadkani and Nozari, 2014). Pederin is expected to act against a wide range of eukaryotic organisms with respect to its unique mode of action. Generally, pederin inhibits mitosis in eukaryotic cells by blocking the synthesis of particular proteins in ribosomes (Pavan, 1963). Therefore, any type of living tissue is expected to be affected by this compound.

We measured the activity of two common digestive enzymes (amylases and proteases) in *E. kuehniella* last larval instar and *T. confusum* adults in response to oral exposure to *P. fuscipes* body extract (Fig. 2). Feeding on pederin resulted in significant decrease in the activity of amylases in both species in comparison with control (Fig. 2a). Similarly, the activity of proteases significantly decreased in both *E. kuehniella* and *T. confusum* as a result of feeding on pederin extract (Fig. 2b). These results imply that pederin strongly interferes with normal function of digestive enzymes in the gut.

Despite the abundance of studies on insecticidal activity and enzyme inhibitory effects of a wide variety of plant-derived toxins/inhibitors (Carlini and Grossi-De-Sá, 2002), the effect of bacterial-derived toxins on digestive enzyme of insects have been rarely studied. However, in several studies, the oral insecticidal effects of bacterial toxins with different modes of actions have been demonstrated (Khandelwal *et al.*, 2004, Hinchliffe *et al.*, 2010, Bowen and Ensign, 1998, Andrejko *et al.*, 2013). For example, a pilin subunit of the nematode symbiotic bacterium, *Xenorhabdus nematophilus*, has been reported to have strong oral toxicity to the larvae of *Helicoverpa armigera* (Hübner) (Lep:

Noctuidae), causing extensive damage to the midgut epithelial membrane (Khandelwal *et al.*, 2004). Transformation of a 537 bp *mrxA* gene, encoding the pilin subunit, into the genome of tobacco and tomato, was reported to cause 100% mortality in larvae of *H. armigera* and reduce damage on genetically manipulated plants (Kumari *et al.*, 2015). Lv *et al.* (2014) reported significant decrease in the amylase activity (Lv *et al.*, 2014), beside increase in protease activity of the 5th larval instar of the oriental armyworm, *Mythimna separata* (Walker) (Lep: Noctuidae) when reared on fraxinellone, a toxin extracted from *Dictamnus dasycarpus*.

Insects have developed a variety of metabolic responses in order to deal with pesticides and other xenobiotic compounds. In most species, these responses include two distinct phases I and II. In phase I, the foreign compound(s) is oxidized to create a reactive site (electrophilic site) by the action of detoxification enzymes such as P_{450s} and esterases, while phase II involves the conjugation of phase I products with endogenous substances such as glutathione. This process transforms the lipophilic xenobiotic compounds into more water soluble derivatives and thus, facilitates excretion of the xenobiotic compound (Mamidala *et al.*, 2011; Reddy *et al.*, 2012). We measured the activity of three major enzymes (monooxygenases, GSTs, and esterases) in *E. kuehniella* larvae and *T. confusum* adults in response to oral exposure to pederin extract (Fig. 3). As expected, feeding on pederin caused significant increase in the activity of P_{450s} monooxygenase of both *E. kuehniella* larvae and *T. confusum* adults when compared with control (*t*-test: df = 4, *t* = 22.99 and 15.58 for *T. confusum* and *E. kuehniella*, respectively, *P* < 0.01) (Fig. 3a). Similarly, the activity of glutathione S-transferases was significantly increased following oral treatment of *T. confusum* with pederin extract (*t*-test: *t* = 7.83, df = 4, *P* < 0.01) and tended to increase in *E. kuehniella* larvae (*t*-test: *t* = 2.44, df

= 4, $P = 0.071$) (Fig. 3b). By contrast, a significant decrease in the activity of carboxyl esterase was detected in both species, when either α -naphthyl-acetate or β -naphthyl acetate was used as substrate for the enzyme (Fig. 3c and d). These results may imply that oxygenases and glutathione S-transferases are the major groups of detoxifying

enzymes involved in metabolism of pederin in treated insects. Feeding on fraxinellone has been shown to result in increased activity of carboxylesterase and glutathione S-transferase, but decreased activity of other detoxifying enzymes such as NADPH-P450 reductase and O-demethylase (Lv et al., 2014).




Figure 2 (a): Relative activity of amylolytic and (b): proteolytic enzymes of *Ephestia kuehniella* last larval instar and *Tribolium confosum* adults in response to oral treatment with pederin, ** show significant difference treatment and control at $P < 0.01$ (t-student test).




Figure 3 Activity of (a): P_{450s} monooxygenase, (b): glutathione S-transferase, and (c): carboxyl esterase and for (d): alpha and beta naphthyl acetate as substrate enzymes of *Ephestia kuehniella* last larval instar and *Tribolium confosum* adults in response to oral treatment with pederin, ** show significant difference between treatment and control at $P < 0.01$ (t-student test).

Altogether, results of the current study indicate high oral toxicity of pederin for two distantly related insects (a beetle and a moth). However, the mechanism(s) and the precise mode of action through which pederin acts against insect digestive tract remains to be discovered in future studies. A possible explanation may be disruption of midgut peritrophic membrane as a barrier that protects the epithelium from mechanical damage by food, exposure to ingested toxins as well as microbial invasion besides concentrating food and digestive enzymes (Shao *et al.*, 2001). Plant and microbial toxins have recently attracted increasing attention as tools for control of insect pests. Current methods of administration are either expression of the toxin in genetically modified plants or dispersing the toxin over plant surfaces (Hinchliffe *et al.*, 2010). Pederin, as a novel compound with unique mode of action, may provide new insights for control of insect pests in agriculture and homes. This persuades researches for development of appropriate formulations in order to deliver sufficient amounts of the toxin to the target pest, and at the same time, avoid the harmful effects of this compound on non-target organisms including natural enemies as well as the end users by whom the crop is intended to be consumed (Hinchliffe *et al.*, 2010). Given the high oral toxicity of pederin to insects and available routes for synthesis (Wan *et al.*, 2011, Mosey and Floreancig, 2012), forthcoming works may evaluate the efficiency of pederin as baited traps for control of specific pests, such as cockroaches, termites, ants, in homes.

Acknowledgements

The authors greatly appreciate entomology laboratory at the University of Tehran and Biotechnology laboratory at Kerman medical University for their generous help in this work.

References

- Abbott, W. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18 (2): 265-267.
- Aktar, W., Sengupta, D. and Chowdhury, A. 2009. Impact of pesticides use in agriculture: their benefits and hazards. *Interdisciplinary Toxicology*, 2 (1): 1-12.
- Andrejko, M., Zdybicka-Barabas, A., Janczarek, M. and Cytryńska, M. 2013. Three *Pseudomonas aeruginosa* strains with different protease profiles. *Acta Biochimica Polonica*, 60 (1).
- Bell, C. and Savvidou, N. 1999. The toxicity of Vikanel (sulfuryl fluoride) to age groups of eggs of the Mediterranean flour moth (*Ephestia kuehniella*). *Journal of Stored Products Research*, 35 (3): 233-247.
- Bernfeld, P. 1955. Amylase α and β . *Methods in Enzymology*, 1: 149-158.
- Bowen, D. J. and Ensign, J. C. 1998. Purification and characterization of a high-molecular-weight insecticidal protein complex produced by the entomopathogenic bacterium *photorhabdus luminescens*. *Applied and Environmental Microbiology*, 64 (8): 3029-3035.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72 (1-2): 248-254.
- Cantrell, C. L., Dayan, F. E. and Duke, S. O. 2012. Natural products as sources for new pesticides. *Journal of Natural Products*, 75 (6): 1231-1242.
- Carlini, C. R. and Grossi-de-Sá, M. F. 2002. Plant toxic proteins with insecticidal properties. A review on their potentialities as bioinsecticides. *Toxicon*, 40 (11): 1515-1539.
- Chan, Y. A., Podevels, A. M., Kevany, B. M. and Thomas, M. G. 2009. Biosynthesis of polyketide synthase extender units. *Natural Product Reports*, 26 (1): 90-114.
- Cox, P., Bell, C., Pearson, J. and Beirne, M. 1984. The effect of diapause on the tolerance of larvae of *Ephestia kuehniella* to methyl bromide and phosphine. *Journal of Stored Products Research*, 20 (4): 215-219.
- Dayan, F. E. and Duke, S. O. 2014. Natural compounds as next-generation herbicides. *Plant Physiology*, 166 (3): 1090-1105.

- Demain, A. L. and Fang, A. 2000. The natural functions of secondary metabolites. History of Modern Biotechnology I. Springer.
- Desneux, N., Decourtye, A. and Delpuech, J. M. 2007. The sublethal effects of pesticides on beneficial arthropods. Annual Review of Entomology, 52: 81-106.
- Duke, S. O., Cantrell, C. L., Meepagala, K. M., Wedge, D. E., Tabanca, N. and Schrader, K. K. 2010. Natural toxins for use in pest management. Toxins, 2 (8): 1943-1962.
- Elpidina, E. N., Vinokurov, K. S., Gromenko, V. A., Rudenskaya, Y. A., Dunaevsky, Y. E. and Zhuzhikov, D. P. 2001. Compartmentalization of proteinases and amylases in *Nauphoeta cinerea* midgut. Archives of Insect Biochemistry and Physiology, 48 (4): 206-216.
- Fields, P. G. and White, N. D. 2002. Alternatives to methyl bromide treatments for stored-product and quarantine insects. Annual Review of Entomology, 47 (1): 331-359.
- Gill, H. K. and Garg, H. 2014. Pesticides: Environmental Impacts and Management Strategies. Pesticides-Toxic Aspects. InTech.
- Habig, W. H., Pabst, M. J. and Jakoby, W. B. 1974. Glutathione S-transferases the first enzymatic step in mercapturic acid formation. Journal of Biological Chemistry, 249 (22): 7130-7139.
- Hinchliffe, S. J., Hares, M. C. and Dowling, A. J. 2010. Insecticidal toxins from the *Photorhabdus* and *Xenorhabdus* bacteria. The Open Toxinology Journal, 3 (1).
- Kellner, R. L. 1998. When do *Paederus riparius* rove beetles (Coleoptera: Staphylinidae) biosynthesize their unique hemolymph toxin pederin? Zeitschrift für Naturforschung C, 53 (11-12): 1081-1086.
- Kellner, R. L. 2002. Molecular identification of an endosymbiotic bacterium associated with pederin biosynthesis in *Paederus sabaeus* (Coleoptera: Staphylinidae). Insect Biochemistry and Molecular Biology, 32 (4): 389-395.
- Kellner, R. L. and Dettner, K. 1995. Allocation of pederin during lifetime of *Paederus* rove beetles (Coleoptera: Staphylinidae): Evidence for polymorphism of hemolymph toxin. Journal of Chemical Ecology, 21 (11): 1719-1733.
- Kellner, R. L. and Dettner, K. 1996. Differential efficacy of toxic pederin in deterring potential arthropod predators of *Paederus* (Coleoptera: Staphylinidae) offspring. Oecologia, 107 (3): 293-300.
- Khandelwal, P., Choudhury, D., Birah, A., Reddy, M., Gupta, G. P. and Banerjee, N. 2004. Insecticidal pilin subunit from the insect pathogen *Xenorhabdus nematophila*. Journal of Bacteriology, 186 (19): 6465-6476.
- Kumari, P., Mahapatro, G. K., Banerjee, N. and Sarin, N. B. 2015. A novel pilin subunit from *Xenorhabdus nematophila*, an insect pathogen, confers pest resistance in tobacco and tomato. Plant Cell Reports, 34 (11): 1863-1872.
- Lv, M., Wu, W. and Liu, H. 2014. Effects of fraxinellone on the midgut enzyme activities of the 5th instar larvae of oriental armyworm, *mythimna separata* walker. Toxins, 6 (9): 2708-2718.
- Mohseni, M., Haromar, S., Chaichi, M.-J. and Alijanpour-Toloti, S.-O. 2014. Dermatitis activity of pederin and isolation of endosymbiont bacterium associated with its biosynthesis from *Paederus fuscipes* beetle of Mazandaran, Iran. Journal of Mazandaran University of Medical Sciences, 23 (2): 2-13.
- Mosey, R. A. and Floreancig, P. E. 2012. Isolation, biological activity, synthesis, and medicinal chemistry of the pederin/mycalamide family of natural products. Natural product reports, 29 (9): 980-995.
- Panizzi, A. R. and Parra, J. R. 2012. Insect Bioecology and Nutrition for Integrated Pest Management, CRC press.
- Pavan, M. 1963. Ricerche biologiche e mediche su pederina e su estratti purificati di *Paederus fuscipes* Curt. (Coleoptera staphylinidae), Ponzio.
- Phillips, T. W. and Throne, J. E. 2010. Biorational approaches to managing stored-product insects. Annual Review of Entomology, 55: 375-397.

- Piel, J. 2002. A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of *Paederus* beetles. *Proceedings of the National Academy of Sciences*, 99 (22): 14002-14007.
- Piel, J., Höfer, I. and Hui, D. 2004. Evidence for a symbiosis island involved in horizontal acquisition of pederin biosynthetic capabilities by the bacterial symbiont of *Paederus fuscipes* beetles. *Journal of Bacteriology*, 186 (5): 1280-1286.
- Pirajá da Silva, M. 1912. Le *Paederus columbinus* est vésicant. *Arch Parasitol*, 15: 429-431.
- Shao, L., Devenport, M. and Jacobs-Lorena, M. 2001. The peritrophic matrix of hematophagous insects. *Archives of Insect Biochemistry and Physiology*, 47 (2): 119-125.
- Stejskal, V. and Hubert, J. 2006. Arthropods as sources of contaminants of stored products: an overview. 9th International Working Conference on Stored Product Protection.
- Subramanyam, B. 1995. *Integrated Management of Insects in Stored Products*, CRC Press.
- Tabadkani, S. M. and Nozari, J. 2014. Relaxed predation hinders development of anti-predator behaviors in an aposematic beetle. *Entomologia Experimentalis et Applicata*, 153 (3): 199-206.
- Van Asperen, K. 1962. A study of housefly esterases by means of a sensitive colorimetric method. *Journal of Insect Physiology*, 8 (4): 401-416.
- Wan, S., Wu, F., Rech, J. C., Green, M. E., Balachandran, R., Horne, W. S., Day, B. W. and Floreancig, P. E. 2011. Total synthesis and biological evaluation of pederin, psymberin, and highly potent analogs. *Journal of the American Chemical Society*, 133 (41): 16668-16679.
- William, G. B. and Janet, C. 1997. Heme peroxidase activity measured in single mosquitoes identifies individuals expressing an elevated oxidase for insecticide resistance. *Journal of the American Mosquito Control Association*, 13 (3): 233-237.

تأثیر پدیرین بر فعالیت آنزیم‌های گوارشی و سم‌زدا در لاروهای *Ephestia kuehniella* و *Tribolium confusum* (Coleoptera: Tenebrionidae) (Lepidoptera: Pyralidae)

فرشته امیرمحمدی^۱، علیرضا بندانی^{۱*}، باقر امیرحیدری^۲ و قدرت‌الله صباحی^۱

۱- گروه گیاه‌پزشکی، پردیس کشاورزی و منابع طبیعی، دانشگاه تهران، تهران، ایران.

۲- مرکز تحقیقات فارماسیوتیکس، پژوهشکده نوروفارماکولوژی، دانشگاه علوم پزشکی کرمان، کرمان، ایران.

پست الکترونیکی نویسنده مسئول مکاتبه: abandani@ut.ac.ir

دریافت: ۱ خرداد ۱۳۹۷؛ پذیرش: ۱۵ آذر ۱۳۹۷

چکیده: در دهه گذشته متابولیت‌های مشتق شده از میکروب و گیاه توجه روبه رشدی را به‌عنوان یک ابزار قدرتمند برای کنترل آفت در کشاورزی دریافت کرده‌است. پدیرین ($C_{25}H_{45}NO_9$) یک ماده تاول‌زاست که از باکتری شبه سودوموناس هم‌زیست با سوسک دونده جنس *Paederus* (Col: Staphylinidae) تولید می‌شود. در این پژوهش سمیت پدیرین بر دو آفت انباری *Ephestia kuehniella* (Lep: Pyralidae) و *Tribolium confusum* (Col: Tenebrionidae) با استفاده از زیست‌سنجی‌های آزمایشگاهی ارزیابی شده‌است. آنالیز پروبیت میانگین غلظت‌های کشنده پدیرین ۱۳۱۱/۹۶ و ۵۹۶/۳۶ (قسمت در میلیون) به‌ترتیب برای لارو سن چهار *E. kuehniella* و بالغ *T. confusum* تخمین زد. ما هم‌چنین فعالیت دو آنزیم گوارشی اصلی (آمیلاز و پروتئاز) و سه آنزیم سم‌زدای اصلی (مونواکسیژنازهای P_{450} ، گلوکونایون‌اس‌ترانسفرازها و کربوکسیل‌استرازها) را در حشراتی که به‌صورت خوراکی با پدیرین تیمار شدند اندازه‌گیری کردیم. تغذیه با پدیرین منجر به کاهش معنی‌دار در فعالیت آنزیم‌های آمیلولیتیک، پروتئولیتیک و کربوکسیل‌استراز شد، اما افزایش معنی‌دار در فعالیت P_{450} ها و گلوکونایون‌اس‌ترانسفرازها را باعث می‌شود. نتایج این پژوهش ممکن است پدیرین را به‌عنوان یک منبع جدید از آفت‌کش‌ها با نحوه تأثیر منحصر به فرد در برنامه‌های مدیریت آفت برجسته کند.

واژگان کلیدی: آنزیم گوارشی، آنزیم سم‌زدا، محصولات طبیعی، *Paederus fuscipes*، پدیرین، سم