

Research Article

Dung flies fauna (Diptera: Scathophagidae) in East Azerbaijan province of Iran, with eight new records

Samad Khaghaninia* and Yaser Gharajedaghi

University of Tabriz, Department of Plant Protection, Faculty of Agriculture, P. O. Box: 51664, Tabriz, I. R. Iran.

Abstract: Based on specimens collected from East Azerbaijan province during 2009-2012, five genera and nine species of the family Scathophagidae are recognized. Eight species including: *Cordilura rufipes* (Meigen, 1826), *Gimnomera montana* Ozerov et Krivosheina, 2013, *Norellisoma spinimana* (Fallén, 1819), *Parallelomma albipes* (Fallén, 1819), *Scathophaga kaszabi* (Šifner, 1975), *Scathophaga lutaria* (Fabricius, 1794), *Scathophaga stercoraria* (Linnaeus, 1758), *Scathophaga taeniopa* (Rondani, 1867) and four genera (*Cordilura* Fallén, 1810, *Gimnomera* Rondani, 1866, *Norellisoma* Wahlgren, 1917, *Parallelomma* Becker, 1894)) are recorded from Iran for the first time. Photos and key to species of the family Scathophagidae occurring in this region are provided.

Keywords: Dung flies, Scathophagidae, new records, East Azerbaijan province, Iran

Introduction

The Scathophagidae (dung flies) is a small family of the superfamily Muscoidea (Diptera; Calyptrata) with about 419 species which belong to two subfamilies (Scathophaginae and Deliniinae) (Ozerov, 2009; Pape et al., 2011; Šifner, 2012). The family Scathophagidae includes more than 57 genera and 250 species, mainly in the Palaearctic region (Šifner, 2008, 2009; Ozerov and Freidberg, 2010). The most species of Scathophagidae are distinguished from the closest Muscoid flies by combinations of the following characters: Male and female eyes dichoptic, palpi present, occiput usually with numerous fine pale setulae, frontal vitta broad and without

setae, lower calipter linear, meron without hairs and setae, ventral surface of scutellum bare, katepisternum usually with one bristle, A_1 usually reaching wing margin, A_2 not curved forward beyond tip of A_1 (McAlpine, 1981; Ozerov, 2009). Biologically, adults are predators, most larvae are phytophagous, some larvae are scavenger or carnivorous (Gorodkov, 1986; Ozerov and Freidberg, 2010). The first contribution to the fauna of Scathophagidae of Iran was made by F. Šifner (1981) who described 2 species belonging to 2 genera (*Paramicroprosopa hoberlandti* (Šifner, 1981) and *Scathophaga jezeki* (Šifner, 1981)). Nevertheless, the long-legged flies of Iran are poorly known. Present study adds four genera and eight species of dung flies to the fauna of Iran.

Materials and Methods

Adult specimens were collected by standard sweep-netting in forest and grassland habitats

Handling Editor: Dr. Ali Asghar Talebi

*Corresponding author, e-mail: skhaghaninia@tabrizu.ac.ir
Received: 25 January 2014, Accepted: 25 February 2014
Published online: 25 February 2014

from East Azerbaijan province of Iran during 2009-2012. The samples were killed in a killing jar containing potassium cyanide. Male genitalia were macerated in 10% KOH. The material examined is deposited in collections of the following institutions:

IMTU: Insect Museum of Tabriz University.

ZMUM: Zoological Museum, Moscow University.

Results

In this study, five genera and nine species of the family Scathophagidae are recognized. Eight species including *Cordilura rufipes* (Meigen, 1826), *Gimnomera montana* Ozerov et Krivosheina, 2013, *Norellisoma spinimana* (Fallén, 1819), *Parallelomma albipes* (Fallén, 1819), *Scathophaga kaszabi* (Šifner, 1975), *Scathophaga lutaria* (Fabricius, 1794), *Scathophaga stercoraria* (Linnaeus, 1758), *Scathophaga taeniopa* (Rondani, 1867) are recorded from Iran for the first time. Of which, four genera (*Cordilura* Fallén, 1810, *Gimnomera* Rondani, 1866, *Norellisoma* Wahlgren, 1917, *Parallelomma* Becker, 1894) are new records to Iran. In addition, all species are recorded for the first time from the province. Species are listed in alphabetic order.

Key to the species of the family Scathophagidae in East Azerbaijan

1. Abdomen with long and dense yellow hairs. 2
 - Abdomen without dense hairs.....4
2. Pteropleoron with hairs*Scathophaga stercoraria* (Linnaeus)
 - Pteropleoron bare3
3. Body yellowish, hind femur with more than 3 long dorsal setae..... *S. lutaria* (Fabricius)
 - Body greenish, hind femur of male with 2 to 3 long dorsal setae..... *S. taeniopa* (Rondani)
4. Body yellow5
 - Body dark green or black6
5. One pair of scutellar setae*Norellisoma spinimana* (Fallén)
 - Two pairs of scutellar setae.....*Gimnomera montana* Ozerov et Krivosheina
6. Body greenish7

- Body black8
- 7. Occiput with some black setae, 4th abdominal sternite long, surstylus wide (Fig. 10, 12)*S. jezeki* (Šifner)
 - Occiput with yellow setae, 4th abdominal sternite broad, surstylus thin (Fig. 11, 13).....*S. kaszabi* (Šifner)
- 8. Femora black*Cordilura rufipes* (Meigen)
 - Femora yellow.....*Parallelomma albipes* (Fallén)

Cordilura rufipes (Meigen, 1826) (Fig. 1)

Material examined: East Azerbaijan: Kandovan, 37°46' N, 46°16' E, 2500 m a.s.l., 15 June 2010 (moist grassland habitat), 1 ♀; Maragheh, 37°25' N, 46°25' E, 1790 m a.s.l., 5 June 2011 (dry grassland habitat), 1 ♀; (2 ♀, IMTU; 1 ♀ ZMUM).

Distribution: Austria, Belgium, China, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Ireland, Lithuania, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Siberia, Slovakia, Spain, Sweden (Šifner, 2008). New to the Iranian insect fauna.

Gimnomera montana Ozerov et Krivosheina, 2013 (Fig. 2)

Material examined: East Azerbaijan: Ajabshir, 37°31' N, 46°07' E, 1660 m a.s.l., 10 July 2009; (dry grassland habitat), 1 ♂; Chichakli, 38°39' N, 46°31' E, 2140 m a.s.l., 5 August 2010 (forest-grassland habitat), 2 ♂; (3 ♂, IMTU; 1 ♂ ZMUM).

Distribution: Armenia, Georgia and Russia (Ozerov and Krivosheina, 2013). New to the Iranian insect fauna.

Norellisoma spinimana (Fallén, 1819) (Fig. 3)

Material examined: East Azerbaijan: Chichakli, 38°39' N, 46°31' E, 2140 m a.s.l., 20 June 2009 (forest-grassland habitat), 2 ♀; Kandovan, 37°45' N, 46°18' E, 2844 m, 10 July 2010, 1 ♀; 37°44' N, 46°19' E, 3005 m, 22 May 2012 (moist grassland habitat), 1 ♀; (4 ♀, IMTU; 1 ♀ ZMUM).




Figure 1-4 Habitus, lateral view. **1.** *Cordilura rufipes* (Meigen, 1826) (dry specimen), **2.** *Gimnomera montana* Ozerov et Krivosheina, 2013 (alcoholic specimen), **3.** *Norellisoma spinimana* (Fallén, 1819) (alcoholic specimen), **4.** *Parallelomma albipes* (Fallén, 1819) (dry specimen).

Distribution: Andorra, Austria, Belgium, Bosnia-Herzegovina, China, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Ireland, Italy, Latvia, Macedonia, Netherlands, Norway, Poland, Romania, Russia, Siberia, Slovakia, Spain, Sweden, Nearctic region (Šifner, 2008). New to the Iranian insect fauna.

***Parallelomma albipes* (Fallén, 1819) (Fig. 4)**

Material examined: East Azerbaijan: Gharadagh, 38°51' N, 46°52' E, 1770 m a.s.l., 15 June 2009 (forests habitat), 2 ♀; (1 ♀, IMTU; 1 ♀ ZMUM).

Distribution: Andorra, Austria, Belgium, Bulgaria, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Ireland, Italy, Lithuania, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Siberia, Slovakia, Spain, Sweden,

Switzerland (Šifner, 2008). New to the Iranian insect fauna.

***Scathophaga jezeki* (Šifner, 1981) (Fig. 5, 10, 12)**

Material examined: East Azerbaijan: Maraghe, 37°25' N, 46°25' E, 1787 m a.s.l., 25 June 2009 (dry grassland habitat), 1 ♂, 1 ♀; Chichakli, 38°35' N, 46°14' E, 1219 m

a.s.l., 29 July 2010 (forest- grassland habitat), 2 ♂; Qurigol, 37°54.975' N, 46°41', 120' E, 1943 m a.s.l., 9 Jul. 2010 (dry grassland habitat), 1 ♂, 1 ♀; Kandovan, 37°44' N, 46°19' E, 3005 m a.s.l., 6 June 2011 (moist grassland habitat), 1 ♂, 2 ♀; (2 ♂, 3 ♀, IMTU; 2 ♂ ZMUM).

Distribution: Iran, United Arab Emirates (Šifner, 2009).



Figure 5-8 Habitus, lateral view. **1.** *Scathophaga jezeki* (Šifner, 1981) (alcoholic specimen), **2.** *Scathophaga kaszabi* (Šifner, 1975) (dry specimen), **3.** *Scathophaga lutaria* (Fabricius, 1794) (dry specimen), **4.** *Scathophaga stercoraria* (Linnaeus, 1758) (Fallén, 1819) (alcoholic specimen).




Figure 9-13 9. *Scathophaga taeniopa* (Rondani, 1867) (Habitus, lateral view, dry specimen). 10. *Scathophaga jezeki* (Šifner, 1981) (4th abdominal sternite), 11. *Scathophaga kaszabi* (Šifner, 1975) (4th abdominal sternite), 12. *Scathophaga jezeki* (Šifner, 1981) (epandrium, dorsal view), 13. *Scathophaga kaszabi* (Šifner, 1975) (epandrium, dorsal view).

***Scathophaga kaszabi* (Šifner, 1975) (Fig. 6, 11, 13)**

Material examined: East Azerbaijan: Gharadagh, 38°51' N, 46°52' E, 1770 m a.s.l., 15 June 2009 (forests habitat), 1 ♂, 3 ♀;

Chichakli, 38°35' N, 46°14' E, 1219 m a.s.l., 29 July 2010 (forest- grassland habitat), 2 ♂, 1 ♀; Qurigol, 37°54.975' N, 46°41', 120 E, 1943 m a.s.l., 9 Jul. 2010 (dry grassland habitat), 1 ♀; Kandovan, 37°44' N, 46°19' E, 3005 m a.s.l., 6

June 2011 (moist grassland habitat), 2 ♂, 3 ♀; (4 ♂, 8 ♀, IMTU; 1 ♂ ZMUM).

Distribution: Mongolia (Šifner, 2008). New to the Iranian insect fauna.

***Scathophaga lutaria* (Fabricius, 1794) (Fig. 7)**

Material examined: East Azerbaijan: Gharadagh, 38°53' N, 46°48' E, 1859 m a.s.l., 20 June 2010 (forests habitat), 1 ♂, 1 ♀; Qurigol, 37°54.975 N, 46°41', 120 E, 1943 m a.s.l., 9 July 2012 (dry grassland habitat), 2 ♂; (2 ♂, IMTU; 1 ♂, 1 ♀, ZMUM).

Distribution: Algeria, Austria, Bosnia-Herzegovina, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Israel, Italy, Lebanon, Netherlands, Norway, Poland, Russia, Siberia, Slovakia, Slovenia, Spain, Switzerland, Tunisia, Turkey (Šifner, 2008). New to the Iranian insect fauna.

***Scathophaga stercoraria* (Linnaeus, 1758) (Fig. 8)**

Material examined: East Azerbaijan: Ajabshir, 37°30' N, 46°01' E, 1437 m, 20 June 2009, 2 ♂, 2 ♀, 37°29' N, 45°52' E, 2037 m 15 July 2010, (dry grassland habitat), 3 ♂, 2 ♀; Maraghe, 37°25' N, 46°25' E, 1787 m a.s.l., 25 June 2009 (dry grassland habitat), 2 ♂; Chichakli, 38°39' N, 46°31' E, 2140 m a.s.l., 5 August 2009, 5 ♂, 8 ♀, 38°39' N, 46°31' E, 2140 m, 28 July 2010, (forest-grassland habitat), 3 ♀; Qaradagh, 38°51' N, 46°52' E, 1770 m a.s.l., 14 June 2010, (forest habitat), 10 ♂, 14 ♀; Horand, 38°59' N, 47°22' E, 1370 m, 15 July 2010, 5 ♂, 3 ♀, 38°56' N, 47°27' E, 1360 m, 5 July 2010 (dry to moist grassland habitat), 2 ♂; Kandovan, 37°44' N, 46°19' E, 3005 m a.s.l., 6 June 2011 (moist grassland habitat), 4 ♂, 7 ♀; Qurigol, 37°54.975 N, 46°41', 120 E, 1943 m a.s.l., 9 July 2012 (dry grassland habitat), 5 ♂, 6 ♀; Isperekhan, 37°46' N, 46°24' E, 2504 m, 10 August 2012, (dry grassland habitat), 2 ♂, 2 ♀; (38 ♂, 44 ♀, IMTU; 2 ♂, 3 ♀, ZMUM).

Distribution: Afghanistan, Andorra, Austria, Bosnia-Herzegovina, China, Czech

Republic, Denmark, Egypt, Estonia, France, Finland, Germany, Great Britain, Hungary, Iceland, Ireland, Italy, Japan, Lithuania, Mongolia, Norway, Netherlands, Poland, Portugal, Romania, Russia, Siberia, Far East, Slovakia, Spain, Sweden, Switzerland, Tunisia, Nearctic region, Oriental region (Šifner, 2008; Ozerov, 2010). New to the Iranian insect fauna.

***Scathophaga taeniopa* (Rondani, 1867) (Fig. 9)**

Material examined: East Azerbaijan: Isperekhan, 37°46' N, 46°24' E, 2504 m a.s.l., 8 Aug. 2010, 4 ♂; Chichakli, 38°38' N, 46°14' E, 1429 m a.s.l., 29 Jul. 2009, 1 ♂.

Distribution: Austria, Czech Republic, France, Germany, Italy, Poland, Russia, Slovakia, Switzerland (Šifner, 2008). New to the Iranian insect fauna.

Acknowledgement

The authors express their sincere thanks to Dr. Andrey L. Ozerov (Zoological Museum, Moscow Lomonosov State University) who kindly helped in identification of the materials.

References

- Gorodkov, K. B. 1986. Family Scathophagidae. P. 11–41. In: Soós, Á. & Papp, L. (Eds.). Catalogue of Palaearctic Diptera. Vol. 11. Scathophagidae-Hypodermatidae. Akadémiai Kiadó, Budapest, 346 p.
- McAlpine, J. F. 1981. Morphology and terminology-adults, pp. 9–63. In: McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R., and Wood, D. M., (Eds.). Manual of Nearctic Diptera, Vol. 1: vi + 674 pp. Agriculture Canada Monograph 27, Ottawa.
- Ozerov, A. L. 2009. New species of Scathophagidae (Diptera). Russian Entomological Journal, 17 (4): 419-427.
- Ozerov A. L. 2010. Review of Afrotropical Scathophagidae (Diptera). Russian Entomological Journal, 18 (4): 299-308.

- Ozerov, A. L. and Freidberg, A. 2010. The Scathophagidae (Diptera) of Israel. Israel Journal of Entomology, 40: 169-185.
- Ozerov, A. L. and Krivosheina, M. G. 2013. New and little known Palaearctic species of the family Scathophagidae (Diptera). Far Eastern Entomologist, 270: 1-6.
- Pape, T., Blagoderov, V. and Mostovski, M. B. 2011. Order Diptera Linnaeus, 1758. In: Zhang, Z-Q., (Ed.). Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa, 3148: 222-229.
- Šifner, F. 1981. Les résultats de l'expédition entomologique Tchécoslovaque-iranienne à l'Iran en 1973. Diptera: Scatophagidae (Avec la description d'une espèce nouvelle du genre Coniosternum d'Albanie et avec le tableau analytique des espèces du genre Coniosternum Becker, 1894 paléarctiques). Acta Entomologica Musei Nationalis Pragae, 40: 95-104.
- Šifner, F. 2008. A catalogue of the Scathophagidae (Diptera) of the Palaearctic region, with notes on their taxonomy and faunistics. Acta Entomologica Musei Nationalis Pragae, 48 (1): 111-196.
- Šifner, F. 2009. Additions and corrections to the Catalogue of the Scathophagidae (Diptera) of the Palaearctic Region. Acta Entomologica Musei Nationalis Pragae, 49 (1): 293-295.
- Šifner, F. 2012. *Mirekiana* gen. nov., a new genus of dung flies (Diptera: Scathophagidae) from the Czech Republic. Journal of the National Museum (Prague), 181 (3): 15-19.

فون مگس‌های سرگین (Diptera: Scathophagidae) در استان آذربایجان شرقی، همراه با گزارش هشت رکورد جدید برای ایران

صمد خاقانی‌نیا* و یاسر قراجه‌داغی

دانشگاه تبریز، دانشکده کشاورزی، گروه گیاهپزشکی، تبریز، ایران.
* پست الکترونیکی نویسنده مسئول مکاتبه: skhaghaninia@gmail.com
دریافت: ۵ بهمن ۱۳۹۲؛ پذیرش: ۶ اسفند ۱۳۹۲

چکیده: نمونه‌های مورد مطالعه طی سال‌های ۱۳۸۹ تا ۱۳۹۲ از استان آذربایجان شرقی جمع‌آوری شدند. هشت گونه شامل *Cordilura rufipes* (Meigen, 1826)، *Gimnomera montana* Ozerov et *Parallelomma albipes* (Fallén)، *Norellisoma spinimana* (Fallén, 1819)، *Krivosheina*, 2013، *Scathophaga lutaria* (Fabricius, 1794)، *Scathophaga kaszabi* (Šifner, 1975)، 1819، *Scathophaga taeniopa* (Rondani, 1867) و *Scathophaga stercoraria* (Linnaeus, 1758) و چهار جنس شامل *Cordilura* Fallén, 1810، *Gimnomera* Rondani, 1866، *Norellisoma* Wahlgren, 1917، *Parallelomma* Becker, 1894 برای اولین بار از ایران گزارش می‌شوند. عکس و کلید گونه‌های خانواده Scathophagidae واقع در این منطقه ارائه شده‌است.

واژگان کلیدی: مگس‌های سرگین، فون، گزارش‌های جدید، استان آذربایجان شرقی، ایران