

# **Research Article Identification of bean root rot casual and associated fungal agents in Khomein county, Markazi province, Iran**

Ghazal Sepehri<sup>1</sup>, Shideh Mojerlou<sup>2\*</sup> and Samira Shahbazi<sup>3</sup>

1. Department of Plant Biotechnology, Faculty of Agriculture, Shahrood University of Technology, Shahrood, Iran.

Department of Horticulture and Plant Protection, Faculty of Agriculture, Shahrood University of Technology, Shahrood, Iran.
 Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute (NSTRI), Atomic Energy Organization of Iran (AEOI), Alborz, Iran.

Abstract: Beans are major feed crops belonging to the family Leguminosae, representing one of the most consumed legumes in Iran. Root rot diseases account for high yield losses in bean crops annually, driving the need to identify causative pathogenic agents. During the cropping season of 2019, samples were collected from rhizosphere and infected plant tissues in Khomein county. Of the total 80 purified isolates, 19 fungal isolates were selected for morphological and molecular identification studies. Fusarium equiseti, Fusarium acuminatum, and Macrophomina phaseolina were isolated from the infected crown and root tissues. Their pathogenicity on bean cultivars (Yaghout, Dorsa, and Koosha) is reported for the first time from Markazi province. However, the pathogenicity of fifteen saprophytic and pathogenic fungi species that are isolated from the rhizosphere needs to be confirmed in further studies. Disease symptoms were found to be more severe on pinto beans than red and white beans. In all cases, molecular studies using ITS and tef genomic region confirmed the morphological results. Briefly, these fungal species are introduced as bean root rot causal agents in Khoemin county, which needs to be considered in upcoming breeding programs.

Keywords: Bean, Fusarium spp., molecular identification, root and crown rot

# Introduction

Regarding food security, legumes are the most important agricultural products after cereals, which account for 71% of global consumption in the human food basket (Hayat *et al.*, 2014; Castro-Rosas *et al.*, 2016). According to the World Food Organization, Iran is ranked the 19<sup>th</sup> country globally in bean production with 220740 tons (FAO, 2018). In the cropping season of 2017-18, Fars, Lorestan, Zanjan, Markazi, and Khuzestan provinces had the highest area under cultivation, respectively (Ahmadi et al., 2019). Plant pathogens are among the principal factors responsible for different levels of yield loss in bean crops and affect revenue and food security worldwide (Burke and Hall, 1991; Cerda et al., 2017; Mardani-Mehrabad et al., 2019). Fungal diseases such as Rhizoctonia root and crown rot (Rhizoctonia solani Kuhn), Fusarium wilt Fusarium oxysporum (Mart.) Sacc., Fusarium root rot Fusarium solani (Mart.) Sacc., Ascochyta blotch Ascochyta rabiei (Kovatsch) Arx., Alternaria blight Alternaria alternata (Fr.) Keissl. and Charcoal rot Macrophomina phaseolina (Tassi) Goid. have been reported as some of the severe diseases of legumes in Iran (Ahari Mostafavi et al., 2012).

Handling Editor: Naser Safaie

<sup>\*</sup> Corresponding author: shideh.mojerlou@shahroodut.ac.ir Received: 05 May 2021, Accepted: 14 July 2021 Published online: 03 December 2021

Bean root rot is one of the most important and common diseases in most bean-growing regions of the world. The disease was first observed in 1916 in New York, USA (Burke and Hall, 1991). Significant disease damage occurs under adverse growing conditions and environmental tensions; however, the crop loss depends on weather conditions, soil, and specific cultural practices in each region (Miller et al., 1995; Harveson et al., 2005). The disease also causes substantial yield loss every year in Iran (Moini et al., 1998). In heavily infected areas, it is responsible for the loss in crop yield of 51 up to 85% (Ahari Mostafavi, 2009). Bean root rot diseases are caused by soilborne fungal pathogens such as R. solani, M. phaseolina, Pythium spp., and Fusarium solani f. sp. phaseoli. R. solani has been reported as one of the primary root rot causal agents and results in significant crop loss in Iran, Turkey (Erper et al., 2007), Kenya (Mwangombe et al., 2007), USA (Harveson et al., 2005), Australia (O'brien et al., 1991) and Canada (Hall and Phillips, 1992). The disease is prevalent in bean cultivation areas in Markazi, Charmahal Bakhtyari, and Zanjan provinces (Heydarian and Ershad, 2002; Naseri, 2008). The infection rate of R. solani in Zanjan province has been reported as 50 % by Moini et al. (1998).

In addition, bean root rot caused by F. solani f. sp. phaseoli has been reported in most beangrowing regions of the world (Naseri and Marefat, 2011).). Fusarium species are important soilborne fungi both as a pathogen (Sahampoor et al., 2020) and elicitor (Torkamani et al., 2014) and has been reported as a causal agent of bean root rot in Iran (Kaiser et al., 1968; Nelson et al., 1983; Saremi, 1998). The importance of this fungus is due to various diseases on a variety of crops (Jarvis and Schoemer, 1978; Burgess et al., 1994; Rezaee et al., 2018). The diversity of species and their adaptation to different climatic conditions have caused considerable damages in other countries. which their distribution varies according to the type of climate (Saremi, 2003). Moini et al. (1998) reported the genus Fusarium as the most important agent of bean root rot. Several species of Fusarium have been reported as causal or associated agents of bean root rot in different parts of the world, including; *F. acuminatum*, *F. anthophilum* (Braun) Wollenweber, *F. avenaceum* (Fries) Sacc., *F. culmorum* (Smith) Sacc., *F. equiseti*, *F. oxysporum*, *F. proliferatum* (Mats) Nirenberg, *F. redolens* Wollenweber, *F. crookwellense*, *F. verticillioides* (Sacc.) Nirenberg (Montiel-González *et al.*, 2005; Asan, 2011).

Moreover, F. solani has been reported from rotten bean roots in Zanjan (Khodagholi *et al.*, 2013). Safarloo and Hemmati (2014) reported *Fusarium* species involved in bean root rot in Zanjan province as F. solani, F. acuminatum, F. equiseti, F. oxysporum, F. crookwellense, and F. sambucinum.

Eventually, widespread incidence of some root disease fungi in important bean growing areas of Iran indicates an urgent need for studying the influential factors affecting the occurrence and spread of these diseases. Therefore, disease damage reduction in bean crops serves as one of the research priorities both in Iran and worldwide. Since Khomein is a bean production center in Markazi province, it was considered representative of the region in this research. Hence, this study aimed to identify the associated and causal fungal agents of bean root rot disease in Khomein county.

## **Materials and Methods**

# Sampling and isolation of fungal isolates

Samples were collected from the rhizosphere and infected plants (roots and crowns) in Khomein county, Markazi province, Iran, during the growing season of 2019 to identify causal and associated agents of bean root rot. The samples were randomly chosen from infected plants with visible wilting, yellowing, root, and crown rot and transferred to the laboratory. After surface disinfestation. fragmentation with disease symptoms was cultured on potato-dextrose-agar (PDA) medium by 1% sodium hypochlorite (active ingredient). Cultures were maintained at  $25 \pm 2$  °C. Fungal colonies grown on agar were sub-cultured and purified on PDA and water-agar (WA) media (Ghaneie et al., 2012). For fungal isolation from the rhizosphere, serial dilutions

were prepared from the soil samples and cultured on PDA and Rose Bengal agar media. The obtained colonies were transferred to PDA medium and purified by the hyphal tip and single spore methods on WA medium.

## Morphological and molecular identification

Morphological identification of the isolates was made based on hyphae, conidiophores, and conidia characteristics using identification keys (Barnett and Hunter, 1998; Leslie and Summerel, 2006). For molecular identification, fungal isolates were first cultured in potatodextrose broth (PDB) medium and placed on a shaker at 135 rpm for 5 to 7 days at room temperature. The fungal mass was then collected using sterile filter paper under vacuum conditions. Total genomic DNA was extracted using Safaie *et al.* (2005) method and used as a template for PCR programs. Finally, the quality and quantity of the obtained genomic DNA were examined using electrophoresis on 0.8% agarose gel and nanodrop (IMPLEN N-60), respectively.

ITS1-5.8s-ITS2 region was amplified using ITS1 and ITS4 primers (White *et al.*, 1990; Haratian *et al.*, 2013).). For accurate identification of *Fusarium* species, the *tef-1a* gene (Translation Elongation Factor 1- $\alpha$ ) was amplified using specific primers TEF-4f and TEF-4r (O'Donnell, 2000). Primer sequences are given in Table 1.

Table 1 Name and sequence of the primers used in this study.

Primer Name	Sequence $(5' \rightarrow 3')$	Annealing Temperature (°C)	Reference
ITS1	TCCGTAGGTGAACCTGCGG	56	White et al., 1990
ITS4	TCCTCCGCTTATTGATATGC	56	White et al., 1990
TEF-4r	GGARGTACCAGTSATCATG	58	O'donnell et al., 2000
TEF-4f	ATGGGTAAGGARGACAAGAC	58	O'donnell et al., 2000

The PCR programs were performed with a final volume of 20  $\mu$ l, including; 10  $\mu$ l of the reaction mixture, 1  $\mu$ l of each forward and reverse primer, 1  $\mu$ l of template DNA (50 ng/ $\mu$ l) and the residual volume of deionized water. PCR process for the ITS region amplification includes; an initial denaturation for 5 min at 94 °C, followed by 35 cycles including 1 min at 94 °C, 30 seconds at 57 °C, 1 min at 72 °C, and final extension at 72 °C for 10 min. Besides, the amplification program for *tef1a* gene consisted of an initial denaturation for 5 min at 95 °C, then 30 cycles including 1 min at 95 °C, 75 seconds at 57 °C, and 1 min at 72 °C and final extension at 72 °C for 10 min.

The PCR products were examined by electrophoresis on 1% and 1.2% agarose gel and then sequenced by Biomagic Gene Company (www. Bmgtechno. com). The obtained sequences were compared with the deposited sequences at the NCBI database. Similar sequences were aligned with the obtained sequences in this study using MEGA X software, and the phylogenetic tree was drawn based on the Neighbor-joining method.

#### Pathogenicity test of the fungal isolates

Before planting the seeds, soil samples were prepared from the experimental farm of the Faculty of Agriculture, Shahrood University of Technology, and then sieved to achieve homogeneous soil and added to the perlite + cocopeat mixture to obtain a soil/perlite/cocopeat ratio of 1: 1: 1, and poured in small pots.

Three cultivars of bean seed, red beans (Yaghout cv.), white beans (Dorsa cv.) and pinto beans (Koosha cv.) were received from Markazi Agricultural and Natural Resources Research and Education Center- Khomein Research Station. The seeds were disinfected with 1% sodium hypochlorite for 15 min and then washed and placed in a dark and suitable environment for 4-5 days to germinate. Three germinated seeds were planted in each pot and placed in a growth chamber with a photoperiod

of 8: 16 (L: D) h. Three replications were considered for each fungal isolate and control. The pathogenicity test was repeated twice.

Petri dishes containing a one-week-old fungal colony were washed with sterile distilled water to prepare spore suspension of Fusarium spp. The number of spores was adjusted to  $2 \times 10^6$  spores/ml using a hemocytometer, and seedlings at the twoleaf stage were inoculated by 10 ml of suspension, which was added to the rhizosphere of each plant (Karimian et al., 2010; Keshavarz et al., 2019). Tap water was used as a control. The inoculation was performed twice at 10 days intervals. Then, disease symptoms in pots kept at  $25 \pm 2$  °C were evaluated after 30 to 40 days by removing the seedlings from the soil. Possible disease symptoms such as vellowing, leaf chlorosis, necrosis, wilting, early leaf drop, narrow red streaks on the hypocotyl, root browning, and hollow stalk were examined (Schwartz et al., 2005). In addition, the disinfected roots of inoculated plants were cultured on PDA medium to recover the fungal isolate.

The inoculation with *M. phaseolina* was performed using the method of Alagheh Bandzadeh (2007) so that the colonized toothpicks by the pathogen were used as inoculum. Consequently, infected toothpicks were inserted into the crown at a depth of 2 cm in the

soil, and un-inoculated toothpicks were used as control. The inoculation was carried out on 5-7-leaf seedlings. After that, the plants were kept in a growth chamber for 30 d at  $27 \pm 2$  °C. Finally, disease symptoms were examined in different treatments and compared with the control.

### Results

In the present study, sampling was conducted at ten fields in Khomein county, and 44 samples of soil and infected plant tissues were collected. Subsequently, 80 fungal isolates were purified, and a total of 19 isolates were selected for further molecular and morphological identification studies. The list of studied isolates is given in Table 2. As shown in the table, three species, including F. equiseti, F. acuminatum, and M. phaseolina were isolated from infected plants' crown and root tissues. Other species were isolated from the plants' rhizosphere as common saprophytic and pathogenic fungi. However, functions as pathogen or saprophyte require further investigation on the host plants. Due to the importance of Fusarium species as causative agents of bean root and crown rot, morphological description and molecular study of the identified species in this research are given below.

**Table 2** List of the identified fungal species in the present study.

No.	Code	Origin	Fungal Species	Accession Number
1	SM1	Soil	Metarhizium anisopliae	MV145182
2	SM2	Soil	Alternaria sp.	MV145183
3	SM3	Soil	Aspergillus europaeus	MV145184
4	SM4	Soil	Alternaria sp.	MV145185
5	SM5	Soil	Acrostalagmus luteoalbus	MV145186
6	SM6	Soil	Penicillium sp.	MV145187
7	SM7	Soil	Penicillium sp.	MV145188
8	SM8	Soil	Fusarium solani	MV145189
9	SM9	Soil	Chaetomium globosum	MV145190
10	SM10	Soil	Chaetomium truncatulum	MV145191
11	SM11	Soil	Acrostalagmus luteoalbus	MV145192
12	SM12	Soil	Chaetomium iranianum	MV145193
13	SM13	Soil	Aspergillus sp.	MV145194
14	SM14	Soil	Acremonium sclerotigenum	MV145195
15	SM15	Soil	Fusarium solani	-
16	RB1	Crown and root	F. equiseti	MW551800
17	RB2	Crown and root	F. equiseti	MW551801
18	WB1	Crown and root	F. acuminatum	MW551802
19	WB2	Crown and root	Macrophomina phaseolina	-

# Identified Fusarium species

*F. acuminatum* Ellis and Everhart

The fungal colony grew relatively slowly on PDA medium and produced white mycelia. Macroconidia were formed in pale orange sporodochium. Red (sometimes brown) pigments were formed on the agar medium. Microconidia were absent, but macroconidia had four to six septa. Although *F. acuminatum* is commonly known as a saprophytic and secondary agent invading necrotic tissues, some

isolates can cause severe root rot in some legume species. The morphological characteristics of WB1 isolate in this study were consistent with the descriptions provided by Leslie and Summerel (2006), and the molecular analysis using the *tef1* region also confirmed the morphological findings. The results of the morphological and molecular studies are shown in Figs. 1c, 2d, and 3, respectively.



Figure 1 Colony of *Fusarium* species on PDA medium 5 days after incubation. a) *Fusarium equiseti* isolate RB1, b) *Fusarium equiseti* isolate RB2, c) *Fusarium acuminatum* isolate WB1, d) *Fusarium solani* isolate SM15.



**Figure 2** Macroconidia of *Fusarium* species using 40X magnification, a and b) *Fusarium solani* isolate SM15, c) *Fusarium acuminatum* isolate WB1, d) *Fusarium equiseti* isolate RB2, e and f) *Fusarium equiseti* isolate RB1.

#### F. equiseti (Corda) Saccardo

Macroconidia were abundant and had a thick wall with a strong posterior dorsal curve and 5 to 7 septa. Chlamydospores were formed rapidly and turned to pale brown, golden, or brown color by aging. The mycelium formed in the PDA medium was initially white and turned brown with age. The morphological characteristics of RB1 and RB2 isolates in this study were consistent with the descriptions provided by Nelson *et al.* (1983), and molecular analysis using the *tef 1* region also confirmed the morphological findings. The results of the morphological study are shown in Fig. 1 (a and b) and Fig. 2 (d, e, and f), and the results of the molecular study are shown in Fig. 3.

F. solani (Martius) Appel and Wollenweber emend.

The fungal colony had white to creamy mycelium on PDA medium. Macroconidia were slightly curved with 3 to 7 septa and rounded ends. Macroconida were abundant and creamy in color. Oval and reniform microconidia were also observed. These characteristics distinguish this species from a similar species, *F. oxysporum*. Chlamydospores usually form quickly within 2-4 weeks. The morphological characteristics of the isolate SM15 in this study were consistent with the descriptions provided in the identification keys (Summerbell, 2003; Larone, 2011), and molecular analysis using the *tef 1* region also confirmed the morphological findings. The results of the morphological and molecular studies are shown in Figs. 1d, 2a, b, and 3, respectively.

## Pathogenicity test of the fungal isolates

The disease symptoms appeared 30 days after inoculation on the upper- and underground parts of the plants of all cultivars with the tested pathogens. Most of the symptoms were in the form of crown wounds and brown discoloration of secondary roots. The symptoms observed on seedlings treated with *F. equiseti*, *F. acuminatum*, and *M. phaseolina* isolates are shown in Fig. 4.

As the last step of the pathogenicity test, the fungal agents were reisolated from the wounds of the infected plants. Morphological studies showed that these isolates were quite similar to the ones initially used for inoculation.

This study showed the most apparent disease symptoms in the plants treated with the pathogens F. acuminatum and F. equiseti in Dorsa, Koosha, and Yaghout cultivars. No disease symptoms were observed in roots, crowns, and aerial parts of the control plants. Symptoms varied depending on the studied species. In inoculated plants with M. phaseolina, hollow stalk and leaf drop were frequently observed before maturity. In infected plants by F. equiseti and F. acuminatum, symptoms such as leaf chlorosis and necrosis, wilting, narrow red streaks on hypocotyl, root, and especially secondary root browning were observed. Besides, vascular tissue browning was usually kept in the cross-section of the stem. The root rot symptom severity varied from reddishbrown to dark brown.



**Figure 3** Phylogenetic position of the studied *Fusarium* species. The most parsimonious tree obtained by *tef-1*  $\alpha$  gene sequence using Neighbor Joining method, MEGA X software. The bootstrap support from 1000 replications is indicated on the branches. *Alternaria alternata* is used as outgroup.



# Discussion

In this study, several pathogenic and associated fungal agents were isolated from the tissue and rhizosphere of the infected bean plants. Among them, the purified fungal isolates from the infected plants were picked for further studies. According to aghout cultivar, b) *Fusarium cuminatum* on Dorsa cultivar, c) *facrophomina phaseolina* on Koosha iltivar. the earlier studies, different *Fusarium* species such as *F. solani*, *F. oxysporum*, *F. crookwellense*, *F.* 

as *F. solani*, *F. oxysporum*, *F. crookwellense*, *F. equiseti*, *F. moniliform*, *F. acuminatum*, and *F. sambacinum* have been reported as root and crown rot causal agents in various hosts (Latani *et al.*, 2006; Khodagholi *et al.*, 2013; Safarloo and Hemmati, 2014).

Azimi *et al.* (2005) isolated 84 isolates belonging to 6 species of *Fusarium* from the bean plants' crown, root, and rhizosphere. The species *F. solani*, *F. oxysporum*, and *F. moniliform* were the most abundant, and *F. equiseti*, *F. semitectum*, and *F. prolifratum* had the lowest frequency. In the present study, *F. acuminatum* and *F. equiseti* were identified as causative agents of bean crown and root rot from Khomein county, consistent with the results of other researchers mentioned above.

As shown in Fig. 3, both F. solani and F. virguliforme were grouped in one cluster along with SM15 isolate. The F. solani complex (FSSC) contains more than 60 phylogenetically distinct species (O'Donnell, 2000; Zhang et al., 2006; O'Donnell et al., 2008; Nalim et al., 2011). The only fusarium species currently recognized as species complex is F. solani. Most group members have been identified as phytopathogens and divided into formae speciales based on their hosts (O'Donnell, 2000; O'Donnell et al., 2008; Coleman, 2016). Soybean sudden death syndrome (SDS) is a root and crown rot disease of soybean Glvcine max (L.) Merr is traditionally caused by Fusarium solani f. sp. glycines (Roy et al., 1997). Four pathogens, including F. virguliforme, F. tucumaniae, F. brasiliense, and F. cuneirostrum, cause SDS and are members of FSSC (Aoki et al., 2003, 2005). The species F. virguliforme has a wide host range and can cause root necrosis on different host species, including alfalfa, pinto and navy beans, white and red clover, pea, and Canadian milk vetch. However, they cause no symptoms on some other plants (Kolander et al., 2012). The current study introduced the isolate SM15 as F. solani species according to morphological characteristics and host species.

*M. phaseolina*, the causative agent of root rot and charcoal rot, is one of the most important plant pathogens in hot and dry regions and has a widespread distribution and extensive host range (Su *et al.*, 2001). The disease severity varies depending on the host susceptibility, suitable climatic conditions including high temperature and low humidity, soil contamination, and irrigation system (Su *et al.*, 2001). Naseri (2007) reported *M*. *phaseolina* for the first time from the bean plants in Zanjan province and proved its pathogenicity on pinto, white and red beans. This pathogen is reported as a causal agent of root rot disease in bean plants from Khomein county in the present study. Also, Jafari Petroudi *et al.* (2014) considered charcoal rot disease caused by *M. phaseolina* as one of the limiting factors of crop cultivation in Iran, which represents the most important fungal disease of soybean, especially in the northern regions of Iran and even in the north of Khuzestan (Orojnia *et al.*, 2021).

Regarding the pathogenicity of Fusarium isolates on bean cultivars, Khodagholi et al. (2013) demonstrated the pathogenicity of F. solani on red beans (Naz cultivar). They evaluated the susceptibility of three-bean cultivars to the pathogen and reported that disease severity was higher on pinto beans than red and white beans. Naseri and Marefat (2011) also reported higher pinto and white beans contamination than red beans from Zanjan Therefore, province. red beans are recommended in areas with a long history of contamination rather than white and pinto beans. In the present study, the contamination in pinto beans was more severe than red and white beans based on disease symptoms and agreed with the results of the two studies mentioned above.

The host plants' symptoms caused by F. solani and F. equiseti have been reported as root and crown rot. Root rot usually appears as dark brown or completely black discoloration, mainly of secondary roots. Also, the rot occurring on the crown area is primarily dark reddish-brown, and sometimes stems turn black and thin, and leaves are small, dark, and completely wrinkled (Azimi et al., 2005). In another study, root and crown necrosis and cracks in infected parts of the plant due to infection caused by F. solani and F. equiseti have been reported. Moreover, in the longitudinal section of roots and crowns, all species show varying degrees of internal tissue discoloration in infected areas. Root and crown rot caused by different species vary in color, from red to light brown to dark. Aerial

symptoms included a range of symptoms such as chlorosis, slow growth, stunting, mosaic, and wilting. The type and severity of symptoms caused by different species are diverse (Safarloo and Hemmati, 2014). The symptoms observed in infected plants with *F. equiseti* and *F. acuminatum* pathogens in the present study were consistent with those of other researchers. They were similar to the symptoms mentioned above.

Finally, due to the importance of Khomein county as a bean seed production center in the country, susceptibility evaluation of typical local and commercial cultivars to *F. acuminatum, F. equiseti, F. solani* pathogens is suggested. Also, the resistance of bean cultivars to these pathogens should be included in breeding programs.

## References

- Ahari Mostafavi, H. 2009. Evaluation of the possibility of biological control of bean root rot disease using non-pathogenic mutants of *Fusarium solani* f. sp. *phaseoli*. Journal of Plant Production Research, 16: 135-149.
- Ahari Mostafavi, H., Mirmajlessi, S. M., Safaie, N., Minassyan, V., Fathollahi, H., Dorri, H.
  R. and Mansouripour, S. M. 2012. The use of a gamma-irradiated mutants of *F. solani* f.
  sp. *phaseoli* with reduced pathogenicity for the biological control of fusarium root rot of bean (*Phaseolus vulgaris*) in field conditions. Journal of Agricultural Science and Technology, 14: 1415-1423.
- Ahmadi, K., Ebadzadeh, H., Hatami, F.,
  Abdeshah, H. and Kazemian, A. 2019.
  Agricultural statistics of the crop year 2017-2018: Volume 1: crops. Ministry of
  Agriculture Jihad Publications, Tehran, Iran.
- Alagheh Bandzadeh, N. 2007. Pathogenic diversity and genetic diversity of *Macrophomina phaseolina* isolates. Master thesis, Faculty of Agriculture, Science and Research Branch, Islamic Azad University, Tehran, Iran.
- Aoki, T., O'Donnell, K. and Scandianim, M. M. 2005. Sudden death syndrome of soybean in South America is caused by four species of

Fusarium: *Fusarium brasiliense* sp. nov., *F. cuneirostrum* sp. nov., *F. tucumaniae*, and *F. virguliforme*. Mycoscience, 46: 162-183.

- Aoki, T., O'Donnell, K., Homma, Y. and Lattanzi, A. R. 2003. Sudden-death syndrome of soybean is caused by two morphologically and phylogenetically distinct species within the *Fusarium solani* species complex-*F. virguliforme* in North America and *F. tucumaniae* in South America. Mycologia, 95: 660-684.
- Asan, A. 2011. Checklist of *Fusarium* species reported from Turkey. Mycotaxon, 116: 479.
- Azimi, S., Farokhi Nezhad, R. and Mousavi Jorf, S. A. 2005. Identification and study on pathogenicity of *Fusarium* spp. associated with foot and root of Faba Bean in Khuzestan. Scientific Journal of Agriculture, 28: 149-164.
- Barnett, H. L. and Hunter, B. B., 1998. Illustrated genera of imperfect fungi (4<sup>th</sup> Ed.). APS Press, St. Paul. pp. 218.
- Burgess, L. W., Summerel, B. A., Bullock, S., Gott, K. P., Backhouse, D. 1994. Laboratory Manual for Fusarium Research (3<sup>rd</sup> Ed.). Department of Crop Science, University of Sydney/Royal Botanic Gardens. pp. 134.
- Burke, D. W. and Hall, R. 1991. Fusarium root rot. In: Hall, R. (Ed.), Compendium of Bean Diseases. APS Press, St. Paul. pp. 9-10.
- Castro-Rosas, J., Aguirre-Santos, E. A., Gómez-Aldapa, C. A., Valle-Cervantes, S., Ochoa-Martínez, L. A., Hernández-Santos, B. and Rodríguez-Miranda, J. 2016. Effect of harvest year on the physical properties, chemical composition and cooking time of three common bean varieties that are grown in Mexico. Quality Assurance and Safety of Crops and Foods, 8: 339-348.
- Cerda, R., Avelino, J., Gary, C., Tixier, P., Lechevallier, E. and Allinne, C. 2017. Primary and secondary yield losses caused by pests and diseases, assessment and modeling in coffee. PloS one, 12: e0169133.
- Coleman, J. J. 2016. The *Fusarium solani* species complex, ubiquitous pathogens of agricultural importance. Molecular Plant Pathology, 17: 146-158.

- Erper, I., Karaca, G. H. and Ozkoc, I. 2007. Root rot disease incidence and severity on some legume species grown in Samsun and the fungi isolated from roots and soils. Archives of Phytopathology and Plant Protection, 41: 501-506.
- FAO. 2018. FAO statistical yearbook, Agricultural production, Food and Agriculture Organization of the United Nations. Available from: http://faostat.fao.org.
- Ghaneie, A., Mehrabi, R., Safaie, N., Abrinbana, M., Saidi, A. and Aghaee, M. 2012. Genetic variation for resistance to *Septoria tritici* blotch in Iranian tetraploid wheat landraces. European Journal of Plant Pathology, 132(2): 191-202.
- Hall, R. and Phillips, L. G. 1992. Effect of crop sequence and rainfall on population dynamics of *Fusarium solani* f. sp. *phaseoli* in soil. Canadian Journal of Botany, 70: 2005-2008.
- Haratian, M., Safaie, N., Sharifnabi, B., Mahmudi, S. B. and Ariana, A. 2013.
  Genetic structure of populations of *Rhizoctonia solani* AG□4 from five provinces in Iran. Plant Pathology, 62(3): 649-656.
- Harveson, R. M., Smith, J. A. and Stroup, W. W. 2005. Improving root health and yield of dry beans in the Nebraska Panhandle with a new technique for reducing soil compaction. Plant Disease, 89: 279-284.
- Hayat, I., Ahmad, A., Masud, T., Ahmed, A. and Bashir, Sh. 2014. Nutritional and health perspectives of Beans (*Phaseolus vulgaris* L.): an overview. Critical Reviews in Food Science and Nutrition, 54: 580-592.
- Heydarian, A. and Ershad, J. 2002. Identification and study of fungi causing crown and root rot of pinto beans in Chaharmahal and Bakhtiari province, In: Proceedings of the 15<sup>th</sup> Iranian Plant Protection Congress.7-11 September 2002, Kermanshah, Iran. pp. 15.
- Jafari Petroudi, S. F., Tajick Ghanbari, M. A., Babaeizad Keshtli, V. and Qasemi Imran, V. 2014. Evaluation of severity of pathogenicity of *Macrophomina phaseolina*

on some soybean cultivars. In: The first national electronic congress of new technologies in Iran with the aim of achieving sustainable development, 4 February 2014, Tehran, Iran. pp. 4977.

- Jarvis, W. R. and Shoemer, R. A. 1978. Taxonomic status of *Fusarium oxysporum* causing foot and root rot of tomato. Phytopathology, 68: 1679-1680.
- Kaiser, W. J., Danesh, D., Okhovvat, M. and Mossahebi, G. H. 1968. Diseases of Pules crop (edible legumes) in Iran. Plant Disease Report, 52: 681-691.
- Karimian, B., Javan-Nikkhah, M., Abbasi, M. and Ghazanfari, K. 2010. Genetic diversity of *Fusarium oxysporum* isolates from common bean and distribution of mating type alleles. Iranian Journal of Biotechnology, 8: 90-97.
- Keshavarz, M., Sadravi, M. and Abdollahi, M. 2019. Biological control of bean Fusarium wilt by silver nanoparticle extract of *Trichoderma harzianum*. Biocontrol in Plant Protection, 6: 57-66.
- Khodagholi, M., Hemmati, R., Naseri, B. and Marefat, A. 2013. Genotypic, phenotypic and pathogenic diversity of *Fusarium solani* isolates causing bean root rot in Zanjan province. Iranian Journal of Cereals Research, 4: 11-125.
- Kolander, T. M., Bienapfl, J. C., Kurle, J. E., and Malvick, D. K. 2012. Symptomatic and asymptomatic host range of *Fusarium virguliforme*, the causative agent of soybean sudden death syndrome. Plant Disease, 96: 1148-1153.
- Larone, D. H. 2011. Medically Important Fungi, A Guide to Identification. ASM Press, Washington. pp. 305.
- Latani, S., Taheri, A. and Rahnama, K. 2006. Identification and pathogenicity of *Fusarium* species isolated from the roots and crown of Gorgan region. Journal of Agricultural Sciences and Natural Resources, 13: 115-126.
- Leslie, J. F. and Summerel, B. A. 2006. Taxonomy and Identification of *Fusarium*. In: Leslie J. F., Summerel, B. A. (ed),

Downloaded from jcp.modares.ac.ir on 2024-10-07 ]

Fusarium laboratory manuals. Blackwell publishing, Iowa, USA.

- Mardani-Mehrabad, H., Rakhshandehroo, F., Shahbazi, S. and Shahraeen, N. 2019. Study on the effect of low doses of gamma irradiation alone or in associated with the salicylic acid and indirect temperature on seed borne infection of Bean common mosaic virus in bean plant. Applied Entomology and Phytopathology, 87: 87-105.
- Miller, S. A., Riedel, R. M. and Rowe, R. C. 1995. Damping-off and Root Rot of Beans. The Ohio State University Extension Fact Sheet, Ohio, USA.
- Moini, M., Ahmadinejad, A. and Shahraeen, N. 1998. Status of bean diseases in Zanjan province. In: Proceedings of the 13th Iranian Plant Protection Congress. 23-27 August, University of Tehran, Karaj, Iran. pp. 154.
- Montiel-González, L., González-Flores, F., Sánchez-García, B. M., Guzmán-Rivera, S., Gámez-Vázquez. F. P., Acosta-Gallegos, J. A., Rodriguez-Gurra, R., Simpson-Wiliamson, J. and Mendoza-Elos, M. 2005. *Fusarium* species on bean (*Phaseolus vulgaris* L.) roots causing rots, in five states of Central Mexico. Revista Mexicana de Fitopatologia, 23: 1-10.
- Mwangombe, A. W., Thiongo, G., Olubayo, F. M. and Kiprop, E. K. 2007. Occurrence of root rot disease of common bean (*Phaseolus vulgaris* L.) in association with bean stem maggot (*Ophyiomia* sp.) in Embu District, Kenya. Plant Pathology Journal, 6: 141-6.
- Nalim, F. A., Samuels, G. J., Wijesundera, R. L. and Geiser, D. M. 2011. New species from the *Fusarium solani* species complex derived from perithecia and soil in the Old-World tropics. Mycologia, 103: 1302-1330.
- Naseri, B. 2007. The first report of *Macrophomina phaseolina*, the causative agent of bean root coal rot in Zanjan, Iran. Journal of Plant Diseases, 43: 276-277.
- Naseri, B. 2008. Root rot of common bean in Zanjan, Iran: major pathogens and yield loss estimates. Australasian Plant Pathology, 37: 546-551.

- Naseri, B. and Marefat, A. 2011. Large-scale assessment of agricultural practices affecting Fusarium root rot and common bean yield. European Journal of Plant Pathology, 131: 179-195.
- Nelson, P. E., Toussoun, T. A. and Marasas, W. F. O. 1983. Fusarium species: an illustrated manual for identification. The Pennsylvania State University Press, University Park. Pennsylvania, USA. pp. 193.
- O'brien, R. G., O'hare, P. J. and Glass, R. J. 1991. Cultural practices in the control of bean root rot. Australian Journal of Experimental Agriculture, 31: 551-555.
- O'Donnell, K. 2000. Molecular phylogeny of the *Nectria haematococca-Fusarium solani* species complex. Mycologia, 92: 919-938.
- O'Donnell, K., Kistler, H. C., Tacke, B. K., Casper, H. H. 2000. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. Proceedings of the National Academy of Sciences, USA, 97: 7905-7910.
- O'Donnell, K., Sutton, D. A., Fothergill, A., McCarthy, D., Rinaldi, M. G., Brandt, M. E., Zhang, N., Geiser, D. M. 2008. Molecular phylogenetic diversity, multilocus haplotype nomenclature, and *in vitro* antifungal resistance within the *Fusarium solani* species complex. Journal of Clinical Microbiology, 46: 2477-2490.
- Orojnia, S., Habibi, D., Shahbazi, S., Paknejad, F. and Ilkaee, M. N. 2021. Investigation the effect of different formulations of mutated Trichoderma with gamma ray on Soybean morphological indices. Romanian Agricultural Research, 3: 97-114.
- Rezaee, S., Gharanjik, Sh. and Mojerlou Sh. 2018. Identification of *Fusarium solani* f. sp. *cucurbitae* races using morphological and molecular approaches. Journal of Crop Protection, 7(2): 161-170.
- Roy, K. W. 1997. *Fusarium solani* on soybean roots: nomenclature of the causal agent of sudden death syndrome and identity and

relevance of *F. solani* form B. Plant Disease, 81: 259-266.

- Safaie, N., Alizadeh, A., Saidi, A. and Rahimian H. 2005. Molecular characterization and genetic diversity among Iranian populations of *Fusarium graminearum* isolates, the causal agent of wheat scab. Iranian Journal of Plant Pathology, 41: 69-74.
- Safarloo, Z. and Hemmati, R. 2014. Identification and study of pathogenicity of *Fusarium* species involved in bean root rot in Zanjan province. Applied Research in Phytomedicine, 3: 99-77.
- Sahampoor, L., Zaker Tavallaie, F., Fani, S. R. and Shahbazi, S. 2020. In vitro efficiency of *Trichoderma harzianum* mutants in biocontrol of *Fusarium oxysporum* f. sp. *radicis-cucumerinum*. Journal of Crop Protection, 9: 285-300.
- Saremi, H. 1998. Ecology and taxonomy of *Fusarium* species, University of Mashhad Press, Iran. pp. 132.
- Saremi, H. 2003. The distribution pattern of *Fusarium* species in different climates. Plant diseases, 39: 83-72.
- Schwartz, H. F., Steadman, J. R., Hall, R. and Forster, R. L. 2005. Compendium of Bean Diseases. American Phytopathological Society Press, MN, USA.

- Su, G., Suh, S. O., Schneider, R. W. and Russin, J. S. 2001. Host specialization in the charcoal rot fungus *Macrophomina phaseolina*. Phytopathology, 91: 120-126.
- Summerbell, R. C., 2003. Aspergillus, Fusarium, Sporothrix, Piedraia and their relatives. Pathogenic fungi in humans and animals, 2: 237-498.
- Torkamani, M., Jafari, M., Abbaspour, N., Heidary, R., and Safaie, N. 2014. Enhanced production of valerenic acid in hairy root culture of *Valeriana officinalis* by elicitation. Open Life Science, 9: 853-863.
- White, T. J., Bruns, T., Lee, S. J. W. T. and Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M. A., Gelfand, D. H., Sninsky, J. J. and White, T. J. (Eds.), PCR Protocols: A Guide to Methods and Applications. PCR Protocols: A guide to methods and applications. Massachusetts, USA. pp. 315-322.
- Zhang, N., O'Donnell, K., Sutton, D. A., Nalim, F. A., Summerbell, R. C., Padhye, A. A. and Geiser, D. M. 2006. Members of the *Fusarium solani* species complex that cause infections in humans and plants are common in the environment. Journal of Clinical Microbiology, 44: 2186-2190.

# شناسایی عوامل قارچی بیماریزا و همراه پوسیدگی ریشه لوبیا در شهرستان خمین (استان مرکزی)

غزل سپهری'، شیده موجرلو<sup>\*\*</sup> و سمیرا شهبازی<sup>"</sup>

۱- گروه بیوتکنولوژی، دانشکده کشاورزی، دانشگاه صنعتی شاهرود، شاهرود، ایران. ۲- گروه باغبانی و گیاهپزشکی، دانشکده کشاورزی، دانشگاه صنعتی شاهرود، شاهرود، ایران. ۳- گروه گیاهپزشکی، پژوهشکده کشاورزی هستهای، پژوهشگاه علوم و فنون هستهای، سازمان انرژی اتمی، کرج، ایران. پست الکترونیکی نویسنده مسئول مکاتبه: shideh.mojerlou@shahroodut.ac.ir دریافت: ۱۵ اردیبهشت ۱۴۰۰؛ پذیرش: ۲۳ تیر ۱۴۰۰

چکیده: لوبیا از محصولات مهم خانواده حبوبات و یکی از پرمصرفترین و مهمترین حبوبات در ایران میباشد. هر ساله بیماری پوسیدگی ریشه لوبیا سبب بروز خسارت میشود، از اینرو شناسایی قارچهای بیماریزا ضروری میباشد. در طول فصل زراعی سال ۱۳۹۸، نمونهبرداری از ریزوسفر و بافت گیاهی آلوده در مزارع شهرستان خمین انجام شد. در مجموع، ۱۹ جدایه از بین ۸۰ جدایه قارچی بهدست آمده و خالصسازی شده برای مطالعات ریختشناسی و مولکولی انتخاب شدند. سه گونه *Eusarium equiseti* و خالصسازی شده برای مطالعات ریختشناسی و مولکولی انتخاب شدند. سه گونه *Eusarium equiseti* و خالصسازی شده برای مطالعات ریختشناسی و مولکولی انتخاب شدند. سه گونه *Eusarium equiseti* و قراری آنها از استان مرکزی گزارش میشود. بیماریزایی این گونهها روی ارقام لوبیا چیتی، سفید و قرمز بررسی و تأیید شد. همچنین، ۱۵ گونه قارچی از خاک اطراف ریشه گیاهان جداسازی شدند کـه شامل قارچهای ساپروفیت و بیمارگر میباشند و تأیید بیماریزایی آنها نیاز به بررسیهای بیشتر و ازمون بیماریزایی روی گیاه میزبان دارد. در این بررسی شدت علایم آلودگی در لوبیا چیتی بیشتر و اوبیا قرمز و سفید بود. در تمامی موارد نتایج بررسیهای مولکولی با استفاده از ناحیه TI و *ter* رازمون بیماریزایی روی گیاه میزبان دارد. در این بررسی شدت علایم آلودگی در لوبیا چیتی بیشتر و تاییدکننده نتایج بررسیهای ریختشناسی بود. به طور خلاصه، این عوامل قارچی برای اولین بار بهعنوان برامه های اصلاحی آینده مورد توجه قرار گیرد.

واژگان کلیدی: لوبیا، پوسیدگی ریشه و طوقه، .Fusarium spp، شناسایی مولکولی