

Short Paper

Distribution and oviposition characteristics of *Orosanga japonica* Melichar (Hemiptera: Ricaniidae) in Guilan province, Iran

Seyedeh Elham Yazdani Badabi* and Ahad Sahragard

Department of Plant Protection, Faculty of Agricultural Sciences, Guilan University, Rasht, Iran.

Abstract: This study aimed to investigate the oviposition behavior, distribution, and host plants of *Orosanga japonica* (Hemiptera: Ricaniidae) in Guilan province, Iran. Random sampling was conducted across various locations within Guilan province. Adult insects typically emerge in late July and deposit their eggs from early August to mid-September. This species prefers laying eggs on young shoots and leaf midribs of various plants, from annual herbaceous species to shrubs. This oviposition behavior can lead to stem desiccation. Additionally, the females' oviposition behavior, sap-sucking activities on stems, and the potential transmission of pathogenic fungi can cause substantial damage to host plants. This research provides information regarding the distribution of pest egg masses across several host plants. Our findings revealed that the mean number of egg masses per twig was higher on chinaberry, averaging 13.75, than on bitter orange, elm trees, and mulberry. The longest average egg mass length, measuring 16.1 mm, was observed on mulberry. Furthermore, this study confirmed the presence of *O. japonica* in 13 cities within Guilan province, including Rasht, Anzali, Amlash, Astaneh, Lahijan, Langroud, Rezvanshahr, Rudsar, Rudbar, Sowme'eh Sara, Siahkhal, Shaft, and Fouman. The insect was predominantly found infesting kiwi, tea, beans, corn, eggplants, peppers, elm leaf blackberries, chinaberries, mulberries, Japanese spindles, cucumbers, citrus plants, fig-trees, rice, cabbage, and grapes.

Keywords: *Orosanga japonica*, Fulgoromorpha, egg masses, distribution, oviposition characteristics, Guilan

Introduction

The planthopper family, Ricaniidae (Hemiptera: Auchenorrhyncha), consists of 67 genera (2.7% of the Fulgoromorpha) and covers 437 species (3.2% of the Fulgoromorpha) (Bourgoin, 2021).

The Family Ricaniidae is mainly distributed in Tropical regions. This family is represented only by the genus *Ricania* in the Palaearctic

region. *Orosanga japonica* (Melichar) (Hemiptera: Ricaniidae) is an invasive species in the western Palearctic region and widespread in the Black Sea area countries. It is a common species in Eastern Asian countries, including Korea and China, and the species has been spread in Ukraine, Russia, Georgia, Turkey, and Azerbaijan (Dlabola, 1967; Nast, 1972, 1987; Gjonov, 2011; EPPO, 2017; Bourgoin, 2017; Demir, 2009, 2018; Ismaylova, 2021).

Handling Editor: Ali Asghar Talebi

* Corresponding author: elyazd2020@gmail.com

Received: 27 February 2022, Accepted: 30 August 2023

Published online: 17 September 2023

It was reported that it caused major damage in tea plantations and kiwi gardens in Turkey's eastern Black Sea coast (Demir, 2009; Ak *et al.*, 2015; Akiner *et al.*, 2019). In Iran, for the first time, this species was recorded in northern Iran (Mazandaran province) in 2010 (Mozaffarian, 2018). Recorded distribution in Iran is in the North, along the shores of the Caspian Sea (Mozaffarian, 2018). The large populations of the nymphs and adults of *O. japonica* cause serious damage to plants. When the females lay their eggs, let them dry. Nymphs and adults cause fumagine by taking the sap of the plant, and it dries out the plant. Furthermore, they cause indirect damage by weakening the plant. This situation makes the plant vulnerable to the fungal pathogen *Pestalotiopsis quepinii* (Eken *et al.*, 2013).

According to the studies conducted in Iran and Turkey, The planthopper overwinters in the egg stage and has five nymphal stages. Depending on climatic conditions, the nymphs emerge from early to late May. The nymphal duration was 41.39 and 38.13 days on *R. ulmifolius* and *M. azedarach*, respectively (Akiner *et al.*, 2019; Altas and Ak, 2019; Karatas *et al.*, 2020; Abbaszadeh *et al.*, 2021). Based on the findings of Ismaylova in Azerbaijan, pest development from the egg to adult stage takes about 2-2.5 months. In late August, females lay eggs under the bark of thin branches and stems (Ismaylova, 2021).

O. japonica is a polyphagous species, and it is found on many plants wherever it is distributed e.g., hazelnut, wild blackberry, chestnut, kiwi, cherry, pineapple, tea, mulberry, common bean, corn, tomatoes, eggplant, cabbage, peppers, cucumbers, citrus plants, common grape vine, apple tree, pear tree, peach tree, fig tree, strawberry, etc. (Gokturk and Aksu, 2014).

So far, several studies have been conducted on the characteristics of spawning of this genus (Choi *et al.*, 2012; Kim *et al.*, 2016; Yu *et al.*, 2016). Ismaylova (2021) photographed eggs of *O. japonica* under an electron microscope and recorded dimensions. However, so far, there is no report about the oviposition characteristics of *O. japonica*. Therefore, this study aims to

investigate females spawning behavior, such as the number and length of egg mass per laying unit (twig, leaf, thorn). This planthopper has quickly spread since the first report of its occurrence in Iran (Mazandaran Province) in 2010. Therefore, this study was carried out to find the current distribution of the pest in Guilan province.

Materials and Methods

To investigate the spawning characteristics of *O. japonica* in the area of Rasht and Anzali, which occurred from August to December of 2021, four host plants, bitter orange, elm tree, chinaberry, and mulberry with egg masses, were selected.

Branches, leaf midribs, and thorns with clutches of eggs were randomly cut with scissors and brought to the laboratory, and number of egg masses and length of egg mass on the substrate were observed in ten replicates. Statistical analysis was performed using descriptive statistics.

To determine the pest distribution in Guilan province, gardens and fields were observed randomly, and each developmental stage of the planthopper was recorded.

Results and Discussion

Results of examining four host plant species of four families indicated that the length of the egg masses laid by the females differed depending on the substrate host. Average length of egg masses was 16.1 mm on mulberry and 11.06 mm on chinaberry, respectively. The number of egg masses per twig of chinaberry was 13.75, the highest compared to other trees, while it was the least, 3.4, on mulberry (Table 1).

Maximum and minimum diameter of twigs were 21.6 and 4 mm on the chinaberry and Elm tree, respectively. The length of the egg mass of *Ricania* spp. Investigated by Choi *et al.* (2012) was 15.2 to 18.0 mm between host plants, which is similar to the result of this study. They reported the number of egg mass 2.2 to 14.5 per twig, which was 3.4 to 13.75 in this study.

Table 1 Oviposition characteristics of *Orosanga japonica* on several kinds of host plants.

Family	Host plants	Mean length of twig/ leaf midrib /thorn (cm)	Mean no. of egg mass per twig/ leaf midrib / thorn	Length of egg mass per twig/ leaf midrib / thorn (mm, Mean \pm SD)	Diameter of twig/thorn where egg mass is counted (mm, Mean \pm SD) ²
Rutaceae	<i>Citrus aurantium</i>	25.3/15.5/4.98	5.5/1.5/1.5	11.6 \pm 1.66	11.9 \pm 1.87
				15.5 \pm 1.88	6.2 \pm 0.34
				12.4 \pm 0.49	
Meliaceae	<i>Melia azedarach</i> ¹	26.0	13.8	11.1 \pm 1.86	21.6 \pm 1.86
Ulmaceae	<i>Ulmus minor</i> ¹	15.5	4.1	14.8 \pm 5.07	4.0 \pm 0.95
Moraceae	<i>Morus alba</i> ¹	28.0	3.4	16.1 \pm 1.34	17.8 \pm 2.23

¹. The data presented in the table refer to twigs.

² Diameter of the twig/thorn at the location where the egg mass is counted.

The mean length of egg masses of *Ricania* sp. recorded on apples and plums was 12.35 and 11.45 mm, respectively (Choi *et al.*, 2011). The mean number of egg masses on mulberry twigs was 3.4, but the length of egg masses was greater (16.1 mm) than on other trees. In Bitter orange, in addition to spawning on branches, it was also observed on leaf midribs and thorns. The length of the egg mass on the leaf midrib was 15.5 mm, while it was 11.6 and 12.4mm on twig and thorn, respectively. The number of egg masses per twig, leaf midrib, and thorn were 5.5, 1.5 and 1.5, respectively. Accordingly, the number of egg masses per leaf midrib and thorn is lower than on twig.

In *Ricania speculum*, eggs are laid in clusters inside the young twig tissues with a diameter rarely

larger than 3 mm or in the leaf midrib (Lucchi and Rossi, 2016). Because the first-year-old branches of chinaberry are less lignified and have a larger twig diameter than other trees, they exhibit a higher mean number of egg masses than the other plants. These results indicate that the degree of tenderness of the host plants affects spawning behavior. The pest was observed in 13 cities, and it was mostly found on kiwi, tea, bean, corn, eggplant, peppers, elm leaf blackberry, chinaberry, mulberry, Japanese spindle, cucumbers, citrus plants, fig-trees, rice, cabbage, and grape. The planthopper has been introduced from Mazandaran to the east of Guilan (Roudsar). It is spreading outwards in all directions to cover a larger area and is dispersing towards the west of Guilan (Fig. 1).

**Figure 1** Distribution of *Orosanga japonica* in Guilan province.

Since chinaberry has the highest number of egg masses per twig compared to the other studied trees, these plants can be trap plants. These results are expected to help manage *O. japonica* by plants that would attract adults and prevent its dispersion.

Acknowledgments

We appreciate Dr. A. Karimi Malati, Department Of Plant Protection, Faculty of Agricultural Sciences, University of Guilan, for her assistance in sampling and valuable guidance. We would like to thank Mr. M. Mohammadi, Guilan Plant Protection Management, Organization of Agriculture Guilan, for his efficient aid in surveying *O. japonica* occurrence.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

References

- Abbaszadeh, S., Karimi, A and Jalali Sendi, J. 2021. Semi-field demographic performance of the invasive planthopper, *Orosanga japonica* (Hemiptera: Ricaniidae) on two host plants. *Journal of Asia-Pacific Entomology*, 24(3): 614-621.
- Ak, K., Güçlü, Ş., Eken, C. and Sekban, R., 2015. Türkiye için yeni bir zararlı *Ricania simulans* (Walker, 1851) (Hemiptera: Ricaniidae). *Türkiye Entomoloji Dergisi*, 39(2): 179-186.
- Akner, M. M., Beriş, F. Ş., Seyis, F., Öztürk, M., Sevgili, H. and Demir, E. 2019. Annual variation of the *Orosanga japonica* Melichar 1898 (Hemiptera: Ricaniidae) populations in the eastern Black Sea region of Turkey and possible molecular separation with based on 28S rDNA sequences from other Ricaniidae groups. *Plant Protection Bulletin*, 59(4): 11-19.
- Altas, K. and Ak, K. 2019. Effect of cultural management methods against fake butterfly [*Ricania japonica* (Hemiptera: Ricaniidae)]. *Artvin Çoruh Üniversitesi Orman Fakültesi Dergisi*, 20(2): 229-238. DOI: 10.17474/artvinofd.583374.
- Bourgoin, T. 2017. FLOW (Fulgoromorpha Lists on TheWeb): a world knowledge base dedicated to Fulgoromorpha. Version 8, updated [06.02.2017]. Accessed March 2017. <http://hemiptera-databases.org/flow/>.
- Bourgoin, T. 2021. FLOW (Fulgoromorpha Lists On the Web): a world knowledge base dedicated to Fulgoromorpha. Version 8, updated 2020-09-17. <http://hemiptera-databases.org/flow/> [Last access 2020-09-21].
- Choi, D. S., Kim, D. I., Ko, S. J., Kang, B. R., Lee, K. S., Park, J. D. and Choi, K. J. 2012. Occurrence ecology of *Ricania* sp. (Hemiptera: Ricaniidae) and selection of environmental friendly agricultural materials for control. *Korean Journal of Applied Entomology*, 51: 141-148.
- Choi, Y. S., Hwang, I. S., Kang, T. J., Lm, J. R. and Choe, K. R. 2011. Oviposition characteristics of *Ricania* sp. (Homoptera: Ricaniidae), a new fruit pest. *Korean Journal of Applied Entomology*, 50(4): 367-372.
- Demir, E. 2009. *Ricania* Germar, 1818 species of western Palaearctic region (Hemiptera: Fulgoromorpha). *Munis Entomology & Zoology Journal*, 4 (1): 271-275.
- Demir, E. 2018. The economically important alien invasive planthoppers in Turkey (Hemiptera: Fulgoromorpha). *Acta Entomologica Slovenica*, 26(2): 231-240.
- Dlabola, J. 1967. Beschreibungen von neuen su"dpala" arktischen Zikadenarten (Homoptera, Auchenorrhyncha). *Acta Entomologica Musei Nationalis Pragae*, 37: 31-50.
- Eken, C., Ak, K., Güçlü, S., Genç, T. and Sekban, R. 2013. Fungal flora of *Ricania simulans* (Hemiptera: Ricaniidae). XI. National Ecology and Environment Congress (01-04 September 2013, Samsun, Turkey), 208 pp.
- EPPO. 2017. *Ricania japonica*: a new polyphagous insect found in the EPPO region. EPPO Global Database. Accessed March 2017, <https://gd.eppo.int/reporting/article-5734>.
- Gjonov, I. 2011. *Ricania japonica* Melichar, 1898—a representative of family Ricaniidae

- (Homoptera, Fulgoromorpha), new to the fauna of Bulgaria. *ZooNotes*, 23: 1-3.
- Gokturk, T. and Aksu, Y. 2014. Morphology, biology and damage of *Ricania simulans* (Walker, 1851) (Hemiptera: Ricaniidae), which harms agricultural and forest areas. Turkey II. Forest Entomology and Pathology Symposium (7-9 April 2014), Antalya. 279-281. (Turkish).
- Ismaylova, G. 2021. New record of *Orosanga japonica* (Melichar, 1898) (Hemiptera: Fulgoromorpha: Ricaniidae) from Azerbaijan. *Punjab University Journal of Zoology*, 36(2): 141-145.
- Karatas, A., Karatas, A., Yavuz, Y. and Genc, M. 2020. Distribution and activity period of the invasive *Orosanga japonica* (Melichar, 1898) (Hemiptera: Ricaniidae) in Turkey. *Zoology in the Middle East*, 66: 246-252.
- Kim, D. H., Kim, H. H., Yang, C. Y., Kang, T. J. Yoon J. B. and Seo, M. H. 2016. Characteristic of Oviposition and Effect of Density Suppression by Yellow-colored Sticky Trap on *Ricania shantungensis* (Hemiptera: Ricaniidae) in Blueberry. *Korean Journal of Pesticide Science*, 20(4): 281-285.
- Lucchi, A. and Rossi, E. 2016. The Egg-Burster in the Asian Planthopper *Ricania speculum* (Walker) (Hemiptera Ricaniidae) *Annals of the Entomological Society of America*, 109(1): 121-126, <https://doi.org/10.1093/aesa/sav108>.
- Mozaffarian, F. 2018. An identification key to the species of Auchenorrhyncha of Iranian fauna recorded as pests in orchards and a review on the pest status of the species. *Zootaxa*, 4420(4):475-501, <https://doi.org/10.11646/zootaxa.4420.4.2>.
- Nast, J. 1972. Palaeartic Auchenorrhyncha (Homoptera). An annotated check list. Polish Scientific Publishers, Warszawa. 550 pp.
- Yu, F. G., Guo, L. N., Qun, Y., Min, L. X., Long, C. W. 2016. Behavior rhythm of *Ricania cacaonis*. *Guizhou Agricultural Sciences*, 44(4): 53-56.

پراکنش و ویژگی‌های تخم‌گذاری (*Orosanga japonica* Melichar (Hemiptera: Ricaniidae) در استان گیلان

سیده الهام یزدانی‌بدابی و احد صحراگرد

گروه گیاه‌پزشکی، دانشکده علوم کشاورزی، دانشگاه گیلان، رشت، ایران.
پست الکترونیکی نویسنده مسئول مکاتبه: elyazd2020@gmail.com
دریافت: ۸ اسفند ۱۴۰۰؛ پذیرش: ۸ شهریور ۱۴۰۲

چکیده: این مطالعه به منظور تعیین ویژگی‌های تخم‌گذاری، پراکنش و میزبان‌های زنجبرک (*Orosanga japonica* (Hemiptera: Ricaniidae) در استان گیلان انجام شد. از مناطق مختلف استان گیلان به صورت تصادفی نمونه‌برداری انجام شد. بالغین در اواخر خرداد ظاهر شده و در اوایل مرداد تا اواسط شهریور تخم‌های خود را می‌گذارند. این‌گونه روی شاخه‌های جوان، رگبرگ‌های میانی گیاهان مختلف شامل گیاهان علفی یکساله و درختچه‌ها تخم‌گذاری می‌کند که موجب خشکیدگی شاخه‌ها می‌شود. بنابراین رفتار تخم‌گذاری ماده‌ها و نیز مکیدن شیره ساقه‌ها و انتقال قارچ‌های بیماری‌زا موجب آسیب جدی به گیاهان میزبان می‌شود. این تحقیق جزئیاتی از توده‌های تخم روی چند گیاه میزبان را فراهم می‌کند. نتایج نشان داد که میانگین تعداد توده تخم در هر شاخه زیتون تلخ ۱۳/۷۵ بوده که نسبت به نارنج، نارون و توت بالاتر می‌باشد. بیشترین طول توده تخم ۱۶/۱ میلی‌متر روی توت بود. حضور *O. japonica* در ۱۱ شهر از استان گیلان شامل رشت، انزلی، املش، آستانه، لاهیجان، لنگرود، رضوان‌شهر، رودسر، رودبار، صومعه‌سرا، سیاهکل، شفت و فومن تأیید شد و بیشتر روی کیوی، چای، لوبیا، ذرت، بادمجان، فلفل، زیتون تلخ، تمشک، توت، شمشاد زینتی، خیار، مرکبات، انجیر، برنج و انگور مشاهده شد.

واژگان کلیدی: *Orosanga japonica*، Fulgoromorpha، توده تخم، پراکنش، ویژگی‌های تخم‌گذاری، گیلان