

## Research Article

# Application of chicken manure and summer plowing to control root-knot nematode *Meloidogyne javanica* in muskmelon, *Cucumis melo* var. *inodorus*, farms

Hadi Karimipour Fard\* and Nazanin Doryanizadeh

Plant Protection Research Department, Kohgiluyeh and Boyer-Ahmad Agricultural and Natural Resources Research and Education Center, AREEO, Yasouj, Iran.

**Abstract:** Root-knot nematode *Meloidogyne javanica* is an economically significant plant parasitic nematode in muskmelon farms in Ardestan, Isfahan province, Iran. A two-year experiment was conducted in two muskmelon farms in this region. We assessed the application of 3, 6, and 9 t/ha fresh (unrotten) chicken manure separately or in combination with summer plowing. To evaluate the efficacy of each treatment, the number of galls, egg masses, reproduction factor, plant growth factors, and yield were recorded at harvesting time. The results showed that combining double summer plowing with nine t/ha chicken manure was the most effective treatment for controlling *M. javanica*. Nematode population indices including: the egg mass number, gall number, total population in root and soil, and reproduction factor, were significantly lower than the control treatment. The application of double plowing with nine t/ha chicken manure treatment caused 83.5%, 79.1%, 80.2%, and 78.3% decreases in egg mass number, gall number, total population per root, and soil, and reproduction factor, respectively. Moreover, total fruit weight and root dry weight in each plot (24 m<sup>2</sup>) increased by 81.8% and 50.4% compared to the control.

**Keywords:** chicken manure, plowing, muskmelon, *Meloidogyne javanica*

## Introduction

Among plant pathogens, plant parasitic nematodes are one of the most devastating agents causing crop damage, and their annual damage is estimated at 100 billion dollars in various crops (Oka *et al.*, 2000). The root-knot nematode *Meloidogyne javanica* (Treub 1885) Chitwood, 1949 parasitizes a wide range of plants in tropical and temperate regions by penetrating plant roots and producing galls (Karssen *et al.*, 2013). The root-knot nematode

reduces photosynthesis and growth and causes nutrient deficiency, yellowing and wilting. It also increases the susceptibility of the root system to invasive fungal/bacterial plant pathogens. The damage to young plants can result in death and severe infection, often preventing the plants from growing to maturity (Jaffee and Muldoon, 1995).

*Meloidogyne* spp. is an important pathogen in vegetable and summer crops (Moens *et al.*, 2009; Collange *et al.*, 2011). In general chemical pesticides, including nematicides, are

Handling Editor: Zahra Tanha Maafi

\* Corresponding author: karimipourfard@yahoo.com

Received: 28 February 2022, Accepted: 14 January 2023

Published online: 19 January 2023

widely used as a control method (Javed *et al.*, 2007). However, due to environmental considerations, chemical control of plant parasitic nematodes has become less desirable (Castillo *et al.*, 2010) and very much restricted in developed countries. Therefore, relying on safe methods such as resistant cultivars, organic materials, and cultivation practices is more emphasized (Widmer *et al.*, 2002). Twice summer plowing and postharvest summer plowing in beet fields resulted in a greater reduction of root-knot nematode (Akhyani *et al.*, 1984; Savage and Bramwell, 2002). Organic amendments such as chicken manure affect soil properties and nematode population (Akhtar and Malik, 2000; Litterick *et al.*, 2004; Nahar *et al.*, 2006; Hu and Qi, 2010; Oka, 2010; McSorley, 2011). They act by stimulating many potential predators or parasites of plant-parasitic nematodes in the soil (Akhtar and Malik, 2000; Oka, 2010). Also, they can enhance nematode-trapping fungi and egg parasitism (Mukerji and Garg, 1988; Singh and Sitaramaiah, 1973). Using chicken manure also improves plant growth and increases crop yield (El-Sherief *et al.*, 2010; Abolusoro *et al.*, 2012). The impact of organic manure depends on many factors, such as organic material type, composting of material, application rate, crop rotation, agronomic practices, soil type, climate, and other environmental factors (McSorley, 2011). Chicken manure is a rich source of organic matter, which, in addition to improving soil conditions, contains nutrients that are effective in plant nutrition. It is especially rich in nitrogen content compared to other animal manures (Hirzell and Walter, 2008). The efficacy of chicken and poultry manure on the reduction of root-knot nematodes has been reported in many types of research (Nasr-Esfahani, 2002; Antonio *et al.*, 2005; Karimipour Fard *et al.*, 2019). Osei *et al.* (2011) reported that chicken manure reduces the root-knot nematode population density by inhibiting egg hatching.

This study evaluated the efficacy of chicken manure and summer plowing for managing *M.*

*javanica* in muskmelon farms. The aims of the study were: *i*: investigating the effect of different levels of chicken manure and summer plowing on root-knot nematode, *ii*: combining the summer plowing and chicken manure in reducing the nematode indices, *iii*: improving the muskmelon yield and growth indices in out-of-season muskmelon cultivation.

## Materials and Methods

The experiment was conducted on two muskmelon fields with a history of root-knot nematode infestation in different regions of Ardestan in Isfahan province, Iran.

### Soil samples

Five randomized core soil samples were taken from each plot (6 × 4 m) to assay the initial nematode population (Pi). The soil samples were collected from experimental fields at the beginning of the first week of July 2009 in a systematic zigzag method, from 0-30 cm depth. Nematodes were extracted by a modified centrifugal floatation method (Jenkins, 1964).

**Application of chicken manure:** The selected plots were tilled twice in July and August 2009. Unrotten chicken manure was applied two months before planting in plots based on the calculated amount. The trials were conducted in a randomized complete block design with eight treatments viz: double plowing, three tonnes chicken manure/ha, double plowing + three tonnes chicken manure/ha, six tonnes chicken manure/ha, double plowing + six tonnes chicken manure/ha, nine tonnes chicken manure/ha, and double plowing + nine tonnes chicken manure/ha, and control without applying plowing and chicken manure with four replications.

**Planting muskmelon:** Muskmelon seeds were sown at the end of the first week of February 2010 in two rows at 3-4 cm depth and 50 cm space between the rows and 70 cm apart along the row. In this winter cropping (out-of-season cultivation), the ridges were covered by plastic tunnel.

**Qualitative and quantitative parameters determination:** Plant height, fresh and dry weight (75 °C in the oven for 48 h) of the aerial shoot (whole aerial shoot), and root (whole root system) were measured. During the harvest season (last week of May 2010 to Mid-June 2010), fruit weight was recorded. From the beginning of the harvest time, the yield of each plot (24 m<sup>2</sup>) was weighed separately. The total weight of the fruit was calculated through the sum of the total yield in consecutive harvests. After 100 days of planting (last week of May 2010), two plants from each plot were unrooted, and the number of egg masses and galls per root system was determined (Taylor and Sasser, 1978). The nematode developmental stages/cm root was determined by staining roots using acid fuchsin solution and Lacto glycerol (Bridge *et al.*, 1982). The number of second-stage juveniles (J2s) was estimated by extracting nematodes from 200 cm<sup>3</sup> of soil samples of each plot using a modified centrifugal floatation method (Jenkins, 1964). The sum of eggs, juveniles, and adult populations of nematodes in the roots and the population of J2s in 200 cm<sup>3</sup> of soil per plot were considered the final population of nematodes in each plot. The Nematode reproductive factor (Rf) was also calculated based on initial and final population density.

**Soil analysis:** Before and after applying the chicken manure, some soil properties, including pH (by pH meter), organic carbon (by Walkley-Black method, Nelson and Sommers, 1996), available P (Olsen's method), and available K (by flame photometer) (Knudsen *et al.*, 1982) were measured.

## Results

The identification of *M. javanica* was confirmed by morphological and morphometric studies. The results showed that double plowing plus nine tonnes of chicken manure/ha was the best treatment for control of the root-knot nematode population in muskmelon fields. This treatment significantly reduced root-knot

nematode population and increased the plant growth characteristics. The total population per root, soil and reproduction factor (RF) was reduced compared to other treatments (Table 1). It caused 83.5%, 79.1%, 80.2%, and 78.3% decreases in egg mass number, gall number, total population per root, and soil and reproduction factor, respectively. This treatment also resulted in significantly higher aerial shoot dry weight, plant and root dry weight by 81.8% and 50.4% increase in the factors of total fruit weight and root dry weight per plot (24 m<sup>2</sup>), respectively. The total fruit weight was also significantly higher than the control (Table 1).

Analysis of soil samples before and after applying chicken manure for two years showed that there was a significant increase in soil salinity (electrical conductivity), organic matter, and the absorbable content of phosphorus and potassium and a significant decrease in pH ( $p \leq 0.05$ ; Figure 1). This study also showed that using chicken manure at the rate of 9t/ha reduced soil pH by 0.31 (Fig. 1), which impacted the improvement of soil conditions and absorption of micronutrients.

## Discussion


In our investigation, the lowest population density of root-knot nematode and the most quantitative characters in fruit yield and plant growth parameters belonged to the treatment of double plowing plus nine tonnes of chicken manure. The application of this treatment led to an increase in plant growth indices and a decrease of 78.3% and 81.8% in the overall reproduction factor and total population of nematodes in soil and roots, respectively.

Our results were in agreement with the results of the works done on this subject, which confirm the effectiveness of chicken manure for root-knot nematode control of different crops (Nasr-Esfahani, 2002; Lopez-Pérez *et al.*, 2005; Oka *et al.*, 2000; El-Sherief *et al.* 2010; McSorley, 2011; Abolusoro *et al.*, 2012; Karimipour Fard *et al.*, 2019).

**Table 1** The effectiveness of chicken manure, plowing, and combined treatments on nematode indices and muskmelon growth parameters in muskmelon fields.

Treatments	No. of Egg /whole root	No. of gall /whole root	Reproduction factor	Total population (in root & soil)	Aerial shoot fresh weight (g)	Aerial shoot dry weight (g)	Plant height (cm)	Root dry weight (g)	Root length (cm)	Total fruit weight (Kg)
Double ploughing+ 3 tonnes of chicken manure	18.62 c	7.87 d	6.14 bc	11516.1 e	1221.25 d	11.60e	112.87 a	3.06 ab	23.92 b	28.50 a
3 tonnes of chicken manure	35.87 b	13.12 bc	7.22 b	15150.6 c	1516.25 c	16.86 d	114.25 a	2.42 ab	22.90 b	19.75 b
Double ploughing + 9 tonnes of chicken manure	17.87 c	6.87 e	4.91 c	8858.4 f	1723.75 b	22.05 a	117.25 a	3.52 a	31.83 a	31.25 a
9 tonnes of chicken manure	30.75 b	12.12 bc	6.17 bc	13641.5 d	2246.25 a	19.91 b	121.62 a	2.83 ab	25.45 ab	27.87 a
Double plowing	18.50 c	11.37 bc	6.77 bc	14337.3 cd	1285.00 d	19.37 bc	113.50 a	3.05 ab	27.39 ab	27.75 a
6 tonnes of chicken manure	29.50 bc	14.12 b	7.83 b	17252.8 b	2173.75 a	18.60 c	110.25 a	3.32 ab	23.97 b	30.06 ab
Double ploughing + 6 tonnes of chicken manure	20.00 cd	10.37 cd	5.28 b	12213.8 e	1586.25 bc	19.57 bc	109.62 a	2.38 ab	22.44 b	30.56 a
Control (no plowing and no chicken manure)	108.50 a	33.00 a	22.67 a	42260.0 a	846.25 e	17.52 d	89.12 b	2.16 b	22.60 b	17.18 b

Note: Values in the same column followed by the same letter(s) are not significantly different ( $p \leq 0.05$ ) using Duncan's Multiple Range test. From each plot of 24 m<sup>2</sup>, two plants with roots were used to measure nematode and plant indices. Each treatment included four replicates; therefore, the values are the average of eight plants.



**Figure 1** Comparing the soil properties before and after nine tonnes of chicken manure application.

In this study, unlike most previous studies, an unrotten and pure form of chicken manure from laying hens was used in a much lower rate than the amounts used in previous studies. Some studies suggest that chicken manure stimulates the decomposition of nematode cuticular layers (Rodriguez-Kabana, 1986; Stirling, 1991). Chicken manure releases toxic

levels of ammonium and probably changes the soil structure during decomposition. It also causes stimulation of antagonistic organisms, increases plant tolerance, and promotes nematode control (Oka *et al.*, 2000; Lopez-Pérez *et al.*, 2005).

Our findings are similar to the results of Everts *et al.* (2006) who indicated that summer

plowing can be considered an effective method to control *M. incognita* in potato fields.

Besides the efficacy of chicken manure for managing root-knot nematodes in muskmelon farms, it showed improved soil fertility through increased organic carbon, P, and K contents in the treated soils.

Cultural practices like amending soil with chicken manure and plowing are environment-friendly management tactics (Oka, 2010). Applying these methods prevents the hazards of chemical nematicides on the environment and on human health (Oka, 2010). Using chicken manure combined with summer plowing can be considered a practical management strategy that could be an alternative to chemical control for the root-knot nematode. Meanwhile, the costs of nematicide application and its effects on non-target species should be considered (Udo and Ugwuoke, 2010).

Use of chicken manure reduced soil pH and raised soil salinity, P, and K of the soil in the treatment of double plowing plus nine tonnes of chicken manure; however, applying pure chicken manure in this study reduced the amount of manure and also prevented high raising of soil salinity as a consequence of using high amounts of fresh (unrotten) chicken manure. Comparison of the salinity (EC) of the soil before and after application of 9 t/ha of chicken manure showed a slight rise of about  $0.07 \text{ ds.m}^{-1}$  in the soil. Although the tolerable salinity levels vary for different crops, this amount for muskmelon is up to  $2.5 \text{ ds.m}^{-1}$  while  $2 \text{ ds.m}^{-1}$  is ideal. However, the soil salinity of  $2 \text{ ds.m}^{-1}$  does not affect yield reduction, but 5% and 20% yield loss occurs at 3 and  $4 \text{ ds.m}^{-1}$ , respectively (Mass, 1986). Using 9 t/ha chicken manure increased the soil organic carbon by 0.92% and significantly increased absorbable potassium and phosphorus. This process, in addition to control of root-knot nematode, will improve the soil conditions and plant growth.

Adding any decomposable organic matter to the soil leads to changes in the physicochemical and biological properties of the soil. It also provides suitable conditions for the activity of antagonists and the establishment of biological control of soil-borne pathogens (Westphal *et al.*,

2016). The use of organic compounds, such as chicken manure that is rich in nitrogen content due to the release and dispersion of ammonia after microbial degradation, causes a rapid reduction in the population of plant parasitic nematodes on the nematode population (Lazarovits *et al.*, 2001). We used fresh (unrotten) chicken manure in this study because of its high fumigant activity (Zhang *et al.*, 2021). Also, it is readily available and is cheaper than chemical fumigants with fewer environmental hazards.

## Conclusion

This research showed that using chicken manure helped to control root-knot nematodes and increased muskmelon yield. In the tropical regions of Iran, summer crops cultivated out-of-season is done. Usually, after harvesting these crops in the summer, the farm is left uncultivated due to high temperatures and water shortages, which can be subjected to summer plowing. Using chicken manure in the recommended amount in these farms would provide a safe and environmentally friendly method for controlling root-knot nematodes. Therefore, according to the conditions and facilities, growers can manage muskmelon root-knot nematode by using chicken manure alone or in combination with summer plowing to enhance its effects.

## Acknowledgments

The Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran, provided funding for this research. We thank our colleagues; Hassan Almasi, Ali Farhadi, and Mojtaba Yahya Abadi; from Isfahan Agricultural and Natural Resources Research and Education Center, Iran, for their collaborations.

## References

- Abolusoro, S. A., Abolussoro, P. F. and Mathew, F. O. 2012. Effects of organic and inorganic manures on the growth attributes

- of root-knot nematode (*Meloidogyne incognita*) infected Ethiopian eggplant (*Solanum aethiopicum*). Libyan Agriculture Research Center Journal International, 3(5): 224-228.
- Akhtar, M. and Malik, A. 2000. Roles of organic soil amendments and soil organisms in the biological control of plant-parasitic nematodes: A review. Bioresource Technology, 74: 35-47.
- Akhyani A., Mojtahedi H. and Naderi A. 1984. Species and physiological races of root knot nematodes in Iran. Iranian Journal of Plant Pathology, 20: 57-70.
- Antonio, J., Perez, L., Roubtsova, T. and Ploeg, A. 2005. Effect of three plant residues and chicken manure used as Biofumigants at three temperature on *Meloidogyne incognita* infestation of Tomato in Greenhouse Experiments. Journal of Nematology, 37(4): 489-494.
- Bridge, J., PagGe S. and Jordan, S. 1982. An Improved Method for staining nematodes in roots. Rothamsted Experimental Station Report for 1981, Part 1.171.
- Castillo, P., Nico, A. I., Navas-Cortés, J. A., Landa B. B., Jiménez-Díaz, R. M. and Vovlas N. 2010. Plant-Parasitic Nematodes Attacking Olive Trees and their Management. Plant Disease, 94: 148-162.
- Collange, B., Navarrete, M., Peyre, G., Mateille, T. and Tchamitchian, M. 2011. Root-knot nematode (*Meloidogyne*) management in vegetable crop production: The challenge of an agronomic system analysis. Crop protection, 30(10): 1251-1262.
- El-Sherief, A. G., Refaei, A. R. and Gad, S. B. 2010. Effects of certain animal manures or plant products alone or integrated with oxamyl on growth of peanut plant infected with *Meloidogyne javanica*. Egyptian Journal of Agronomatology, 9(1): 30-39.
- Everts, K. L., Sardenelli, R., Kratochvil, R. J., Armentrout, D. K. and Gallagher L. E. 2006. Root-knot and Root-lesion nematode suppression by Cover crops, Poultry litter and Poultry litter compost. Plant Disease, 90(4): 487-492.
- Hirzell, J. and Walter, I. 2008. Availability of nitrogen, phosphorus and potassium from poultry litter and conventional fertilizers in a volcanic soil cultivated with silage corn. Chilean Journal of Agricultural Research, 68(3): 264-273.
- Hu, C. and Qi, Y. C. 2010. Abundance and diversity of soil nematodes as influenced by different types of organic manure. Helminthologia, 47(1): 58-66.
- Jaffee, B. A. and Muldoon, A. E. 1995. Susceptibility of root-knot and cyst nematodes to the nematode-trapping fungi *Monacrosporium ellipsosporum* and *M. cionopagum*. Soil Biology & Biochemistry, 27(8): 1083-1090.
- Javed, N., Gowen, S. R., Inam-ul-Haq, M., Abdullah, K. and Fayyaz, S. 2007. Systemic and persistent effect of neem (*Azadirachta indica*) against root knot nematode, *Meloidogyne javanica* and their storage life. Crop Protection, 26(7): 911-916.
- Jenkins, W. R. 1964. A rapid centrifugal flotation for separating nematodes from soil. Plant Disease Reporter, 48(9): 692-692.
- Karimipour Fard, H., Saeidi, K. and Doryanizadeh, N. 2019. Effects of chicken manure and summer ploughing on root-knot nematode (*Meloidogyne javanica*) management in cantaloupe (*Cucumis melo* var. *cantalupensis*). Archives of Phytopathology and Plant Protection, 52(15-16): 1193-1205.
- Karssen, G., Wesemael, W. and Moens, M. 2013. Root-knot nematodes. In: Perry R. N. and Moens, M. (Eds). Plant nematology, 2<sup>nd</sup> edition. CABI, Wallingford, UK. pp. 73-108.
- Knudsen, D., Peterson, G. A., Pratt, P. F. 1982. Lithium, Sodium and Potassium Methods of Soil Analysis, Part 2 Chemical and Microbiological Properties. Agronomy Monograph No: 9 ASA-SSSA. Wisconsin, USA.
- Lazarovits, G., Tenuta, M. and Conn, K. L. 2001. Organic amendments as a disease control strategy for soil borne disease of high-value agricultural crops. Australasian Plant Pathology, 30(2): 111-117.
- Litterick, A. M., Harrier, L., Wallace, P., Watson, C. A. and Wood, M. 2004. The role of

- uncomposted materials, composts, manures, and compost extracts in reducing pest and disease incidence and severity in sustainable temperate agricultural and horticultural crop production—A review. *Critical Reviews in Plant Sciences*, 23(6): 453-479.
- López-Pérez, J. A., Roubtsova, T. and Ploeg, A. 2005. Effect of three plant residues and chicken manure used as biofumigants at three temperatures on *Meloidogyne incognita* infestation of tomato in greenhouse experiments. *Journal of Nematology*, 37(4): 489-494.
- Mass, E. V. 1986. Salt tolerance in plants. *Applied Agricultural Research*, 1: 12-26.
- McSorley, R. 2011. Overview of organic amendments for management of plant-parasitic nematodes, with case studies from Florida. *Journal of Nematology*, 43(2): 69.
- Moens, M., Perry, R. N. and Starr, J. L. 2009. *Meloidogyne* species: A diverse group of novel and important plant parasites. In: Perry, R. N, Moens, M. and Starr, J. L. (Eds.) *Root-Knot Nematodes*. CABI Publishing, Wallingford. pp. 1-17.
- Mukeji, K. G. and Garg, K. L. 1988. *Biocontrol of Plant Disease (Vol.1)*. CRC press. Florida, USA.
- Nahar, M. S., Grewal, P. S., Miller, S. A., Stinner, D., Stinner, B. R., Kleinhenz M. D., Wszelaki, A. and Doohan, D. 2006. Differential effects of raw and composted manure on nematode community and its indicative value for soil microbial, physical and chemical properties. *Applied Soil Ecology*, 34(2-3): 140-151.
- Nasr-Esfahani, M. 2002. Evaluation of non-chemical control of root-knot nematode of cucumber. Final report. Plant pests and diseases research institute, Agricultural Research Center of Isfahan. 19 p. (in Persian).
- Nelson, D. W. and Sommers, L. E. 1996. Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 3. Chemical Methods*, 5: 961-1010.
- Oka, Y. 2010. Mechanisms of nematode suppression by organic soil-amendments —a review. *Applied Soil Ecology*, 44(2): 101-115.
- Oka, Y., Koltai, H., Bar-Eyal, M., Mor, M., Sharon, E., Chet, I. and Spiegel, Y. (2000). New strategies for the control of plant parasitic nematodes. *Pest Management Science*, 56(11): 983-988.
- Osei, K., Addico, R., Nafeo, A., Edu-Kwarteng, A., Agyemang, A., Danso, Y. and Sackey-Asante, J. 2011. Effect of some organic waste extracts on hatching of *Meloidogyne incognita* eggs. *African Journal of Agricultural Research*, 6(10): 2255-2259.
- Rodriguez-Kabana, R. 1986. Organic and inorganic nitrogen amendments to soil as nematode suppressants. *Journal of Nematology*, 18(2): 129.
- Savage, I. and Bramwell, R. K. 2002. Root diseases, weeds, and nematodes with poultry litter and conservation tillage in a sweet corn–snap bean double crop. *Crop Protection*, 21(10): 963-972.
- Singh, R. S. and Sitaramaiah, K. 1973. Control of Plant Parasitic Nematodes with Organic amendments of soil. GB Plant University of Agriculture and Technology, India.
- Stirling, G. R. 1991. *Biological control of Plant Parasitic Nematodes*. CAB International. Redwood Press Ltd, Melksham.
- Udo, A. I. and Ugwuoke, K. I. 2010. Pathogenicity of *Meloidogyne incognita* Race 1 on Tumeric as Influenced by Inoculum Density and Poultry Manure Amendment. *Plant Pathology Journal*, 9(4): 162-168.
- Westphal, A., Kücke, M. and Heuer, H. 2016. Soil amendment with digested from bio-energy fermenters for mitigating damage to *Beta vulgaris* sub spp. by *Heterodera schachtii*. *Applied Soil Ecology*, 99: 129-136.
- Widmer, T. L., Mitkowski, N. A. and Abawi, G. S. 2002. Soil organic matter and management of plant-parasitic nematodes. *Journal of Nematology*, 34(4): 289-295.
- Zhang, D., Cheng, H., Hao, B., Li, Q., Fang, W., Ren, L., Yan, D., Ouyang, C., Li, Y., Wang, Q. and Jin, X., 2021. Effect of fresh chicken manure as a non-chemical soil fumigant on soil-borne pathogens, plant growth and strawberry fruit profitability. *Crop Protection*, 146: 105653.

## کاربرد کود مرغی و شخم تابستانه به منظور کنترل نماتد ریشه گری *Meloidogyne javanica* در مزارع خربزه

هادی کریمی‌پور فرد\* و نازنین دریانی‌زاده

بخش تحقیقات گیاه‌پزشکی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان کهگیلویه و بویراحمد، سازمان تحقیقات، آموزش و ترویج کشاورزی، یاسوج، ایران. پست الکترونیکی نویسنده مسئول مکاتبه: karimipourfard@yahoo.com دریافت: ۹ اسفند ۱۴۰۰؛ پذیرش: ۲۴ دی ۱۴۰۱

**چکیده:** نماتد ریشه‌گری *Meloidogyne javanica* یکی از نماتدهای انگل گیاهی مهم از لحاظ اقتصادی در مزارع خربزه اردستان در استان اصفهان است. آزمایشی دو ساله در دو مزرعه خربزه آلوده به این نماتد در منطقه مذکور، با کاربرد کود مرغی در مقادیر سه، شش و نه تن در هکتار به‌طور جداگانه یا همراه با شخم تابستانی جهت کنترل نماتد ریشه گری، انجام شد. برای ارزیابی تأثیر هر یک از تیمارها روی *M. javanica*، تعداد گال، توده تخم، فاکتور تولیدمثلی، شاخص‌های مربوط به رشد گیاه و همچنین عملکرد محصول در زمان برداشت ثبت شد. نتایج نشان داد که ترکیب دو بار شخم تابستانه با نه تن کود مرغی نپوسیده، مؤثرترین تیمار برای کنترل *M. javanica* است. شاخص‌های جمعیتی نماتد شامل تعداد توده تخم، تعداد گال، جمعیت کل در ریشه و خاک و فاکتور تولیدمثلی در این تیمار به‌طور معنی‌داری کمتر از تیمار شاهد (بدون کود مرغی و شخم تابستانه) بود. اعمال تیمار مذکور به‌ترتیب باعث کاهش ۸۳/۵، ۷۹/۱، ۸۰/۲ و ۷۸/۳ درصد در تعداد توده تخم، تعداد گال، کل جمعیت نماتد در ریشه و خاک و فاکتور تولیدمثلی شد. همچنین وزن کل میوه و وزن خشک ریشه در هر کرت (به ابعاد ۲۴ مترمربع) نسبت به شاهد ۸۱/۸ درصد و ۵۰/۴ درصد افزایش یافت.

**واژگان کلیدی:** کود مرغی، شخم زدن، نماتد ریشه‌گری، خربزه، *Meloidogyne javanica*