

Research Article

First report of the genus *Hoplotylus* s'Jacob, 1960 (Nematoda: Hoplolaimidae) and two species of the genus *Malenchus* Andrássy, 1968 (Nematoda: Tylenchidae) from Iran

Parisa Jalalinasab and Ramin Heydari*

Department of Plant Protection, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran.

Abstract: In order to identify plant-parasitic nematodes of eastern forests in Guilan province, soil and root samples were collected and the extracted nematodes were studied based on morphological and morphometric characters. Twenty eight species of plant parasitic nematodes were identified. A population of the genus *Hoplotylus* was recovered and identified as *H. femina*. The identification was also verified by partial sequencing of the large subunit ribosomal DNA (rDNA). In addition, two populations of the genus *Malenchus* were also recovered and based on morphological and morphometric characteristics were identified as *M. solovjovae* and *M. acarayensis*. The two species *Hoplotylus femina* and *Malenchus solovjovae* are new records for Iran.

Keywords: Hoplotylus femina, Malenchus solovjovae, M. acarayensis, large subunit ribosomal DNA

Introduction

Guilan province is one of the northern provinces of Iran with a subtropical climate and very heavy rainfall. Large parts of the province are mountainous, and covered with natural forests. So far, several nematode species have been reported from the province (Alem et al., 2010; Aliramaji et al., 2012; Divsalar et al., 2011; Eskandari et al., 2009; Nassaj Hosseini et al., 2004; Pedramfar et al, 2001; Shahabi et al., Soleymanzadeh et al., 2016; Tanha Maafi et al., 1998, 2002, 2006). In a recent attempt to identify the plant parasitic nematodes of forests in eastern part of the province, several species were recovered and identified. Two species namely Hoplotylus femina s'Jacob,

Handling Editor: Ebrahim Pourjam

*Corresponding author, e-mail: rheydari@ut.ac.ir Received: 18 February 2017, Accepted: 19 December 2017 Published online: 9 January 2018 1960 and *Malenchus solovjovae* Brzeski, 1989 are new records from Iran and the species *Malenchus acarayensis* Andrássy, 1968, previously reported from Iran (Namadipour *et al.*, 2014), are all illustrated herein.

The genus *Hoplotylus* was originally described by s'Jacob (1960) from the surrounding soil of the roots of *Quercus robur* L. and Chamaecyparis lawsoniana Parl in Wageningen, The Netherlands. Species of the genus Hoplotylus are endoparasitic plant nematodes. Based on head shape and mode of parasitism, Hoplotylus is close to members of the family Pratylenchidae Thorne, 1949 and based on stylet shape and position of the phasmid (near to anus) it is similar to members of the family Hoplolaimidae Filipjev, 1934. In the taxonomic framework of Siddiqi (2000), the genus *Hoplotylus* was included Pratylenchidae under the subfamily Radopholinae Allen & Sher, 1967 and was distinguished from other members of the

subfamily by females with stout and rather long stylet, having rounded to tulip-shaped knobs, males with developed bursa not enveloping tail and high and asymmetrical head. A recent study shows close relationship of this genus with members of the family Hoplolaimidae (Palomares-Rius *et al.*, 2016).

The genus *Malenchus* was originally described by Andrássy (1968) with M. machadoi as the type species. Species of the Malenchus are characterized prominent body annuli. In some species, lateral fields have two incisures under light microscope which, in scanning electron microscopy (SEM), shows very longitudinal lines and in some species, four or six incisures can be seen (Siddiqi, 2000). Historically, Andrássy (1981) described seven species and proposed Neomalenchus as a synonym of Malenchus. Sumenkova (1988) established Paramalenchus Sumenkova, 1988 for the species P. anthrisulcus Sumenkova, 1988 but, it was synonymised with Malenchus by Ebsary (1991). Siddiqi (2000) introduced subgenus Telomalenchus and also considered Neomalenchus and *Malenchus* as subgenera for this genus. According to Geraert (2008), the genus includes two subgenera, Malenchus Siddiqi, 2000 and Telomalenchus Siddiqi, 2000. Telomalenchus includes three unusual Malenchus species namely M. williamsi Geraert & Raski, 1986, M. parthenogeneticus Geraert & Raski, 1986 and M. leiodermis Geraert & Raski, 1986 which differ from the subgenus Malenchus in having wider lateral field with four or six incisures, distinct vulval lateral membrane and straight amphidial apertures (sinuate in subgenus Malenchus). Here we followed Geraert's (2008) scheme for classification of the recovered species of the genus Malenchus.

Materials and Methods

Soil and root samples were randomly collected from different regions of forests in the eastern parts of Guilan province during 2014-2015. The nematodes were recovered from soil samples using tray method (Whitehead and Hemming 1965) and the extracted nematodes were killed by gentle heat, fixed in a solution of TAF (8% formalin and 2% triethanolamine in distilled water) and then processed to anhydrous glycerin (De Grisse, 1969). Permanent slides were made and examined using an Olympus BH2 light microscope. Morphometric data were obtained and line drawings were made using a drawing tube attached to a Nikon E200 light microscope. Photomicrographs were taken using a digital camera attached to the same microscope.

The partial sequence of the D2/D3 expansion segment of 28S rRNA gene was obtained for the *Hoplotylus* population; total DNA was extracted using worm lysis buffer containing proteinase K (Williams *et al.*, 1992). The forward D2A (5'–ACAAGTACCGTGAG GGAAAGTTG–3') and reverse D3B (5'–TCGGAAGGAACCAGCTACTA–3') primers (Nunn, 1992) were used for amplification and sequencing of this region. Polymerase Chain Reaction (PCR) assays and sequencing of DNA products were performed according to Heydari *et al.* (2014).

Results

During this study, 28 plant parasitic nematode species belonging to 18 families were identified (Table 1). A population of the genus Hoplotylus was isolated and based morphological, morphometric molecular characteristics, identified as H. femina; being the first representative of the genus from Iran.. Two species of the genus Malenchus (subgenus Malenchus) i.e. M. solovjovae (as a new record) and M. acaravensis were also identified. complementary description was provided for the latter species, previously reported from Iran without description (Namadipour et al., 2014).

Table 1 List of nematode species identified from the forest in eastern Parts of Guilan province in the present study.

Species	Locality	Associated plants
Aphelenchus avenae	Langarud, Lahijan, Deylaman	Blackberry
Paraphelenchus myceliophthorus	Deylaman	Caucasian Wingnut
Aphelenchoides centralis	Lahijan	Caucasian elm
Aphelenchoides fuchsi	Deylaman	Caucasian Wingnut
Aphelenchoides parabicaudatus	Langarud	Moss
Aphelenchoides varicaudatus	Deylaman	Pine
Seinura hyrkania	Deylaman	Pine
Ditylenchus triformis	Deylaman	Elm
Nothotylenchus affinis	Deylaman	Caucasian Wingnut
Nothotylenchus medians	Deylaman	Caucasian elm
Helicotylenchus pseudodigonicus	Deylaman	Elm
Rotylenchus agnetis	Deylaman	Fern
Pratylenchus caffeae	Lahijan	Maple
Pratylenchus mediterraneus	Deylaman	Elm
Hoplotylus femina	Deylaman	Caucasian elm
Coslenchus costatus	Siahkal	Elm
Cephalenchus leptus	Deylaman, Siahkal, Langarud	Caucasian Wingnut, Caucasian elm, Willow, Black Locust, Fern
Boleodorus thylactus	Siahkal	Elm
Filenchus afghanicus	Lahijan, Deylaman	Maple, Caucasian Wingnut
Filenchus discrepans	Siahkal	Caucasian Wingnut, Elm
Filenchus misellus	Amlash	Beech
Filenchus thornei	Langarud	Beech
Malenchus acarayensis	Deylaman	Caucasian elm
Malenchus solovjovae	Deylaman	Pine
Deladenus durus	Deylaman	Caucasian elm
Paratylenchus lepidus	Siahkal	Walnut
Paratylenchus straeleni	Deylaman	Caucasian elm
Xiphinema hymenense	Deylaman	Caucasian elm

Iranian population of *Hoplotylus femina* s'Jacob, 1960

(Figs. 1 and 2; Table 2)

Description

Female: Body cylindrical. Lip region slightly set off from body contour with heavily sclerotized labial framework and three or occasionally four annuli. Lateral field with four incisures at mid-body, the outer lines areolated all along the body. Stylet well developed, basal tulip-shaped. Orifice knobs of dorsal pharyngeal gland at $6-7\mu m$ distance posterior to stylet base. Glandular region of pharynx overlapping intestine dorsally. Hemizonid at 1-2 annuli anterior to excretory pore. Reproductive system monodelphic, ovary outstretched with one row of oocytes. Spermatheca spherical, filled with sperm. Vulva a transverse slit, post-vulval uterine sac small. Tail conical, its tip pointed or finely rounded, having hyaline region at terminus with smooth or coarse irregularly annulated outline. Phasmids pore-like, four annuli posterior to anus.

Male: Body shorter and more slender than that of females. Sexual dimorphism present in anterior region, *i.e.* males have higher and asymmetrical head with three annuli. Stylet and pharynx reduced. Basal stylet knobs small and rounded. Testis single, outstretched. Spicules slightly curved ventrally. Bursa not enveloping tail, with phasmidial extension and three incisures in this region. Tail conical, with a pointed terminus.




Figure 1 Iranian population of *Hoplotylus femina* s'Jacob, 1960. A: Male head end; B: Female head end; C: Female entire body; D: Male entire body; E-G: Posterior region of female tail showing hyaline portion; H-J: Male posterior end showing spicules and caudal alae. (Scale bars: A, B, E-J = $10\mu m$; C, D = $20\mu m$).




Figure 2 Iranian population of *Hoplotylus femina* s'Jacob, 1960. A: Female head end; B: Male head end; C: Female pharyngeal region showing excretory pore; D: Female posterior end showing vulval region and post uterine sac (PUS); E: Lateral field at mid-body; F, G: Posterior region of female tail showing hyaline portion; H: Male posterior end showing spicules; I: Male caudal alae. (Scale bars = 10μm).

Table 2 Morphometrics of *Hoplotylus femina* from Iran and the data from other reports.

Characters	Guilan province		s'Jacob, 1960		Palomares-Rius et al., 2016	
	Female	Male	Female	Male	Female	Male
n	7	8	29	10	10	2
L	$527 \pm 68.6 (429-598)$	$421 \pm 38.3 (378-478)$	536 (450-656)	580	$611 \pm 60 (525-694)$	520 (480-560)
a	$26.9 \pm 2.1 (24.4-29.9)$	$30.6 \pm 3.8 (25.3 - 36.1)$	26.2 (23.1-30.6)	44	$28.3 \pm 4.2 (23.3-36.5)$	40.3 (32.0-48.7)
b	$5.8 \pm 0.6 (4.7 - 6.8)$	$5.2 \pm 0.6 (4.3 - 6.2)$	5.9 (5.1-7.5)	6.8	$6.2 \pm 0.6 (5.5 - 7.0)$	6.5 (5.9-7.1)
b'	$3.5 \pm 0.4 (2.9 - 4.0)$	$4.0 \pm 0.4 (3.4 - 4.6)$	-	5.8	$5.0 \pm 0.6 (4.5 \text{-} 6.0)$	5.1 (4.3-5.8)
c	$14.8 \pm 2.0 (13.2 - 18.6)$	$18.1 \pm 4.3 \ (10.6-21.3)$	16.1 (14.2-19.2)	13.5	$16.6 \pm 1.9 (12.5 - 19.0)$	15.1 (14.1-16.0)
c'	$3.0 \pm 0.3 (2.6 - 3.4)$	$2.3 \pm 0.6 (1.7-3.4)$	-	3.6	$2.7 \pm 0.2 (2.5 - 3.0)$	3.7 (3.2-4.1)
V or T	$85.2 \pm 1.7 (83.7-88.7)$	$38.5 \pm 9.8 (17.6 - 46.0)$	85.9 (84.0-87.3)	58	$86.2 \pm 1.7 (82.5 - 88.0)$	15.1 (10.1-20.0)
V'	$91.4 \pm 1.1 (90.4-93.7)$	-	-	-	-	-
Stylet	$23.6 \pm 1.5 (21-26)$	$13.3 \pm 0.5 (13.0-14.5)$	25 (21-26)	18	$21.2 \pm 2.6 (18.5 - 24.5)$	14.3 (12.5-16.0)
Excretory	$95 \pm 8.3 (84-104)$	$72 \pm 4.8 (65-78)$	-	94	$94 \pm 10.4 (85 \text{-} 108)$	75 (67-82)
Pore Pharynx	90.6 ± 7.5 (76-99)	$81 \pm 7.3 (70-91)$	-	-	$100 \pm 14.2 (84-122)$	80 (79-81)
MB	69.1 ± 5.2 (63.9-77.6)	$61.1 \pm 8.6 (53.3-80.8)$	-	53	$67.6 \pm 7.6 (57.2-75.4)$	70.2 (69.9-70.4)
PUS	$10.7 \pm 1.6 (9-13)$	-	-	-	$19.8 \pm 3.3 (16.5 - 24.0)$	-
Head- vulva	$449 \pm 62.1 (364-514)$	-	-	-		-
Vulva- anus	$41.7 \pm 6.0 (33-48)$	-	-	-	$42 \pm 10.4 (22-51)$	-
Tail	$36 \pm 5.7 (29-44)$	24 ± 7.3 (19-37)	-	49	$37 \pm 4.7 (30-43)$	34 (34-35)
Spicules	-	$14.6 \pm 0.7 (14-16)$	-	17	,	14.3 (14.0-14.5)
Gubernac ulum	-	4.8 ± 0.5 (4-5)	-	5		4.8 (4.5-5.0)

Measurements are in μm and in the form mean \pm SD (range).

Discussion

Hoplotylus femina was originally described by s'Jacob (1960) as the type species of the genus from the rhizosphere of *Quercus robur* and Chamaecyparis lawsoniana in Wageningen, The Netherlands. It is characterised by having well developed stylet with tulip-shape knobs, four areolated lateral lines, conical tail with smooth or coarse irregular annulation in distal region, males with higher and asymmetrical head and developed bursa. In the original description of the species (s'Jacob, 1960), sperm cells or functional spermatheca was not seen in the female reproductive system, but a spherical to slightly oblong spermatheca with spheroid to slightly elongate sperm cells were commonly observed in other reports, even in populations apparently lacking males (Sturhan and Lis kova, 2007). In Iranian population of H. femina, a functional spermatheca filled with spheroid sperm cells was observed in the female reproductive system.

The genus *Hoplotylus*, has four other known species namely H. silvaticus Bernard & Niblack, 1982., H. sjacobi Bernard & Niblack, 1982., H. montanus Minagawa, 1984, and H. triversus (Minagawa, 1984) Siddiqi, 1986. H. femina is morphologically and morphometrically similar to H. silvaticus, H. montanus and H. sjacobi. It can be distinguished from H. silvaticus by its longer tail (c' = 2.6-2.9 vs. 1.7-2.5 and c = 14.2-19.2 vs. 15-28.7) and also tail shape (posterior tail region is smooth or coarsely and irregularly annulated in H. femina vs. tail tip annulated dorsally in H. silvaticus). It differs from H. montanus by having shorter stylet (21-26 vs. 26-30.3µm), more posterior location of phasmids in females and different tail end in males (pointed vs. clavate). This species can also be distinguished from H. *sjacobi* by having shorter tail (c' = 2.6-2.9 vs. 3.3-4.3 and c = 14.2-19.2 vs. 12.1-14) and morphology of stylet knobs (closely appressed to shaft and edges in profile are parallel in H. femina vs. stylet knobs extending away from shaft and edges in profile diverge anteriorly in *H. sjacobi*).

In this study, *H. femina* was recovered from the rhizosphere of Caucasian elm (*Zelkova carpinifolia*) in Deylaman region, Guilan province. Morphological and morphometric characters of the Iranian population of *H. femina* fit well with the data in original description of the species by s'Jacob (1960) and the population reported from Spain by Palomares-Rius *et al.* (2016).

Molecular analysis

Amplification of D2-D3 expansion segment of the 28S rRNA gene from a single nematode produced a fragment of 686 nt (accession number KY614234). The Blast search using this sequence revealed the highest identity with other available sequences of this region for *H. femina* isolates from Spain (98% identity with KU513546, 98% identity with KU513544 and 95% identity with KU513543).

Iranian population of *Malenchus* (*Malenchus*) solovjovae Brzeski, 1989 (Figs 3 and 4; Table 3) Description

Female: Body straight or slightly ventrally curved upon fixation, tapering towards both ends. Lateral field with two lines under light microscope, originating at level of stylet, near middle of stylet shaft. Cuticle with distinct annuli, annulus 2.0-2.5µm wide at mid-body. Head narrower than body contour, with 4-5 fine annuli. Amphidial apertures large and sinuous shaped. Stylet delicate, with small knobs. Orifice of dorsal pharyngeal gland behind stylet knobs at about 1µm from base of stylet. Median bulb fusiform, valvular apparatus not Excretory pore at level of anterior part of pharyngeal bulb. Reproductive monodelphic, ovary outstretched with one row of oocytes. Vulva not sunk in body, with epiptygmata. Spermatheca rounded, offset and filled with globular sperm. Post vulval uterine sac 12-14µm long. Prophasmids located at 3-9 annuli anterior to vulva. Tail conical with a short filiform tip.

Male: Similar to females except for reproductive system and rather longer tail. Testis single, spermatocytes arranged in one row. Spicules sickle shaped. Bursa prominent, cloacal. Gubernaculum 5 μm long. Tail with pointed tip.

Discussion

Malenchus solovjovae was originally described by Brzeski (1988) from various shrubs and birch tree (*Betula* sp.) roots in Karelia, USSR. It is distinguished by annuli 2.0-2.5µm wide at mid-body, fusiform, more elongated and narrow median bulb, prophasmids located at 2-9 annuli anterior to vulva, 27-36 annuli between vulva and anus, and long tail in relation to vulva-anus distance. This species is morphologically and morphometrically similar to M. undulatus Andrássy, 1981, M. pachycephalus Andrássy, 1981 and M. machadoi (Andrássy, 1963) Andrássy, 1968. It most closely resembles M. undulatus but it differs from it by having wider annuli (2.0-2.5 vs. 1.7-1.9µm), different shape of median bulb (fusiform and narrow vs. relatively well developed), positions of prophasmids (2-9 vs. 7-15 annuli anterior to vulva) and less annuli between vulva and anus (27-36 vs. 44-47 annuli). It can be distinguished from M. pachycephalus by having narrower head, position of the prophasmids (2-9 vs. 11 annuli anterior to vulva), longer tail in relation to vulva-anus distance and shape of spicules (sickle shaped vs. straight). And finally, this species differs from M. machadoi by having wider annuli (2.0-2.5 vs. 1.6-1.8µm), longer tail in relation to vulva-anus distance, less annuli between vulva and anus (27-36 vs. 52-54 annuli), longer and different shaped spicules (18-19 vs. 15-16µm, sickle shaped vs. arcuate).

In this survey, this species was recovered from wood sample of a dead pine tree (*Pinus* sp.) in Siahkal region, Guilan province. The Iranian population of *M. solovjovae* is morphologically and morphometrically similar to the type population, except for a slightly longer stylet (10-11 vs. 8.0-9.5µm) and number of annuli between vulva and anus (35-39 vs. 27-36).




Figure 3 Iranian population of *Malenchus solovjovae* Brzeski, 1989. A: Female entire body; B: Male entire body; C: Female pharyngeal region showing excretory pore; D: Female head end; E: Origination of Lateral field; F: Vulval region showing prophasmid; G: Female tail; H: Male posterior end showing spicules and caudal alae. (Scale bars: $C-H = 10\mu m$; A, $B = 20\mu m$).




Figure 4 Iranian population of *Malenchus solovjovae* Brzeski, 1989. A: Beginning of lateral field; B, C: Female head end; D: Amphidial fovea; E: lobed spermatheca with sperm cells; F: Vulval region showing post uterine sac (PUS, arrowhead); G: Lateral field at mid-body; H: Female tail; I: Male caudal alae; J: Spicules; K: Prophasmid in female. (all scale bars = $10\mu m$).

Table 3 Morphometrics of *Malenchus solovjovae* from Iran and the data from type population.

Characters	Guilan province		Brzeski, 1989			
	Female	Male	Velika Manga population		Kroshnozero population	
			Female	Male	Female	
n	8	3	31	5	5	
L	$461 \pm 25.8 (428-504)$	$445.3 \pm 17.2 (427-461)$	439 (403-466)	427 (412-451)	424 (397-444)	
a	$24.8 \pm 1.2 (23.4-27.0)$	$26.8 \pm 2.0 (24.9 \text{-} 28.8)$	19 (18-21)	22 (21-23)	19 (17-20)	
b	$4.5 \pm 0.2 (4.3 - 4.9)$	$4.4 \pm 0.2 (4.2 - 4.6)$	4.8 (4.5-5.2)	4.7 (4.5-4.8)	4.8 (4.5-5.1)	
c	$6.3 \pm 0.3 \ (6.0 - 6.8)$	$5.0 \pm 0.1 (4.9 - 5.1)$	5 (4.5-6.3)	3.9 (3.8-4.1)	4.9 (4.5-5.6)	
c'	$7.2 \pm 0.3 (6.9 - 7.7)$	$7.6 \pm 0.2 (7.5 - 7.9)$	8.7 (7.6-10)	8.7 (8.1-9.5)	8.2 (6.7-8.9)	
V or T	$68.3 \pm 0.7 (67.2-69.5)$	$36.1 \pm 1.7 (34.2-37.3)$	66 (64-70)	-	65 (64-68)	
V'	$81.1 \pm 1.1 \ (80.1-83.4)$	-	82 (81-84)	-	82	
Stylet	$10.4 \pm 0.5 (10-11)$	$10.3 \pm 0.6 (10\text{-}11)$	9 (8.5-9.5)	9	8.4 (8-9)	
MB	$44.0 \pm 0.9 (42.6 - 45.0)$	$43.5 \pm 1.4 (42.2 - 45.0)$	48 (43-51)	48-50	47 (46-48)	
Excretory pore	$83.8 \pm 6.9 (74-98)$	$78.7 \pm 2.9 (77-82)$	84 (78-94)	79-88	81 (77-87)	
Pharynx	$101.5 \pm 3.7 (98-109)$	$100.3 \pm 1.5 (99-102)$	91 (85-96)	87-95	89 (88-92)	
Annuli width	$2.1 \pm 0.1 (2.0 - 2.2)$	-	2.3 (2.0-2.5)	-	2.3 (2.0-2.5)	
PUS	$0.7 \pm 0.0 (0.6 \text{-} 0.8)$	-	-	-	-	
Head-vulva	$314.5 \pm 17.9 (291-347)$	-	289 (264-304)	-	277 (252-300)	
Vulva-anus	$73.3 \pm 6.2 (63-80)$	-	62 (54-70)	-	61 (57-65)	
Tail	$72.9 \pm 4.7 (64-77)$	$89.0 \pm 1.7 (87-90)$	89 (67-98)	105-133	86 (79-91)	
Spicules	-	$19.3 \pm 0.6 (19-20)$	-	18-21	-	
Gubernaculum	-	$5.0 \pm 0.0 (5-5)$	-	3-5	-	

Measurements are in μm and in the form mean \pm SD (range).

Malenchus acarayensis (Malenchus) Andrássy, 1968

(Figs 5 and 6; Table 4)

Description

Female: Body straight or slightly ventrally curved upon fixation. Lateral field originating at about one-third of procorpus with two lines under light microscope Cuticle with distinct annuli, annulus 1.0-1.7 µm wide at the middle of the body. Head 4-5µm wide, with 5-6 fine annuli. Amphidial apertures large and sinuous in shape. Stylet delicate with small knobs. Orifice of dorsal pharyngeal gland about 1µm from knobs. Median bulb rather weak, valvular apparatus small. Excretory pore at level of anterior part of Reproductive pharyngeal bulb. system monodelphic, ovary outstretched with one row of oocytes. Vulva with epiptygmata. Spermatheca oval, offset and filled with rounded sperm cells. Post vulval uterine sac about half vulval body width long. Prophasmids at 7-14 annuli anterior to vulva. Tail straight with sharp to thread-like tip.

Males: Similar to females except for rather finer cuticular annuli (0.7 µm at middle of the

body). Bursa cloacal and finely crenate. Tail similar to female with fine and indistinguishable cuticular annuli.

Discussion

Malenchus acarayensis was originally described by Andrássy (1968) from Paraguay. Males of this species were described by Knobloch (1976) from low bog-like area in association with white birch, Betula papyrifera Marsh, Michigan, USA. It is characterised by 1.2-1.4µm wide annuli at mid-body. rather weak median bulb prophasmids at 7-14 annuli anterior to vulva and long tail in relation to vulva-anus distance. Morphological and morphometric characters of the Iranian population of M. acarayensis fit well with the original description. M. acarayensis is morphologically and morphometrically similar to M. shaheenai Khan & Ahmad, 1991. It can be distinguished from M. shaheenai by having different origination of lateral field (1/3 to mid procorpus vs. at level of median bulb) and shape of spermatheca (oval vs. oblong pouch-like).




Figure 5 Iranian population of *Malenchus acarayensis* Andrássy, 1968. A: Beginning of lateral field; B: Female head end; C: Male entire body; D: Female entire body; E: Vulval region showing prophasmid; F: Lateral field at mid-body; G: Female tail; H: Male posterior end showing spicules and caudal alae. (Scale bars: A, B, E-J = $10\mu m$; C, D = $20\mu m$).




Figure 6 Iranian population of *Malenchus acarayensis* Andrássy, 1968. A: Origination of Lateral field; B: Female head end; C: Amphidial fovea; D: Female pharyngeal region showing excretory pore (arrowhead); E: Vulval region showing prophasmid; F: Lateral field at mid-body; G: Posterior region of female showing filled spermatheca, vulval region and post uterine sac (PUS, arrowhead); H: Male caudal alae; I: Spicules; J: Female tail. (Scale bars = $10\mu m$).

Table 4 Morphometrics of *Malenchus acarayensis* from Iran and the data from other reports.

Characters	Guilan province		Andràssy, 1968	Knobloch, 1976	Gomez-Barcina et al.,	1992
	Female	Male	Female	Male	Female	Male
n	4	2	-	5	20	5
L	$345.3 \pm 9.2 (340-359)$	331, 342	280-410	300 (270-330)	$353 \pm 23.5 (305-389)$	$344 \pm 23 (306-365)$
a	$22.3 \pm 0.7 (21.3 - 22.7)$	36.8, 38	19-26	25 (21-29)	$21.8 \pm 2.3 (18.2 - 28.2)$	$24.4 \pm 2.6 (22.4-27.8)$
b	$4.4 \pm 0.2 (4.2 - 4.5)$	4.5, 4.6	-	4.3 (3.8-4.7)	$4.5 \pm 0.3 (3.9-5.0)$	$4.5 \pm 0.3 (4.1-5)$
c	$4.2 \pm 0.1 \ (4.1 - 4.4)$	4.6, 4.7	3.8-5.5	4.1 (3.6-4.5)	$4.5 \pm 0.5 (3.8 - 5.5)$	$3.8 \pm 0.2 (3.6-4)$
c'	$9.7 \pm 0.5 \ (9.1 \text{-} 10.4)$	10.1, 10.6	7.5-10	-	$8.8 \pm 1.4 (7.0 - 11.9)$	$9.8 \pm 1.2 (8.2 - 11)$
V or T	$61.5 \pm 1.0 (60.2 - 62.5)$	15.7, 21.3	61-66	37 (32-41)	$62 \pm 2.2 (60-66)$	$43 \pm 4.6 (37-48)$
V'	$80.7 \pm 1.7 (79.1-82.6)$	-	-	-	$80 \pm 1.5 (78-80)$	-
Stylet	8.5 ± 0.6 (8-9)	7	8-8.5	8	$8.5 \pm 0.5 (8 \text{-} 9.5)$	$8.6 \pm 0.7 (7.5-9)$
MB	$47.0 \pm 0.4 (46.8 - 47.6)$	44.6, 45.9	47	-	$47 \pm 2.4 (41-50)$	$48 \pm 3.2 (42-50)$
Excretory pore	$60.8 \pm 5.0 (56-66)$	53, 54	-	-	$62 \pm 7.2 (50-78)$	$64 \pm 4.9 (57-69)$
Pharynx	$78.8 \pm 2.4 (77-82)$	74	70-80	-	$78 \pm 6.7 (70-96)$	$76 \pm 6.9 (67-84)$
Annuli width	$1.3 \pm 0.1 (1.2 - 1.4)$	0.7	1-1.7	-	1.4-1.7	1.2 ± 0.06 (1.2-1.3)
PUS	$7.5 \pm 1.3 (6-9)$	-	-	-	5.5-8	-
Head-vulva	$212.3 \pm 3.0 (209-216)$	-	-	-	-	-
Vulva-anus	$50.8 \pm 5.7 (45-57)$	-	-	-	$55 \pm 4.1 (51-64)$	-
Tail	$82.3 \pm 3.3 (78-86)$	71, 73	70-76	-	$79 \pm 6.6 (69-89)$	$90 \pm 7.6 (77-96)$
Tail/vulva-anus		-	1.1-1.5	-	$1.4 \pm 0.1 (1.3 \text{-} 1.6)$	-
Spicules	-	12	-	15 (13-16)	-	$15.3 \pm 1.5 (13-16)$
Gubernaculum	-	3, 4	-	4 (4-5)	-	$4.2 \pm 0.8 (3-5)$

Measurements are in μm and in the form mean \pm SD (range).

In the present study it was recovered from the rhizosphere of Caucasian elm ($Zelkova\ carpinifolia$) in Deylaman region, Guilan province. The Iranian population of M. acarayensis is similar to the type populatiopn based on its morphology and morphometric data; however, it has slightly shorter spicules (12 vs. 13-16 μ m). Namadipour $et\ al.$ (2014) reported this species from Zanjan province as a short note.

References

Alem, M., Jamali, S., Tanha Maafi, Z. and Pedramfar, H. 2010. Identification of three species of the genus *Xiphinema* in the stone fruit trees in Guilan province. Proceeding of the 19th Iranian Plant Protection Congress, 1-3 August 2010, Tehran, Iran. p. 640.

Aliramaji, F., Pourjam, E., Karegar, A. and Eskandari, A. 2012. Three species of *Nagelus* (Tylenchina: Merliniinae) from Iran. Iranian Journal of Plant Protection Science, 43 (2): 301-312. (In Persian).

Andrássy, I. 1968. Fauna Paraguayensis. 2. Nematoden aus den Galeriewaldern des Acaray-Flusses. Opuscula Zoologica Budapest, 8 (2): 167-312.

Andrássy, I. 1981. The genera and species of the family Tylenchidae "Orley, 1880 (Nematoda). The genus *Malenchus* Andrássy, 1968. Acta Zoologica-Academiae Scientiarum Hungaricae, 27: 1-47.

Bernard, E. and Niblack, T. 1982. Review of *Hoplotylus* S'JACOB (Nematoda: Pratylenchidae). Nematologica, 28 (1): 101-109.

Brzeski, M. 1988. *Malenchus parvus* sp. n., *M. solovjovae* sp. n. and observations on *M. leiodermis* (Nematoda, Tylenchidae). Nematologica, 34 (1): 47-56.

De Grisse, A. T. 1969. Redescription ou modification de quelques techniques dans L'etude des nematodes phytoparasitaires. Mede. Rijks. fak. LandbWet Gent, 34: 351-369.

Divsalar, N., Jamali, S., Pedramfar, H. and Komeili, R. 2011. Identification of plant

- parasitic nematodes of citrus orchards in Guilan east and Mazandaran west province. Journal of Plant Protection. 25 (2): 168-177. (In Persian).
- Ebsary, B. A. 1991. Catalog of the order Tylenchida (Nematoda). Ottawa, Canada, Research Branch Agriculture Canada.
- Eskandari, A., Karegar, A. and Pourjam, E. 2009. Three species of sheath nematodes from Iran. Journal of Animal Biology, 3: 1-7. (In Persian).
- Geraert, E. 2008. The Tylenchidae of the World- Identification of the Family Tylenchidae (Nematoda). Gent, Belgium, Academia Press.
- Geraert, E. and Raski, D. 1986. Unusual Malenchus species (Nematoda: Tylenchidae). Nematologica, 32 (1): 27-55.
- Gómez-Barcina, A., Geraert, E., Castillo, P. and Pais, G. 1992. Three Malenchus species from Spain (Nemata: Tylenchidae) with a note on the amphidial opening in the genus. Fundamental and Applied Nematology 15: 153-157.
- Heydari, R., Maafi, Z. T. and Decraemer, W. 2014. Morphological and molecular characterisation of *Trichodorus golestanensis* (Nematoda: Trichodoridae), a new species from Iran. European Journal of Plant Pathology, 140 (2): 329-340.
- Knobloch, N. A. 1976. The genus *Malenchus* (Nematoda: Tylenchida) with descriptions of two species from Michigan. Journal of Nematology, 8 (1): 53.
- Minagawa, N. 1984. New species of *Hoplotylus* and *Pratylenchoides* (Tylenchida: Pratylenchidae) from Japan. Japanese Journal of Nematology, 14: 15-19.
- Namadipour, F., Eskandari, A. and Asghari, R. 2014. Identification of two nematode species of the genera *Malenchus* and *Criconema* from Lorestan Province. Proceeding of the 21st Iranian Plant Protection Congress, 23-26 August 2014, Urmia, Iran. P. 310.
- Nasaj Hoseini, S. M., Pourjam, E. and Seraji, A. 2004. Identification of two ectoparasitic nematode species from tea plant rhizosphere in Guilan province. Proceeding of the 16th

- Iranian Plant Protection Congress, 28-31 August 2004, Tabriz, Iran. p. 318.
- Nunn, G. B. 1992. Nematode molecular evolution. Ph.D. Dissertation, University of Nottingham, Nottingham.
- Palomares-Rius, J. E., Cantalapiedra-Navarrete, C., Archidona-Yuste, A., and Castillo, P. 2016. *Hoplotylus femina* s' Jacob, 1960 (Nematoda: Pratylenchidae) from Spain with molecular phylogenetic relationships inferred by D2-D3 expansion fragments of 28S and the partial 18S rRNA gene sequences. Nematology, 18 (5) 559-569.
- Panahandeh, Y., Pourjam, E., Aliramaji, F. and Pedram, M. 2015. Data on some members of the family Tylenchidae (Nematoda: Tylenchina) from Iran. Biologia, 70 (10): 1376-1387.
- Pedramfar, H., Kheiri, A. and Pourjam, E. 1998. Occurrence of *Heterodera oryzae* in rice fields of Guilan Province with morphological characters. Proceedings of the 13th Iranian Plant Protection Congress, 23-27 August 1998, Karaj, Iran. p. 87.
- Pedramfar, H., Pourjam, E. and Kheiri, A. 2011. Plant parasitic nematodes associated with rice in Guilan Province, Iran. Journal of Plant Pathology, 37: 258-301. (In Persian).
- s'Jacob, J. 1959. *Hoplotylus femina* ng, n. sp. (Pratylenchinae: Tylenchida) associated with ornamental trees. Nematologica, 4 (4): 317-321.
- s'Jacob, J. 1979. The description of the male of *Hoplotylus*, collected from an oak forest. Nematologica, 25 (1): 147-151.
- Shahabi, S., Kheiri, A., Rakhshandehroo, F. and Jamali, S. 2016. Occurrence and distribution of nematodes in rice fields in Guilan province, Iran and the first record of *Mylonchulus polonicus* (Stefanski, 1915) Cobb, 1917 (Nematoda: Mononchina). Journal of plant Protection Research, 56 (4): 420-429.
- Siddiqi, M. R. 1986. Tylenchida: Parasites of Plants and Insects. Commonwealth Agricultural Bureaux, Farnham Royal, Slough, UK.

- Siddiqi, M. R. 2000. Tylenchida Parasites of Plants and Insects, 2nd Edition. Wallingford, UK, CAB International.
- Soleymanzadeh, M., Pedram, M., Pourjam, E., and Álvarez-Ortega, S. 2016. Description of *Lelenchus brevislitus* n. sp. (Nematoda: Tylenchidae), an example of a cryptic species from Iran and its phylogenetic relationships with other species in the family. Nematology, 18(8): 987-998.
- Sturhan, D. and Liskova, M. 2007. Notes on morphology, distribution and ecology of *Hoplotylus femina* (Tylenchida, Pratylenchidae). Nematologia Mediterranea, 35 (2): 243-246.
- Sumenkova, N. I. 1988. *Paramalenchus anthrisculus* gen. et sp. n. (Nematoda, Tylenchidae) from the rhizosphere of plants in a floodland meadow. Zoologichesky Zhurnal, 67: 1073-1076.
- Tanha Maafi, Z. 1998. *Pratylenchus neglectus* and *P. thornei* endoparasite nematodes associated with wheat in Gilan province. Proceedings of the 13th Iranian Plant

- Protection Congress, 23-27 August 1998, Karaj, Iran. p. 66.
- Tanha Maafi, Z. and Decraemer, W. 2002. Description of *Trichodorus gilanensis* n. sp. from a forest park in Iran and observations on Paratrichodorus tunisiensis (Siddiqi, 1963) Siddiqi, 1974 (Nematoda: Diphtherophorina). Nematology, 4 (1): 43-54.
- Tanha Maafi, Z. Ebrahimi, N., Khozaini, F., Barooti, S and Karegar, A. 2006. Occurrence of *Scutellonema brachyurus* (Steiner, 1938), Andrasy, 1958 in Damghan, Gilan and Golestan regions. Journal of Plant Pathology, 42 (1): 197-200 (In Persian).
- Whitehead, A. G. and Hemming, J. R. 1965. A comparison of some quantitative methods of extracting small vermiform nematodes from soil. Annual Applied Biology, 55: 25-38.
- Williams, B. D., Schrank, B., Huynh, C., Shownkeen, R. and Waterston, R. H. 1992. Α genetic mapping system in Caenorhabditis elegans based on polymorphic sequence tagged sites. Genetics, 131: 609-624.

اولین گزارش از جنس (Nematoda: Hoplolaimidae) و توصیف دو گونه از جنس (Malenchus (Nematoda: Tylenchidae) در ایران

يريسا جلالي نسب و رامين حيدري*

گروه گیاهپزشکی، پردیس کشاورزی و منابع طبیعی، دانشگاه تهران، کرج، ایران. * پست الکترونیکی نویسنده مسئول مکاتبه: rheydari@ut.ac.ir دریافت: ۳۰ بهمن ۱۳۹۵؛ پذیرش: ۲۸ آذر ۱۳۹۶

چکیده: به منظور بررسی فون نماتدهای انگل گیاهی جنگلهای شرق استان گیلان، نمونههایی از خاک و ریشه گیاهان این مناطق جمعآوری شده و نماتدهای موجود در آنها جداسازی شدند. شناسایی براساس ویژگیهای ریختسنجی و ریختشناسی انجام شد. در بین نماتدهای شناسایی شده جمعیتی از جنس H. femina جداسازی شد که سپس به عنوان DNA شناسایی شد. شناسایی مرفولوژی به به وسیله توالی بخشهایی از زیرواحد بزرگ DNA ریبوزومی نیز تأیید گردید. همچنین دو جمعیت از جنس solovjovae جداسازی شد که براساس خصوصیات ریختسنجی و ریختشناسی، M. solovjovae و میشوند. گونه شدند. دو گونه H. femina و H. femina برای اولین بار از ایران گزارش میشوند. گونه گونه نیز توصیف از ایران گزارش شده بود، لذا این گونه نیز توصیف و تشریح شده است.

واژگان کلیدی: M. acarayensis Malenchus solovjovae Hoplotylus femina زیرواحد بزرگ DNA ریبوزومی