

Research Article

Encyrtidae (Hymenoptera: Chalcidoidea) in southern area of Kerman province with some new generic and specific records

Mahla Shojaey¹, Mohammad Khayrandish^{1*}, Seyed Massoud Madjdzadeh² and Hossein Lotfalizadeh³

1. Department of Plant Protection, Faculty of Agriculture, Shahid Bahonar University of Kerman, Kerman, Iran.

2. Department of Biology, Faculty of Sciences, Shahid Bahonar University of Kerman, Kerman, Iran.

3. Plant Protection Research Department, East-Azarbaijan Agricultural and Natural Resources Research & Education Center, AREEO, Tabriz, Iran.

Abstract: The present study provides information about parasitoid wasps of the family Encyrtidae (Hymenoptera: Chalcidoidea) collected from south of Kerman province. The specimens were caught using Malaise traps during April to September 2017. Totally 16 species belonging to 10 genera of encyrtid wasps were collected and identified. Among them, *Cerchysiella* Girault, 1914 and *Charitopus* Foerster, 1856 are new generic records for the fauna of Iran. Four species viz. *Charitopus manukyani* Sakhnov, 1993; *Cheiloneurus submuticus* Thomson, 1876; *Mayridia myrlea* (Walker, 1838) and *Prionomitus tiliaris* (Dalman, 1820) were recorded for the first time from Iran. Diagnostic characteristics are mentioned for the newly recorded species. In addition, biological associations as well as geographical distribution for all reported species are presented. An updated list of encyrtid wasps of Kerman province is also cataloged.

Keywords: encyrtid wasp, new record, Iran, fauna, Kerman province

Introduction

Encyrtidae (Hymenoptera: Chalcidoidea) with 506 genera and over 4000 known species worldwide is a large family which contains two subfamilies, Encyrtinae and Tetracneminae (Noyes, 2020). Some species are considered as primary endoparasitoids and hyperparasitoids of wide variety of hosts, including species of Arachnida and many insect orders (Noyes and Hayat, 1994). They have been successfully used as biological control agents in some parts of the world (Dahlsten *et al.*, 2005; Bauer *et al.*, 2008; Noyes, 2020). Encyrtids are the most efficient parasitoids of the scale insects pest species in the

families Coccidae, Diaspididae and Pseudococcidae, and less frequently, Aphidoidea and Psylloidea (Hemiptera) (Guerrieri and Noyes, 2000, 2005, 2009; Guerrieri and Viggiani, 2005; Evans and Abd-Rabou, 2013). They are distributed successfully in nearly all agroecosystems but their greatest diversity is in the tropical and subtropical areas (Grissell and Schauf, 1990; Noyes *et al.*, 1997).

Despite the fact that Iran has various geographical regions and diverse types of vegetation, the fauna of Iranian Encyrtidae is poorly known. The first reliable checklist of Iranian Encyrtidae was published by Fallahzadeh and Japoshvili (2010) who reported 93 species in 32 genera from different parts of Iran. Since then more contributions were made to the encyrtids fauna of Iran (Lotfalizadeh *et al.*, 2010a, b; Golpayegani *et al.*, 2010; Fallahzadeh *et al.*, 2011, 2014; Hesami *et al.*, 2012; Fallahzadeh

Handling Editor: Ali Asghar Talebi

*Corresponding author: m.khayrandish@uk.ac.ir

Received: 02 April 2020, Accepted: 02 July 2020

Published online: 18 July 2020

and Japoshvili, 2013; Moghbeli-Gharaei *et al.*, 2013; Ebrahimi *et al.*, 2014; Lotfalizadeh and Bab-Morad, 2015; Japoshvili *et al.*, 2015, 2016; Lotfalizadeh *et al.*, 2016; Mohammadpour *et al.*, 2016; Feli kohikheili *et al.*, 2016; Fallahzadeh and Japoshvili, 2017; Fallahzadeh *et al.*, 2018; Japoshvili and Fallahzadeh, 2018; Morravej *et al.*, 2018; Yousefi *et al.*, 2018) that increased the number of species and genera to 173 and 51, respectively.

The aim of this work was to increase the knowledge on encyrtids fauna of Iran, as a part of our ongoing project on taxonomy and biodiversity of Iranian Chalcidoidea.

Materials and Methods

The sampling was carried out in south of Kerman province, southeast of Iran during April to September 2017 (Fig. 1). Wasps were collected using Malaise traps in different habitats and were preserved in 75% Ethanol until they were mounted on cards and identified. Prior to mounting the specimens were treated with Acetic

acid' vapor in order to avoid collapsing (Noyes, 1982). The identifications were made using Noyes and Hayat (1984), Noyes *et al.* (1997) and Zhang and Huang (2004). The morphological terminology and classification follows Grissell and Schauff (1990, 1997). External morphology was illustrated using an Olympus™ SZH, equipped with an Omax (18Mp) A35180U3 digital camera. The images were then processed using Combine ZM and Adobe Photoshop® CS6 programs. The voucher specimens are deposited in Zoological Museum of Shahid Bahonar University of Kerman, Kerman, Iran (ZMSBUK).

Results

Totally, 65 specimens of Encyrtidae belonging to the subfamily, Encyrtinae representing 10 genera and 16 species were collected and identified, of which two genera and five species are new records for Iranian fauna. The genera and species are listed alphabetically. Short description of characteristics is presented for the newly recorded species.




Figure 1 Geographic map of Kerman province in southeast of Iran. Red star indicates the sampling sites in south of Kerman province.

Family: Encyrtidae Walker, 1837

Subfamily: Encyrtinae Walker, 1837

Genus: *Blastothrix* Mayr, 1876

Blastothrix sp. (Fig. 2)

Material examined: Iran: Kerman province: Jiroft, Dalfard, Bondar (29°01'31.4" N, 57°36'56.1" E, 2390m), 17.VII-27.VIII.2017, 1♀, Malaise trap, leg.: S.M. Madjzadeh.

Short description: Body dark blue (Fig. 2A); antennal formula 1, 1, 6, 3 (scape, pedicel, funicle, clava), basal antennal segment about 3.5 times longer than width, pedicel longer than first funicular segment, clava rounded to apex (Fig. 2B), ventral margin of torulus lower than ventral margin of eye (Fig. 2C); mesoscutum without notauli and scutellum without apical tuft of setae (Fig. 2D), mesopleuron enlarged posteriorly and more or less touching base of metasoma, forewing with dark spot on the anterior margin, marginal vein longer than wide and as long as stigmal vein (Fig. 2E).

Genus: *Cerchysiella* Girault, 1914

Cerchysiella sp. (Fig. 3)

Material examined: Iran: Kerman province: Jiroft, Bahram Abad (28°18'54.4" N, 57°37'1.5" E, 682m), 09-29.IV.2017, 1♀, Malaise trap, leg.: S.M. Madjzadeh.

Short description: Body dark blue (Fig. 3A); antennal formula 1, 1, 6, 3, all funicular segments longer than broad, clava apically rounded (Fig. 3B), ventral margin of torulus lower than ventral margin of eye (Fig. 3C); mesoscutum without notauli and scutellum without apical tuft of setae, mesoscutum and scutellum with coarse silvery hairs, scutellum nearly entirely smooth (Fig. 3D), forewing hyaline, lineal calva of forewing posteriorly open, marginal vein as long as postmarginal vein, stigmal vein longer than marginal vein (Fig. 3E), ovipositor distinctly protrudes.

Comment: The genus *Cerchysiella* is recorded for the first time from Iran. This species seems to be an undescribed species but for its description, further specimens are needed.

Genus: *Charitopus* Foerster, 1856

Charitopus manukyani Sakhnov, 1993 (Fig. 4)

Material examined: Iran: Kerman province: Bam, Hemat Abad (29°08'19.6" N, 57°58'05.1" E, 1673m), 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Short description: Head and mesosoma dark green (Fig. 4A); antennal formula 1, 1, 6, 3, all funicular segments longer than broad, some segments of antennal funicle more than 2 times longer than wide, clava apically rounded (Fig. 4B), torulus lower on head, its dorsal margin well below ventral margin of eye (Fig. 4C); notauli present (Fig. 4D), mesopleuron clearly separated from base of metasoma by propodeum, the propodeum touching metacoxa, metacoxa entirely yellow, marginal vein longer than postmarginal vein and stigmal vein longer than marginal vein (Fig. 4E); metasoma yellow with two dark lines in lateral margins; hypopygium extending to apex of metasoma (Fig. 4F).

Distribution in Iran: Kerman province.

General distribution: Palaearctic region (Noyes, 2020; Iran- New record).

Comment: This is the second report of this species from the Palaearctic region.

Genus: *Cheiloneurus* Westwood, 1833

Cheiloneurus submuticus Thomson, 1876 (Fig. 5)

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 22.V-04.VII.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Short description: Body dark blue (Fig. 5A); antennal formula 1, 1, 6, 3, all funicular segments not longer than broad, clava strongly obliquely truncate (Fig. 5B), dorsal margin of torulus well below ventral margin of eye (Fig. 5C); mesoscutum without notauli, scutellum apically without tuft of setae (Fig. 5D), mesopleuron enlarged posteriorly and more or less touching base of metasoma; forewing infusate (except base of forewing), marginal vein much longer than postmarginal vein and stigmal vein (Fig. 5E); hypopygium not extending to apex of metasoma.

Distribution in Iran: Kerman province.

General distribution: Palaearctic region (Noyes, 2020; Iran- New record).




Figure 2 *Blastothrix* sp., Female: A. Lateral view, B. Antenna, C. Head (frontal view), D. Mesosoma in dorsal view, E. Forewing.




Figure 3 *Cerchysiella* sp., Female: A. Lateral view, B. Antenna, C. Head (frontal view), D. Mesosoma in dorsal view, E. Forewing.




Figure 4 *Charitopus manukyani* Sakhnov, 1993, Female: A. Lateral view, B. Antenna, C. Head in frontal view, D. Mesosoma in dorsal view, E. Forewing, F. Metasoma in lateral view.




Figure 5 *Cheiloneurus submuticus* Thomson, 1876, Female: A. Lateral view, B. Antenna, C. Head in latero-frontal view, D. Mesosoma in dorsal view, E. Forewing.

Genus: *Homalotylus* Mayr, 1876***Homalotylus flaminus* (Dalman, 1820) (Fig. 6A)**

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 04.VII-26.VIII.2017, 1♀, Malaise trap, leg.: M. Purrezaali; Bam, Dehbakri, Bidkhun (29°07'22.6" N, 57°52'56.8" E, 2220m), 04-16.VII.2017, Malaise trap, leg.: M. Purrezaali; Bam, Hemat Abad (29°08'19.6" N, 57°58'05.1" E, 1673m), 22.V-04.VII.2017, 1♀, Malaise trap, leg.: M.

Purrezaali; Anbar Abad, Rudfargh (28°29'41" N, 58°09'56.2" E, 1429m), 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Distribution in Iran: Fars (Fallahzadeh *et al.*, 2006), Kermanshah (Ghahari *et al.*, 2010), Qazvin (Rastegar *et al.*, 2017), Khuzestan (Moravvej *et al.*, 2018) and Kerman provinces (Present study).

General distribution: Afrotropical, Australasia, Nearctic, Neotropical, Oriental and Palaearctic regions (Noyes, 2020).




Figure 6 Lateral view of collected female species of Encyrtidae in south of Kerman province. A. *Homalotylus flaminus* (Dalman, 1820), B. *Homalotylus nigricornis* Mercet, 1921.

***Homalotylus nigricornis* Mercet, 1921 (Fig. 6B).**

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 22.V-04.VII.2017, 1♀ & 04.VII-26.VIII.2017, 1♀, Malaise trap, leg.: M. Purrezaali; Anbar Abad, Bardeh, 05.VI-05.VII.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Distribution in Iran: Ardabil (Lotfalizadeh and Ebrahimi, 2001) and Kerman provinces (Present study).

General distribution: Palaearctic region (Noyes, 2020).

***Homalotylus near platynaspidis* Hoffer, 1963 (Fig. 7)**

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 04.VII-26.VIII.2017, 2♀, Malaise trap, leg.: M. Purrezaali; Bam, Hemat Abad (29°08'19.6" N, 57°58'05.1" E, 1673m), 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Short description: Body dark blue (Fig. 7A); antennal formula 1, 1, 6, 1, the second funicular segment and clava white (Fig. 7B), ventral margin of torulus lower than ventral margin of eye (Fig. 7C); pronotum, mesoscutum and scutellum with coarse silvery hairs, notauli present (Fig. 7D), mesopleuron clearly separated from base of metasoma by propodeum; forewing infusate and

with two hyaline areas, one hyaline area extending across entire breadth of wing in distal half of first third and the other hyaline area joining the outer side of large median dark spot, submarginal vein in distal third broadened and curved, marginal vein as long as postmarginal vein (Fig. 7E); hypopygium extending to apex of metasoma.

Distribution in Iran: Kerman province (Present study).

Comment: This specimen is similar to *Homalotylus platynaspidis*. But in *H. platynaspidis*, scape, pedicel and funicular

segments 1-5 black; segment 6 and club yellowish brown, in some specimen yellowish white; scape as long as pedicel and three and one-half basal funicular segments together; funicular segments quadrate, the basal three being a trifle elongate, remaining three segments as long as wide; postmarginal distinctly longer than stigmal vein (Hoffer, 1963). Therefore, this species maybe an undescribed species.

***Homalotylus quaylei* Timberlake, 1919** (Fig. 8).

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 04.VII-26.VIII.2017, 1♀, Malaise trap, leg.: M. Purrezaali.




Figure 7 *Homalotylus* near *platynaspidis* Hoffer, 1963, Female: A. Lateral view, B. Antenna, C. Head in frontal view, D. Mesosoma in dorsal view, E. Forewing.



Figure 8 Lateral view of collected female *Homalotylus quaylei* Timberlake, 1919 in south of Kerman province.

Distribution in Iran: Tehran (OILB, 1971), Khuzestan (Asadeh and Mossadegh, 1991; Novin *et al.*, 2000), Mazandaran (Maafi *et al.*, 1998) Fars (Lotfalizadeh, 2010), East Azarbaijan (Ghahari *et al.*, 2010) and Kerman provinces (Present study).

General distribution: Afrotropical, Oriental and Palearctic regions (Noyes, 2020).

Genus: *Leptomastix* Foerster, 1856

Leptomastix sp. (Fig. 9)

Material examined: Iran: Kerman province: Jiroft, Dalfard (29°00'33" N, 57°35' 32.4" E, 2121m), 13-30.VIII.2016, 1♀, Malaise trap, leg.: M. Changizi.

Short description: Head and mesosoma dark blue (Fig. 9A); antennal formula 1, 1, 6, 3, all funicular segments longer than broad, first

funicular segment longer than pedicel (Fig. 9B), ventral margin of torulus lower than ventral margin of eye (Fig. 9C); tegula dark, scutellum apically without tuft of setae, mesopleuron enlarged posteriorly and more or less touching base of metasoma, wings not reduced, forewing hyaline and extended beyond tip of metasoma, marginal vein longer than stigma, postmarginal vein longer than marginal vein, glabrous oblique stripe on disk of forewing more or less broadly interrupted (Fig. 9D); metasoma dark brown, hypopygium extending to apex of metasoma.

Genus: *Mayridia* Mercet, 1921

Mayridia myrlea (Walker, 1838) (Fig.10)

Material examined: Iran: Kerman province: Bam, Hemat Abad (29°08'19.6 "N, 57°58'05.1 "E, 1673m), 13-22.V.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Short description: Body metallic green (Fig. 10A); antennal formula 1, 1, 6, 3, all funicular segments not longer than broad, two last funicular segments squared, clava obliquely truncate and equal last three funicular segments (Fig. 10B), ventral margin of torulus lower than ventral margin of eye (Fig. 10C); mesosutum without notauli (Fig. 10D), scutellum with sculpturing same as that of mesoscutum, mesopleuron clearly separated from base of metasoma by propodeum, forewing with 2 longitudinal infusate rays, apex of forewings pale, marginal vein longer than postmarginal vein (Fig. 10E); hypopygium not extending to apex of metasoma.

Distribution in Iran: Kerman province.

General distribution: Palearctic region (Noyes, 2020; Iran- New record).

Genus: *Ooencyrtus* Ashmead, 1900

Ooencyrtus masii (Mercet, 1921) (Fig. 11A)

Material examined: Iran: Kerman province: Bam, Dehbakri, Bidkhun (29°07'22.6" N, 57°52'56.8" E, 2220m), 22.V-04.VII.2017, 1♀, Malaise trap, leg.: M. Purrezaali; Bam, Hemat Abad (29°08'19.6" N, 57°58'05.1" E, 1673m), 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali.




Figure 9 *Leptomastix* sp., Female: A. Lateral view, B. Antenna, C. Head in frontal view, D. Forewing.



Figure 10 *Mayridia myrlea* (Walker, 1838). Female: A. Lateral view, B. Antenna, C. Head in latero-frontal view, D. Mesosoma in dorsal view, E. Forewing.

Distribution in Iran: Fars as *Ooencyrtus* cf. *masii* (Sabahi *et al.*, 1998, 1999), Khojir protected area (Omid *et al.*, 2005) and Kerman provinces (Present study).

General distribution: Palaearctic region (Noyes, 2020).

Comment: This species is similar to *Ooencyrtus ocneriae* Hayat and Mehrnejad, 2018. But in *O. ocneriae* scutellum with very fine, polygonal reticulate sculpture much shallower than that on the mesoscutum, and smooth in about apical third; mid tibia (also fore and hind tibiae) white, without a sub-basal dark band; and stigmal vein 1.2 times as long as combined length of the marginal and postmarginal veins (Hayat and Mehrnejad, 2018).

***Ooencyrtus telenomicida* (Vassiliev, 1904)** (Fig. 11B)

Material examined: Iran: Kerman province: Bam, Dehbakri, Bidkhun (29°07'22.6" N, 57°52'56.8" E, 2220m), 22.V-04.VII.2017, 6♀, 04-17.VII.2017, 9♀, 17.VII-26.VIII.2017, 14♀, Malaise trap, leg.: M. Purrezaali; Bam, Hemat Abad (29°08'19.6"

N, 57°58'05.1" E, 1673m), 22.V-04.VII.2017, 1♀, Malaise trap, leg.: M. Purrezaali; Jiroft, Dalfard, Bondar (29°01'31.4" N, 57°36'56.1" E, 2390m), 17.VII-27.VIII.2017, 7♀, Malaise trap, leg.: M. Purrezaali; Jiroft, Dalfard, Bondar (29°00'36" N, 57°36'39.1" E, 2232m), 07-17.VII.2017, 2♀, Malaise trap, leg.: M. Purrezaali.

Distribution in Iran: Alborz, Tehran, Hamadan, Lorestan, Mazandaran, Markazi (Chodjai, 1968, 1989; Davatchi and Chodjai, 1969; Radjabi and Amir-Nazari, 1989; Modarres Awal, 1997); East Azarbaijan (Iranipour *et al.*, 1998), Kerman (Hashemi Rad *et al.*, 2002), Qazvin (Noori *et al.*, 2003), Kerman (Mohammadpour *et al.*, 2015, 2016), Guilan (Lotfalizadeh *et al.*, 2016).

General distribution: Oriental and Palaearctic regions (Noyes, 2020).

***Ooencyrtus* sp. 1** (Fig. 12)

Material examined: Iran: Kerman province: Bam, Hemat Abad (29°08'19.6"N, 57°58'05.1"E, 1673m), 22.V-04.VII.2017, 1♀, Malaise trap, leg.: M. Purrezaali.




Figure 11 Lateral view of collected female species of Encyrtidae in south of Kerman province. A. *Ooencyrtus masii* (Mercet, 1921), B. *Ooencyrtus telenomicida* (Vassiliev, 1904).



Figure 12 *Ooencyrtus* sp. 1, Female: A. Lateral view, B. Antenna, C. Forewing.

Short description: Head and mesoscutum dark green (Fig. 12A); ventral margin of torulus lower than ventral margin of eye, antennal formula 1, 1, 6, 3, antennal scape subcylindrical, all funicular segments longer than broad (Fig. 12B), first funicular segment not longer than pedicel, clava apically rounded and its length equal to the last three segments of funicle; mesonotum without notauli, scutellum dark brown and nearly entirely smooth, mesopleuron well developed and more or less touching base of metasoma, forewing hyaline, all coxa dark green, postmarginal vein longer than marginal vein and stigma, lineal calva not interrupted by setae (Fig. 12C); metasoma dark brown, hypopygium not extending to apex of metasoma.

***Ooencyrtus* sp. 2** (Fig. 13)

Material examined: Iran: Kerman province: Bam (29°06'01.7" N, 58°19'44" E, 1111m), 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali.

Short description: head and mesosoma dark blue (Fig. 13A); ventral margin of torulus lower than ventral margin of eye, antennal formula 1, 1, 6, 3, basal antennal segment more than 4 times longer than width, pedicel longer than first funicular segment, clava apically rounded (Fig. 13B); mesoscutum without notauli, scutellum with sculpturing barely same as that of mesoscutum, scutellum dark brown, mesopleuron well developed and more or less touching base of metasoma (Fig. 13C), forewing hyaline, lineal calva of forewing posteriorly open, all coxa yellowish brown, postmarginal vein longer than marginal vein, stigmal vein longer than postmarginal vein (Fig. 13D); metasoma dark brown, hypopygium not extending to apex of metasoma.

Genus: *Prionomitus* Mayr, 1876

***Prionomitus tiliaris* (Dalman, 1820)** (Fig. 14)

Material examined: Iran: Kerman province: Anbar Abad, Rudfargh (28°29'41" N, 58°09'56.2" E, 1429m), 27.VI-16.VII.2017, 1♀, 26.VIII-21.IX.2017, 1♀, Malaise trap, leg.: M. Purrezaali; Manujan (27°29'53.1" N, 57°33'43.4" E, 358m), 07.IV-05.V.2017, 1♀, Malaise trap, leg.: M. Purrezaali.




Figure 13 *Ooencyrtus* sp. 2, Female: A. Lateral view, B. Antenna, C. Body (lateral view), D. Forewing.

Short description: Head and mesosoma greenish blue (Fig. 14A); antennal formula 1, 1, 6, 3, first funicular segment shorter than pedicel, all funicular segments not longer than broad (Fig. 14B), ventral margin of torulus lower than ventral margin of eye (Fig. 14C); mesonotum without notauli (Fig. 14D), scutellum with sculpturing deeper than that of mesoscutum, mesopleuron clearly separated from base of metasoma by propodeum, forewings hyaline, lineal calva of forewing posteriorly open, postmarginal vein longer than marginal vein and shorter than stigmal vein (Fig. 14E); metasoma dark brown.




Figure 14 *Prionomitus tiliaris* (Dalman, 1820). Female: A. Lateral view, B. Antenna, C. Head in lateral view, D. Mesosoma in dorsal view, E. Forewing.

Distribution in Iran: Kerman province.

General distribution: Australasia, Nearctic and Palaearctic regions (Noyes, 2020; Iran-New record).

Genus: *Psyllaephagus* Ashmead, 1900

Psyllaephagus claripes Trjapitzin, 1967 (Fig. 15)

Material examined: Iran: Kerman province: Bam, Hemat Abad (29°08'19.6" N, 57°58'05.1" E, 1673m), 26.VIII-21.IX.2017, 2♀, Malaise trap, leg.: M. Purrezaali.

Distribution in Iran: Hamadan (Rajabi Mazhar and Sadeghi, 2006; Rajabi Mazhar et al., 2007) and Kerman provinces (Present study).

General distribution: Palaearctic region (Noyes, 2020).



Figure 15 Lateral view of collected female *Psyllaephagus claripes* Trjapitzin, 1967 in south of Kerman province.

Discussion

In the course of this survey of Encyrtidae in southern parts of Kerman province, 16

species belonging to one subfamily and 10 genera were identified. In total, 10 species were recorded for the first time from Kerman province. Two genera and five species were recorded for the first time for the fauna of Iran.

It seems *Homalotylus flaminus* is a cosmopolitan species and *Charitopus manukyani*, *Cheiloneurus submuticus*, *Homalotylus nigricornis*, *Mayridia myrlea*, *Ooencyrtus masii*, *Psyllaephagus claripes* are distributed only in the Palaearctic region. *Charitopus manukyani* is recorded only from Mongolia and this is the second record of this species from the Palaearctic region. Other species are distributed in the Palaearctic and other zoogeographic regions. Also *Charitopus manukyani* and *Psyllaephagus claripes* have not been recorded from neighbouring countries but other species have been reported from at least one of the neighbouring countries. The known hosts of Encyrtidae of Kerman province are species of the orders Hemiptera, Hymenoptera and Neuroptera (Fallahzadeh and Japoshvili, 2017). According to the literatures, most known species of Kerman province are parasitoids of Coccidae, Psyllidae and Diaspididae and some species are hyperparasitoid (Table. 1). The common species of this study were compared with neighboring countries and it was found that the study area has the largest number of common species with Turkey. Also, the number of common species reported from Kerman province to Turkey is higher than other neighbouring countries. The total number of encyrtids species reported from Iran including the present study increased to 179 species in 53 genera and from Kerman province including the present study increased to 29 species.

Table 1 The species of Encyrtidae and their hosts in Kerman province.

Species	Hosts	Reference
<i>Adelencyrtus intersectus</i> (Fonscolombe, 1832)	Soft scale	Yazdani and Rajabi (1993)
<i>Anagyrus aligarhensis</i> Agarwal and Alam, 1959	<i>Phenacoccus</i> sp.	Abd-Rabou <i>et al.</i> (2011)
<i>Blastothrix sericea</i> (Dalman, 1820)	<i>Anapulvinaria pistaciae</i> (Bodenheimer, 1926)	Yazdani and Rajabi (1993)
<i>Cerapterocerus mirabilis</i> Westwood, 1833	<i>Aonidiella orientalis</i> Newstead, 1894	Chodjai (1968, 1989); Davatchi and Chodjai (1969); Behdad (1991); Modarres Awal (1997)
<i>Cerchysiella</i> sp.	Unknown	Present study
<i>Charitopus manukyani</i> Sakhnov, 1993	Unknown	Present study
<i>Cheiloneurus pistaciae</i> Manickavasagam and Mehmejad, 2008	<i>Chelonus kermakiae</i> (Tobias, 2001)	Manickavasagam <i>et al.</i> (2008)
<i>Cheiloneurus submuticus</i> Thomson, 1876	Unknown	Present study
<i>Eupoecilopoda perpunctata</i> (Masi, 1942)	Chrysopidae	Yazdani and Mehrnejad (1993)
<i>Habrolepis dalmanni</i> (Westwood, 1837)	<i>Melanaspis inopinata</i> (Leonardi, 1913)	Yazdani and Rajabi (1993)
<i>Homalotylus flaminus</i> (Dalman, 1820)	Unknown	Present study
<i>Homalotylus nigricornis</i> Mercet, 1921	Unknown	Present study
<i>Homalotylus</i> near <i>platynaspidis</i> Hoffer, 1963	Unknown	Present study
<i>Homalotylus quaylei</i> Timberlake, 1919	Unknown	Present study
<i>Leptomastix flava</i> Mercet, 1921	<i>Eulecanium rugulosum</i> (Archangelskaya, 1937)	Yazdani and Rajabi (1993)
<i>Mayridia myrlea</i> (Walker, 1838)	Unknown	Present study
<i>Ooencyrtus egeria</i> Huang and Noyes, 1994	<i>Brachynema germarii</i> (Kolenati, 1846), <i>Acrosternum arabicum</i> Wagner, 1959 and <i>A. breviceps</i> (Jakovlev, 1889)	Mohammadpour <i>et al.</i> (2016)
<i>Ooencyrtus masii</i> (Mercet, 1921)	Unknown	Present study
<i>Ooencyrtus ocnariae</i> Hayat and Mehmejad, 2018	<i>Ocnaria terebinthina</i> Strg.	Hayat and Mehrnejad (2018)
<i>Ooencyrtus pistaciae</i> Hayat and Mehrnejad, 2016	<i>Brachynema germarii</i>	Hayat and Mehrnejad (2016)
<i>Ooencyrtus pityocampae</i> (Mercet, 1921)	<i>Brachynema signatum</i>	Mohammadpour <i>et al.</i> (2013, 2014, 2015)
<i>Ooencyrtus telenomicida</i> (Vassiliev, 1904)	<i>Brachynema</i> spp., <i>Acrosternum arabicum</i> and <i>A. breviceps</i>	Hashemi Rad <i>et al.</i> (2002); Mohammadpour <i>et al.</i> (2016)
<i>Prionomitus tiliaris</i> (Dalman, 1820)	Unknown	Present study
<i>Prochiloneurus bolivari</i> Mercet, 1919	Unknown	Abd-Rabou <i>et al.</i> (2011)
<i>Psyllaephagus claripes</i> Trjapitzin, 1967	Unknown	Present study
<i>Psyllaephagus pistaciae</i> Ferrière, 1961	<i>Agonoscena pistaciae</i> Burckhardt and Lauterer, 1989	Ebrahimi <i>et al.</i> (1998); Emami and Mehrnejad (2002, 2004, 2006); Mehrnejad (2002a, b, c, 2008); Mehrnejad and Copland (2006, 2007)
<i>Psyllaephagus procerus</i> (Mercet, 1921)	<i>Diaphorina citri</i> Kuwayama, 1908	(Moghbeli-Gharaei <i>et al.</i> (2013)
<i>Psyllaphycus diaphorinae</i> Hayat, 1972	<i>Agonoscena pistaciae</i>	Mehrnejad and Emami (2005)
<i>Syrphophagus aphidivorus</i> (Mayr, 1876)	Hyperparasitoid of <i>Agonoscena pistaciae</i>	Yazdani and Mehrnejad (1993); Ebrahimi <i>et al.</i> (1998); Mehrnejad (2002b); Emami and Mehrnejad (2002, 2004, 2006); Mehrnejad and Emami (2005)

Acknowledgments

This research was supported by Shahid Bahonar University of Kerman and East-Azarbaijan

Agricultural and Natural Resources Research and Education Center, which is greatly appreciated. Our special thanks are expressed to Mr. M. Purrezaali and Mrs. M. Changizi for

helping us in sample collection. The authors are grateful to anonymous reviewers for their valuable comments and recommendations on the earlier version of this paper.

References

- Abd-Rabou, Sh., Ghahari, H., Hedqvist, K. J. and Ostovan, H. 2011. Parasitoid-host relationship between *Comperiella bifasciata* Howard (Hymenoptera: Encyrtidae) and *Aonidiella orientalis* (Newstead) (Hemiptera: Diaspididae) with a list of Iranian Encyrtidae. *Journal of Entomological Research*, 35(1): 1-8.
- Asadeh, Gh. A. and Mossadegh, M. S. 1991. An investigation of the mealybug (*Pseudococcus* spp.) natural enemies fauna in the Khuzestan's province. *Proceedings of the 10th Iranian Plant Protection Congress*, Kerman, 1-5 September 1991. University of Shahid Bahonar, Kerman, pp. 3.
- Bauer, L. S., Liu, H.P., Miller, D. L., and Gould, J. 2008. Developing a classical biological control program for *Agrilus planipennis* (Coleoptera: Buprestidae), an invasive ash pest in North America. *Newsletter of the Michigan Entomological Society*, 53: 38-39. Available from <http://www.nrs.fs.fed.us/pubs/1439> [accessed 11 August 2014].
- Behdad, E. 1991. *Pests of Fruit Crop in Iran*. 2nd Edition. Neshat Publishing, Esfahan, 822 pp.
- Chodjai, M. 1968. Résultats de l'étude faunistique des Hyménoptères parasites (Terebrants) en Iran et l'importance de leur utilisation dans la lutte biologique. *Proceedings of the first Iranian Plant Protection Congress*, 13-18 September 1968. Tehran University, Karaj, pp. 25-35.
- Chodjai, M. 1989. *Entomology (Ethology, Social Life and Natural Enemies) (Biological Control)*. Vol. III. 2 Edition. Tehran University Publishing, Tehran, 406 pp.
- Dahlsten, D. L., Daane, K. M., Paine, T. D., Sime, K. R., Lawson, A. B., Rowney, D. L., Roltsch, W. J., Andrews, J. W., Kabashima, J. N., Shaw, D. A., Robb, K. L., Gelsel, P. M., Chaney, W. E., Ingels, C. A., Varela, L. G., Bianchi, M. L. and Taylor, G. 2005. Imported parasite wasps helps control red gum lerp psyllid. *California Agriculture*, 59(4): 229-234.
- Davatchi, A. and Chodjai, M. 1969. *Les Hyménoptères entomophages de l'Iran (études faunistiques)*. Vol. 107. Faculty Agriculture, Tehran University Publishing, Tehran, 88 pp.
- Ebrahimi, E., Tavakoli-Korghond, G. R., Mianbandi, K., Mahmoodi, H., Mohammadipour, K. and Noyes, J. 2014. *Ooencyrtus ferdowsii* sp. n. (Hymenoptera: Encyrtidae), an egg parasitoid of *Osphranteria coerulescens* (Coleoptera: Cerambycidae) in Iran. *Zoology in the Middle East*, 61(1): 45-49.
- Ebrahimi, S. H., Shahid Zandi, K. H. and Mehrnejad, M. R. 1998. Population fluctuation of natural enemies of pistachio psylla (*Agonoscaena targionii* Lichtenstein). *Proceedings of the 13th Iranian Plant Protection Congress*, Karaj, 23-27 August 1998. Karaj Junior College of Agriculture, Karaj, pp. 106.
- Emami, S. Y. and Mehrnejad, M. R. 2002. Host selection behavior in three hyperparasitoids of the common pistachio psylla, *Agonoscaena pistaciae*. *Proceedings of the 15th Iranian Plant Protection Congress*, Kermanshah, 7-11 September 2002. Razi University of Kermanshah, Kermanshah, pp. 176.
- Emami, S. Y. and Mehrnejad, M. R. 2004. The situation of weeds aphids parasitoids in pistachio orchards of Kerman province. *Proceedings of the 16th Plant Protection Congress of Iran*, Tabriz 28 August-1 September 2004. Tabriz, Iran, University of Tabriz, pp. 19.
- Emami, S. Y. and Mehrnejad, M. R. 2006. Weed aphids and their parasitoids in pistachio orchards in Kerman Province. *Acta Horticulturae*, 726: 437-439.
- Evans, G. and Abd-Rabou, S. 2013. An annotated list of the encyrtids of Egypt (Hymenoptera: Chalcidoidea: Encyrtidae). *Acta Phytopathologica et Entomologica Hungarica*, 48(1): 107-128.

- Fallahzadeh, M. and Japoshvili, G. 2010. Checklist of Iranian Encyrtids (Hymenoptera: Chalcidoidea) with descriptions of new species. *Journal of Insect Science*, 10: 1-24.
- Fallahzadeh, M. and Japoshvili, G. 2013. Corrections to the list of Encyrtidae (Hymenoptera: Chalcidoidea) from Iran. *Journal of the Entomological Research Society*, 15(2): 117-121.
- Fallahzadeh, M., and Japoshvili, G. 2017. An updated checklist of Iranian Encyrtidae (Hymenoptera: Chalcidoidea). *Zootaxa*, 4344(1): 1-46.
- Fallahzadeh, M., Japoshvili, G. and Guerrieri, E. 2018. Notes and corrections to Guerrieri and Gahari's (2018) list of Iranian Encyrtidae (Hymenoptera: Chalcidoidea) (*Zootaxa*, 4444(3)). *Zootaxa*, 4527(2): 295-296.
- Fallahzadeh, M., Japoshvili, G., Abdmaleki, R. and Saghaei, N. 2014. New records of Tetracneminae (Hymenoptera: Chalcidoidea: Encyrtidae) from Iran. *Turkish Journal of Zoology*, 38: 515-518.
- Fallahzadeh, M., Japoshvili, G., Saghaei, N. and Daane, K. M. 2011. Natural enemies of *Planococcus ficus* (Hemiptera: Pseudococcidae) in Fars province vineyards, Iran. *Biocontrol Science and Technology*, 21(4): 427-433.
- Fallahzadeh, M., Shojaei, M., Ostovan, H. and Kamali, K. 2006. The first report of two parasitoid wasps on the larvae of *Nephus bipunctatus* (Coleoptera: Coccinellidae) from Iran. *Journal of Entomological Society of Iran*, 26(1): 95-96.
- Feli kohikheili, Z., Damavandian, M. R., Barimani varandi, H. and Barari, H. 2016. First record of 3 species of Parasitoid wasps on Coccoidea from Iran. *Journal of Entomological Society of Iran*. 35(2): 93-96.
- Gahari, H., Abd-Rabou, Sh., Sakenin, H., Hedqvist, K.J. and Ostovan, H. 2010. A contribution to some Chalcidoidea wasps (Hymenoptera) from Iran. *Journal of Biological Control*, 24(1): 17-21.
- Golpayegani, Sh., Talebi, A. A., Lotfalizadeh, H. and Rakhshani, E. 2010. Faunistic study and host range of encyrtid parasitoids (Hymenoptera: Encyrtidae) collected in some parts of Lorestan and Markazi provinces of Iran. *Journal of Entomological Research*, 1(4): 315-324.
- Grissell, E. E. and Schauff, M. E. 1990. A handbook of the families of Nearctic Chalcidoidea (Hymenoptera). *Entomological Society of Washington Handbook No 1*: 85 pp.
- Guerrieri, E. and Noyes, J. S. 2000. Revision of European species of genus *Metaphycus* Mercet (Hymenoptera: Chalcidoidea: Encyrtidae), parasitoids of scale insects. *Systematic Entomology*, 25: 147-222.
- Guerrieri, E. and Noyes, J. S. 2005. Revision of the European species of *Copidosoma* Ratzeburg (Hymenoptera: Encyrtidae), parasitoids of caterpillars (Lepidoptera). *Systematic Entomology*, 30(1): 97-174.
- Guerrieri, E. and Noyes, J. S. 2009. A review of the European species of the genus *Trechnites* Thompson (Hymenoptera: Chalcidoidea: Encyrtidae), parasitoids of plant lice (Hemiptera: Psylloidea) with description of a new species. *Systematic Entomology*, 34: 252-259.
- Guerrieri, E. and Viggiani, G. 2005. A review of the encyrtid (Hymenoptera: Chalcidoidea, Encyrtidae) parasitoids of Dryinidae (Hymenoptera: Chrysidoidea) with description of *Cheiloneurus olmii* sp. nov. *Systematics and Biodiversity*, 2(3): 305-317.
- Hashemi Rad, H., Soleyman Najadian, E. and Radjabi, G. 2002. Population fluctuation of green stink bugs (*Brachynema* spp. and *Acrosternum* spp.) and their egg parasitoids in Rafsanjan. *Proceedings of the 15th Iranian Plant Protection Congress*, Kermanshah, 7-11 September 2002. Razi University of Kermanshah, Kermanshah, pp. 179.
- Hayat, M. and Mehrnejad, M. R. 2016. *Ooencyrtus* (Hymenoptera: Encyrtidae), egg parasitoids of the pistachio green stink bugs (Hemiptera: Pentatomidae) in Iran. *Zootaxa*, 4117(2): 198-210.

- Hayat, M. and Mehrnejad, M. R. 2018. A new species of *Ooencyrtus* (Hymenoptera: Encyrtidae), parasitoid in eggs of *Ocneria terebinthina* (Lepidoptera: Lymantriidae) infesting pistachio plants in Iran. *Journal of Insect Systematics*, 5(1 and 2): 14-18.
- Hesami, Sh., Mohammadi Khoramabadi, A., Japoshvili, G. and Seyedebrahimi, S. 2012. *Metaphycus bogdanovikatkovi* (Hymenoptera: Encyrtidae), a new record for the Iranian fauna. *Journal of Entomological Society of Iran*, 33(3): 85-86.
- Hoffer, A. 1963. Descriptions of new species of the family Encyrtidae from Czechoslovakia (Hymenoptera, Chalcidoidea) I. *Sborník Entomologického Oddelení Národního Musea v Praze*, 35:549-592.
- Iranipour, Sh., Kharrazi Pakdel, A., Esmaili, M. and Rajabi, G. 1998. Introduction of a chalcid species from Eupelmidae for Iran. *Proceedings of the 13th Iranian Plant Protection Congress*, Karaj, 23-27 August 1998. Karaj Junior College of Agriculture, Karaj, pp. 5.
- Japoshvili, G. and Fallahzadeh, M. 2018. A new species of *Anagyrus* (Hymenoptera: Encyrtidae) from Iran, parasitoid of *Planococcus ficus* (Hemiptera: Pseudococcidae) on *Vitis vinifera*. *Zoology in the Middle East*, 64(4): 340-344.
- Japoshvili, G., Jalilvand, K., Vahedi, H.A. and Fallahzadeh, M. 2015. *Metaphycus parthenolecanii* sp. n. (Hymenoptera: Encyrtidae), a parasitoid of *Parthenolecanium corni* (Hemiptera: Coccidae) in Iran. *Zoology in the Middle East*, 61(2): 161-165.
- Japoshvili, G., Rakhshani, H. and Khajehali, J. 2016. A new species of *Copidosoma* (Hymenoptera: Encyrtidae) from Iran. *Zoology in the Middle East*, 62(3):261-264 .
- Lotfalizadeh, H. 2010a. The genus *Metaphycus* Mercet, 1917 (Hymenoptera: Encyrtidae) of the Iranian fauna with description of a new species. *North-Western Journal of Zoology*, 6(2): 255-266.
- Lotfalizadeh, H. 2010b. Some new data and corrections on Iranian encyrtid wasp (Hymenoptera: Chalcidoidea, Encyrtidae) fauna. *Biharean Biologist*, 4(2): 173-178.
- Lotfalizadeh, H. and Ebrahimi, E. 2001. New report of *Homalotylus nigricornis* Mercet (Hymenoptera: Encyrtidae) in Iran. *Journal of Entomological Society of Iran*, 21(1): 116.
- Lotfalizadeh, H., and Bab-Morad, M. 2014. *Copidosoma bouchenum* Ratzeburg (Hymenoptera: Encyrtidae): new record from Iran. *Turkish Journal of Zoology*, 39: 185-187.
- Lotfalizadeh, H., Bayegan, Z. A. and Zargaran, M. R. 2016. Species diversity of Chalcidoidea (Hymenoptera) in the rice fields of Iran. *Journal of the Entomological Research Society*, 18(1): 99-111.
- Maafi, S., Rajabi, G. and Jafari, M. E. 1998. Introduction of the natural enemies of *Planococcus citri* in Mazandaran. *Proceedings of the 13th Iranian Plant Protection Congress*, Karaj, 23-27 August 1998. Karaj Junior College of Agriculture, Karaj, pp. 153.
- Manickavasagam, S., Mehrnejad, M. R. and Noyes, J. S. 2008. *Cheiloneurus pistaciae* sp. nov. (Hymenoptera: Encyrtidae) a facultative hyperparasitoid of *Kermania pistaciella* Amsel (Lepidoptera: Tineidae), a pest of pistachio trees in Iran. *Zootaxa*, 1958: 61-64.
- Mehrnejad, M. R. 2002a. Negative agents harming the population of *Psyllaephagus pistaciae*, the parasitoid of the common pistachio psylla. *Proceedings of the 15th Iranian Plant Protection Congress*, Kermanshah, 7-11 September 2002. Razi University of Kermanshah, Kermanshah, pp. 175.
- Mehrnejad, M. R. 2002b. Bionomics of the common pistachio psylla, *Agonoscena pistaciae*, in Iran. *Acta Horticulturae*, 591: 535-539.
- Mehrnejad, M. R. 2002c. Overwintering sites of the parasitoid, *Psyllaephagus pistaciae* and its host, *Agonoscena pistaciae*. (Abstract 077). In: *Abstracts, VIIth European Congress of Entomology*, October 7-13 2002, Thessaloniki, Greece. Hellenic Entomological Society, Athens, pp. 66.
- Mehrnejad, M. R. 2008. Seasonal biology and abundance of *Psyllaephagus pistaciae*

- (Hymenoptera: Encyrtidae), a biocontrol agent of the common pistachio psylla *Agonoscena pistaciae* (Hemiptera: Psylloidea). *Biocontrol Science and Technology*, 18(4): 409-417.
- Mehrnejad, M. R. and Copland, M. J. W. 2006. Biological parameters of parasitoid *Psyllaephagus pistaciae* and its host *Agonoscena pistaciae* in relation to temperature. *Journal of the Entomological Research Society*, 8(1): 1-20.
- Mehrnejad, M. R. and Copland, M. J. W. 2007. Host discrimination by the endoparasitoid *Psyllaephagus pistaciae* (Hymenoptera: Encyrtidae): a case of time-dependent ability. *Biocontrol Science and Technology*, 17(3/4): 401-411.
- Mehrnejad, M. R. and Emami, S. Y. 2005. Parasitoids associated with the common pistachio psylla, *Agonoscena pistaciae*, in Iran. *Biological Control*, 32(3): 385-390.
- Modares Awal, M. 1997. List of Agricultural Pests and Their Natural Enemies in Iran. Revised Edition. Ferdowsi University Press, Mashhad, 429 pp.
- Moghbeli-Gharaei, A., Ziaaddini, M., Jalali, M. A. and Lotfalizadeh, H. 2013. Report of *Psyllaephagus procerus* Mercet (Hymenoptera: Encyrtidae) as parasitoid of the *Diaphorina citri* (Hemiptera: Psyllidae) from Iran. *Applied Entomology and Phytopathology*, 80(2): 185-186.
- Mohammadpour, M., Jalali, M. A., Michaud, J. P., Ziaddini, M. and Hashemirad, H. 2014. Multiparasitism of stink bug eggs: competitive interactions between *Ooencyrtus pityocampae* and *Trissolcus agriope*. *BioControl*, 59(3): 279-286.
- Mohammadpour, M., Ziaddini, M., Jalali, M. A. and Hashemirad, H. 2013. Some biological characteristics of *Ooencyrtus pityocampae*, the egg parasitoid of *Brachynema signatum* in laboratory conditions. *Proceedings of the Conference of Biological Control in Agriculture and Natural Resources*, 27-28 August 2013. University of Tehran, Tehran, pp. 208.
- Mohammadpour, M., Ziaddini, M., Jalali, M. A., Hashemirad, H. and Lotfalizadeh, H. 2015. Identification of the egg parasitoids of green pistachio stink bug, and the effect of diet on adult longevity of the two parasitoids, *Ooencyrtus pityocampae* (Mercet) and *Trissolcus agriope* (Kozlov and Lê). *Journal of Entomological Research*, 7(3): 277-287.
- Mohammadpour, M., Ziaddini, M., Jalali, M. A., Hashemirad, H. and Mohammadi-Khoramabadi, A. 2016. Egg parasitoids of the pistachio green stink bug, *Brachynema germari* (Hemiptera: Pentatomidae) in Kerman province, Iran. *Zoology and Ecology*, 26(1): 28-34.
- Moravvej, S. A., Lotfalizadeh, H. and Shishehbor, P. 2018. A contribution to the study of Encyrtidae (Hymenoptera: Chalcidoidea) of Khuzestan in southwestern Iran. *Journal of Insect Biodiversity and Systematics*, 4(1): 13-23.
- Noori, H., Asgari, S., Moslehi-Rad, H. 2003. Study and identification on *Eurygaster integriceps* Puton egg parasitoids in Ghazvin Province. *Journal of Agricultural Sciences, Islamic Azad University*, 9(2): 141-150.
- Novin, M. 2000. The biology and population fluctuations of *Nipaecoccus viridis* (Newstead) and its natural enemies in citrus orchards of Dezful. M. Sc. thesis of Entomology, Shahid Chamran University, Ahwaz, 152 pp.
- Novin, M., Mossadegh, M. S., Karami Nejad, M. and Ghasemi Nejad, M. 2000. Natural enemy of *Nipaecoccus viridis* in the North of Khuzestan. *Proceedings of the 14th Iranian Plant Protection Congress*, Isfahan, 5-8 September 2000. Isfahan University of Technology, Isfahan, pp. 264.
- Noyes, J. S. 1982. Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History*, 16: 315-334.
- Noyes, J. S. 2020. Universal Chalcidoidea Database. The Natural History Museum, London. Available from: <http://www.nhm.ac.uk/chalcidoids> (Accessed 20th March 2020).

- Noyes, J. S. and Hayat, M. 1984. A review of the genera of Indo-Pacific Encyrtidae (Hymenoptera: Chalcidoidea). Bulletin of the British Museum (Natural History) (Entomology), 48: 131-395.
- Noyes, J. S., Woolley, J. B. and Zolnerowich, G. 1997. Chapter 8, Encyrtidae. In: Gibson, G. A. P., Huber, J. T and Woolley, J. B. (Eds.), Annotated keys of the genera of Nearctic Chalcidoidea (Hymenoptera). pp. 170-320. National Research Council Research Press, Ottawa, Canada.
- OILB. 1971. Liste d'identification des entomophages. Vol. 8. OILB, Genève, 64 pp. [Ent. Library-Bound with volume 11 of Entomophaga].
- Omid, R., Azizkhani, E., Askary, H. and Moniri, V. R. 2005. Biological investigation of *Ocneria terebinthina* in Khojir Protected Area. Iranian Journal of Forest and Range Protection Research, 2(2): 209-211.
- Radjabi, G. and Amir-Nazari, M. 1989. Egg parasites of the sunn pest in the central part of the Iranian plateau. Applied of Entomology and Phytopathology, 56(1, 2): 1-12.
- Rajabi Mazhar, N. and Sadeghi, S. E. 2006. Studying biology of *Psyllopsiopsis repens* Loginova and identification of its natural enemies in Hamadan Province. Proceedings of the 17th Iranian Plant Protection Congress, Karaj, 2-5 September 2006. Campus of Agriculture and Natural Resources, University of Tehran, Karaj, pp. 236.
- Rajabi Mazhar, N., Sadeghi, S. E. and Yarmand, H. 2007. Report of *Psyllaephagus clarips* (Hymenoptera: Encyrtidae) on *Psyllaephagus repens* in Iran. Iranian Journal of Forest and Range Protection Research, 1(2): 211.
- Rastegar, J., Sakenin, H. and Ghods, P. 2012. A study of the Encyrtidae (Hymenoptera: Chalcidoidea) from Iran. Calodema, 224: 1-3.
- Sabahi, Q., Kharrazi Pakdel, A. and Esmaili, M. 1999. Study on the biology of *Ooencyrtus* cf. *masii* (Hymenoptera: Encyrtidae), a new egg parasitoid of pistachio defoliator *Ocneria terebinthina* (Lepidoptera: Lymantriidae). Journal of Entomological Society of Iran, 18(1, 2): 1-5.
- Sabahi, Q., Kharrazi Pakdel, A., Esmaili, M. and Abaii, M. 1998. Studies on the biology of *Ooencyrtus* cf. *masii* (Hymenoptera: Encyrtidae), an egg parasitoid of pistachio defoliator *Ocneria terebinthina* Staudinger (Lepidoptera: Lymantriidae) in Fars province. Proceedings of the 13th Iranian Plant Protection Congress, Karaj, 23-27 August 1998. Karaj Junior College of Agriculture, Karaj, pp. 138.
- Yazdani, A. and Mehrnejad, M. R. 1993. The first record of one psyllid species and several Hymenoptera parasitoids on pistachio psyllid from Iran. Proceedings of the 11th Plant Protection Congress of Iran, Rasht, 28 August-2 September 1993. University of Guilan, Rasht, pp. 210.
- Yazdani, A. and Rajabi, A. 1993. The first record of Hymenoptera parasitoids of scale insects on pistachio in Iran. Proceedings of the 11th Plant Protection Congress of Iran, Rasht, 28 August-2 September 1993. University of Guilan, Rasht, pp. 212.
- Yousefi, M., Vahedi, H. A., Ebrahimi, E. and Sharifi, R. 2018. Identification of parasitoid wasps of Scale insects (Chalcidoidea) in fruit orchards of Markazi province. Applied Entomology and Phytopathology, 85(2): 251-264.
- Zhang, Y. and Huang, D. 2004. A review and an illustrated key to genera of Encyrtidae (Hymenoptera: Chalcidoidea) from China. Science Press, Beijing, China. 166 pp.

زنبورهای خانواده Encyrtidae (Hymenoptera: Chalcidoidea) در نواحی جنوبی استان کرمان به همراه گزارش جدید چند جنس و گونه

مهلا شجاعی^۱، محمد خیراندیش^{۱*}، سیدمسعود مجدزاده^۲ و حسین لطفعلی زاده^۳

۱- گروه گیاه پزشکی، دانشکده کشاورزی، دانشگاه شهید باهنر کرمان، کرمان، ایران.

۲- گروه زیست شناسی، دانشکده علوم، دانشگاه شهید باهنر کرمان، کرمان، ایران.

۳- بخش تحقیقات گیاه پزشکی، مرکز تحقیقات کشاورزی و منابع طبیعی استان آذربایجان شرقی، تبریز، ایران.

پست الکترونیکی نویسنده مسئول مکاتبه: m.khayrandish@uk.ac.ir

دریافت: ۱۴ فروردین ۱۳۹۹؛ پذیرش: ۱۲ تیر ۱۳۹۹

چکیده: در این پژوهش، زنبورهای پارازیتوئید خانواده Encyrtidae (Hymenoptera: Chalcidoidea) از مناطق جنوبی استان کرمان جمع آوری و مورد بررسی قرار گرفته است. نمونه ها با استفاده از تله مالیز، طی ماه های فروردین تا شهریور ۱۳۹۶ جمع آوری گردید. در مجموع، ۱۶ گونه متعلق به ۱۰ جنس از خانواده Encyrtidae جمع آوری و شناسایی شد. جنس های *Charitopus* و *Cerchysiella* Girault, 1914 *Chritopus manukyani* Foerster, 1856 برای فون ایران جدید می باشند. هم چنین چهار گونه *Chritopus manukyani* Foerster, 1856، *Cheiloneurus submuticus* (Walker, 1838)، *Sakhnov*, 1993 و *Mayridia myrlea* Thomson, 1876، *Prionomitus tiliaris* (Dalman, 1820) برای فون ایران گزارش جدید می باشند. خصوصیات مرفولوژیک برای گونه های جدید و خصوصیات زیست شناسی و پراکنش جغرافیایی برای همه گونه ها ذکر شده است. هم چنین لیست به روز شده از گونه های خانواده Encyrtidae از استان کرمان ارائه شده است.

واژگان کلیدی: encyrtid wasp، گزارش جدید، ایران، فون، استان کرمان