

Research Article

First record of the subfamily Lycorininae (Hymenoptera: Ichneumonidae) from Iran

Ali Ameri^{1*}, Ebrahim Ebrahimi¹, Aliasghar Talebi² and Pascal Rousse³

1. Insect Taxonomy Research Department, Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

2. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

3. Maison de la Bio, 55 chemin de la Castelette BP 11283, Avignon Cédex 9, France.

Abstract: The present study is based on the material collected from East Azerbaijan province (Northwestern Iran) between July-August 2019. The specimens were collected using Malaise traps from a variety of habitats. *Lycorina triangulifera* Holmgren, 1859 were collected and identified. This species is the first record of the subfamily Lycorininae Cushman and Rohwer, 1920 (Hymenoptera: Ichneumonidae) in Iran. Morphological descriptions, as well as detailed illustrations of this species, are presented.

Keywords: Lycorininae, new record, distribution, Iran

Introduction

Lycorininae Cushman and Rohwer, 1920 is a small subfamily of Ichneumonidae, encompassing the single genus *Lycorina* Holmgren 1859 and 34 described species distributed in almost all biogeographical regions (Yu *et al.* 2016). The subfamily can be easily distinguished by central triangular areas delimited by deep lateral grooves on tergites 2–4 and the expanded anterolateral corner of the propodeum engaging the metanotum with a small hook (Gauld *et al.*, 1997; Quicke *et al.*, 2009). *Lycorina* species are koinobiont ectoparasitoids of various concealed micro-sized lepidopteran caterpillars (Coronado-Rivera *et al.* 2004; Shaw, 2004). Some species have been reared from Crambidae in leaf rolls or webs (Doerksen and Neunzig, 1974; Finlayson, 1976) and Tortricidae (Chao, 1980; Shaw, 2004).

Faunal studies on the family Ichneumonidae have recently been carried out in various parts of Iran (Amiri *et al.*, 2015a, b, 2016a, b, 2017; Barahoei *et al.*, 2012; Hooshyar *et al.* 2014; Mohammadi-Khoramabadi, 2015; Mohammadi-Khoramabadi and Talebi, 2013; Mohammadi-Khoramabadi *et al.*, 2011, 2013a, b; Riedel *et al.*, 2019; Zardoue Heydari *et al.*, 2020a, b, c, d) which led to the identification of about 580 species and 27 subfamilies. *Lycorina triangulifera* is recorded for the first time from Iran in the present study.

Materials and Methods

The examined specimens were collected from the Arasbaran protected area of East Azerbaijan province, northwestern Iran, bearing a high mountainous region ranging from 256 m to 2896 m above sea level. Arasbaran encompasses part of the Caucasus mountains with diverse natural landscapes, including highlands, steep valleys, high and steep mountainsides, forests, rivers, and agricultural rangelands (Mohammadi Fazel, 2020) (Fig. 1A-D). The specimens were collected using

Handling Editor: Ehsan Rakhshani

* Corresponding author: ameri@iripp.ir

Received: 14 September 2020, Accepted: 28 March 2021

Published online: 02 April 2021

Malaise traps. The captured specimens were extracted from the collecting jars, then treated with a mixture of ethanol(60%) /xylene (40%) for two days, followed by amyl acetate for two days (AXA), and finally placed on filter paper for drying (van Achterberg, 2009). Photographs were taken using a stereomicroscope (Olympus SZH, Tokyo, Japan) equipped with a 650D Canon digital camera. Most morphological terms are defined on the HymaToL website (<http://www.hymatol.org>).

The identifications were mainly carried out using reliable keys (Chao, 1980; Gauld, 1984; Gauld *et al.*, 1997; Kasparyan, 2007; Rousse and van Noort, 2013, 2014; Choi *et al.*, 2014). All examined specimens are deposited in the Insect Taxonomy Research Department of the Iranian Research Institute of Plant Protection(Hayk Mirzayans Insect Museum, HMIM). The geographical and biological records are extracted from the Taxapad (Yu *et al.*, 2016).



Figure 1 Field sampling with malaise traps in Iran East Azerbaijan Province. A) Orchards, Kaleybar; B) General view, Aynalo; C) Forest edge, Arasbaran Protected Area; D) Forests and pastures, Kaleybar.

Results

Two adults (female) of *L. triangulifera* were collected and identified. A morphological description is provided here.

Lycorina triangulifera Holmgren, 1859

Female: Body length 7.2 mm, forewing 6.0 mm, ovipositor 3.0 mm. Face moderately densely punctured, 1.3x wider than the high, central part of face convex in lateral view; clypeus moderately flat, with the round apical edge, separated from face by a transverse groove; malar space equal to the width of the mandible (Fig. 2A, D-F); frons

with a longitudinal carina; scape 1.7x wider than the first flagellomere; flagellum with 29 flagellomeres, first flagellomere, 4.5 longer than wide, apical flagellomere square. Mesoscutum 1.3x longer than wide, mesonotum sparsely punctate; scutellum carinate and convex; notaulus shallow; epicnemial carina incomplete (Fig. 2F) Metasoma depressed, terga dorsally closely and coarse punctation; raised central triangular areas large and strongly transverse (Fig. 2 B-C). Ovipositor sheath 1.7x as long as the hind tibia. Forewing with open areolet, 2m-cu with one bulla, pterostigma broadly triangular, marginal

cell moderately long, hindwing with 1/Rs longer than r-m (Fig. 3A-B). Body black with the following exceptions: Clypeus, posterior orbits, and two spots on the face and hind part of scutellum yellow. Ovipositor and flagellum ventrally brown, the hind tibia and ovipositor sheath dark brown, hind trochantellus to femur reddish brown.

Male: Body length 7.0 mm, fore wing 6.0 mm, flagellum with 34 segments, all coxae and trochanters black; fore, mid, hind femora, tibia and tarsus dark brown (Fig. 2 D, B).

Material examined: Iran, East Azerbaijan Province, Arasbaran Protected Area (Kalaybar, Aynalo) July 16, 2019 (38°54'10.62"N 46°47'0.74"E), 1263 m, August, 1-19, 2019, Malaise trap, 1♂ 1♀, Leg.: A. Ameri.

Distribution: Austria, Azerbaijan, Belarus, Belgium, Bulgaria, China, Czech Republic, Denmark, Estonia, Finland, France, Germany, Hungary, Ireland, Italy, Japan, Korea, Lithuania, Moldova, Mongolia, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Sweden, Ukraine, United Kingdom (Yu *et al.*, 2016), Iran (New record).



Figure 2 *Lycorina triangulifera*, A) General habitus, female, lateral view; B) Metasoma, male, dorsal view; C) Metasoma, female, dorsal view; D) General habitus, male, lateral view; E) Head, frontal view; F) Head and Mesosoma, lateral view.



Figure 3 *Lycorina triangulifera*; A) Forewing, B) Hindwing.

Discussion

This species is the first record of the subfamily in Iran, bringing to 27 the number of recorded ichneumonid subfamilies (Barahoei *et al.* 2012; Hooshyar *et al.*, 2014; Mohammadi-Khoramabadi, 2013, 2015; Riedel *et al.*, 2019). Among neighboring countries, this species was so far known only from Azerbaijan (Abdinbekova, 1963).

The specimens collected in this study were captured using Malaise traps not to add host records. Some species of Lycorininae are known as parasitoids of Gelechiidae, Cerambycidae, and Tortricidae (Yu *et al.*, 2016). Owing to its host range, this species is a parasitoid of some important agricultural pests and can be considered as a most potential biological control agent of *Yponomeuta malinella* Zeller, 1838 and *Archips rosana* (Linnaeus, 1758), which are important

pests in orchard plants in Iran (Hatmaker, 1985; Alizadeh and Shayan, 2017).

Comparing the number of recorded species (current study) of the subfamily Lycorininae in Iran to some countries shows that more investigations are needed to obtain better knowledge. The number of Lycorininae is probably higher than presently known since several species from some countries can be expected to occur also in Iran (Yu *et al.*, 2016). Most of the collections are made in the Arasbaran protected area of Iran, and the inadequate sampling coverage of southern and central parts of the country indicates that the number of Iranian species of Lycorininae could increase.

Acknowledgments

Thanks to the Iranian Research Institute of Plant Protection for providing financial support of this

research project under code number 2-16-16-047-990334. The authors thank the Department of Environment, Islamic Republic of Iran, for cooperation and permission for sampling in the protected areas.

References

- Abdinbekova, A. A. 1963. The Ichneumonid fauna of the Kuba-Khachmassk zone of Azerbaidzhan. *Izvestiya Akademii Nauk Azerbaidzhanskoi SSR*, 1963(5): 43-51.
- Alizadeh, P. and Shayan, A. 2017. Executive instruction of citrus leafroller, *Archips rosanus* (Linnaeus) (Lep: Tortricidae). Office of prediction and Pest Control, Plant Protection Organization, Tehran, Iran, 4p. (In Persian).
- Amiri, A., Talebi, A. A., Jussila, R., Rakhshani, E. and Hajiqanbar, H. 2015a. A study of the Iranian Cremastinae (Hymenoptera: Ichneumonidae). *Journal of Insect Biodiversity and Systematics*, 1(2): 87-100.
- Amiri, A., Talebi, A. A., Rakhshani, E. and Hajiqanbar, H. 2017. First report of the genus *Cymodusa* (Ichneumonidae: Campopleginae) from Iran. *Journal of Insect Biodiversity and Systematics*, 3(2): 81-89.
- Amiri, A., Talebi, A. A., Riedel, M., Rakhshani, E. and Hajiqanbar, H. 2015b. A survey of Metopiinae (Hymenoptera: Ichneumonidae) in southern Iran, with three new records. *Journal of Crop Protection*, 4(4): 519-531.
- Amiri, A., Talebi, A., Jussila, R., Rakhshani, E. and Hajiqanbar, H. 2016a. Study of the subfamily Ophioninae (Hymenoptera: Ichneumonidae) in southern Iran. *Journal of Entomological Society of Iran*, 35(4): 53-67.
- Amiri, A., Talebi, A., Rey Del Castillo, C., Rakhshani, E. and Hajiqanbar, H. 2016b. Study of the genus *Lissonota* (Hymenoptera: Ichneumonidae: Banchinae) in southern Iran. *Journal of Entomological Society of Iran*, 36(2): 89-99.
- Barahoei, H., Rakhshani, E. and Riedel, M. 2012. A checklist of Ichneumonidae (Hymenoptera: Ichneumonoidea) from Iran. *Iranian Journal of Animal Biosystematics*, 8(2): 83-133.
- Chao, H. F. 1980. Description of *Lycorina spilonotae* Chao, sp. nov., with notes on its final-instar larvae (Hymenoptera: Ichneumonidae: Lycorinae). *Entomotaxonomia*, 2: 165-168. [In Chinese].
- Choi, J. K., Cha, J. Y. and Lee, J. W. 2014. Taxonomic study of the subfamily Lycorininae (Hymenoptera: Ichneumonidae) from Korea. *Animal Systematics, Evolution and Diversity*, 30(1): 49-54.
- Coronado-Rivera, J., Gonzalez Herrera, A., Gauld, I. D. and Hanson, P. 2004. The enigmatic biology of the Ichneumonid subfamily Lycorininae. *Journal of Hymenoptera Research*, 13: 223-227.
- Cushman, R. A. and Rohwer S. A. 1920. Holarctic tribes of the Ichneumon-flies of the subfamily Ichneumoninae (Pimplinae). *Proceedings of the United States National Museum* 57: 379-396.
- Doerksen, G. P. and Neunzig, H. H. 1974. The biology of the apple leaf skeletonizer, *Psorosina hammondi*, on *Crateagus* (Lepidoptera: Pyralidae: Phycitinae). *Annals of the Entomological Society of America*, 67: 146-147.
- Finlayson, T. 1976. Cephalic structures and spiracles of finalinstar larvae of the genus *Toxophoroides* (Hymenoptera: Ichneumonidae: Lycorininae). *Canadian Entomologist*, 108: 981-984.
- Gauld, I. D. 1984. The Pimplinae, Xoridinae, Acaenitinae and Lycorinae (Hymenoptera: Ichneumonidae) of Australia. *Bulletin of the British Museum (Natural History), Entomology series*, 49: 235-239.
- Gauld, I. D., Wahl, D., Bradshaw, K., Hanson, P. and Ward, S. 1997. The Ichneumonidae of Costa Rica, 2. Introduction and keys to species of the smaller subfamilies, Anomaloninae, Ctenopelmatinae, Diplazontinae, Lycorininae, Phrudinae, Tryphoninae (excluding Netelia), and Xoridinae, with an appendices on the Rhyssinae. *Memoirs of the American Entomological Institute*, 57: 1-485.

- Hatmaker, E. D. 1985. Data Sheet on *Yponomeuta malinellus*. USDA-APHIS-PPQ-CPHST-PERAL, New Pest Advisory Group.
- Hooshyar, H., Vafaie-Shoushtari, R. and Barimany-Varandy, H. 2014. First report of *Stilbops vetulus* (Hym.: Ichneumonidae: Stilbopinae) from Iran. Journal of Entomological Society of Iran, 34: 103-104.
- Kasparyan, D. R. and Khalaim, A. I. 2007. Pimplinae, Tryphoninae, Eucerotinae, Xoridinae, Agriotypinae, Lycorininae, Neorhacodinae, Ctenopelmatinae, Phrudinae, Ophioninae, Acaenitinae, Collyriinae, Mesochorinae. In: Lelej, A. S. (Ed.), Key to the Insects of Russia Far East. Vol. IV. Neuropteroidea, Mecoptera, Hymenoptera. Pt. 5, (Ed., Lelej AS). Dalnauka, Vladivostok, pp. 1-1052 (in Russian).
- Mohammadi Fazel, A. 2020. Protected Areas of the East Azerbaijan Province of Iran, 2nd Edition of the Atlas of Protected Areas of Iran, 37 pp.
- Mohammadi-Khoramabadi, A. 2015. First records of the subfamily Adelognathinae (Hym.: Ichneumonidae) from Iran, Biharean Biologist, 9: 164-165.
- Mohammadi-Khoramabadi, A. and Talebi, A. A. 2013. A study of the genus *Orthocentrus* (Hymenoptera: Ichneumonidae, Orthocentrinae) in Gilan and Tehran provinces of Iran, with first records of seven species and one subspecies, Applied Entomology and Phytopathology, 80(2): 29-39.
- Mohammadi-Khoramabadi, A., Talebi, A. A. and Farahani, S. 2011. *Hybrizon buccatus* (de Brebisson, 1825), the first record of the subfamily Hybrizontinae (Hymenoptera: Ichneumonidae) from northern Iran. Biharean Biologist, 5(2): 162-163.
- Mohammadi-Khoramabadi, A., Talebi, A. A. and Zwakhals, K. 2013a. A study of the subfamily Pimplinae (Hymenoptera: Ichneumonidae) in the north of Iran, with eleven new species records. Entomofauna, 34(2): 29-56.
- Mohammadi-Khoramabadi, A., Talebi, A. A. and Zwakhals, K. 2013b. Study on Diplazontinae (Hymenoptera: Ichneumonidae) in the north central of Iran. Journal of Crop Protection, 2(3): 241-261.
- Quicke, D. L. J., Laurenne, N. M., Fitton, M. G. and Broad, G. R. 2009. A thousand and one wasps: a 28S rDNA and morphological phylogeny of the Ichneumonidae (Insecta: Hymenoptera) with an investigation into alignment parameter space and elision. Journal of Natural History 43(23-24): 1305-1421.
- Riedel, M., Ameri, A., Talebi, A. A. and Ebrahimi, E. 2019. Contribution to the Ichneumonidae (Hymenoptera) of Iran, with descriptions of seven new species. Linzer Biologische Beiträge, 51 (1): 361-390.
- Rousse, P. and van Noort, S. 2013. Revision of the Afrotropical Lycorininae (Ichneumonidae; Hymenoptera), with description of a new species from South Africa. Zootaxa, 3666: 252-266.
- Rousse, P. and van Noort, S. 2014. Revision of the Afrotropical Lycorininae (Ichneumonidae; Hymenoptera) II. Three new *Lycorina* species and additional distribution records. Zootaxa, 3884: 222-234.
- Shaw, M. R. 2004. Notes on the biology of *Lycorina triangulifera* Holmgren (Hymenoptera: Ichneumonidae: Lycorininae). Journal of Hymenoptera Research, 13(2): 302-308.
- van Achterberg, C. 2009. Can Townes type Malaise traps be improved some recent developments. Entomologische Berichten Amsterdam, 69(4): 129-135.
- Yu, D. S., Van Achterberg, K. and Horstmann, K. 2016. Taxapad 2016. World Ichneumonoidea 2015. Taxonomy, Biology, Morphology and Distribution Ontario: Nepean, Canada: On USB Flash Drive.
- Zardouei Heydari, M. Z., Rakhshani, E., Rousse, P., Mokhtari, A. and Schwarz, M. 2020a. A new species of *Pristomerus* Curtis, 1836 (Hymenoptera, Ichneumonidae, Cremastinae) and key to the known species in Iran. Zootaxa, 4809(1): 187-194.
- Zardouei Heydari, M., Rakhshani, E., Mokhtari, A. and Schwarz, M. 2020b. The genus *Anomalon* Panzer (Hymenoptera:

- Ichneumonidae: Anomaloninae) of Iran, with description of a new species. *European Journal of Taxonomy* 662: 1-19.
- Zardouei Heydari, M., Rakhshani, E., Mokhtari, A. and Schwarz, M. 2020c. New data on the genus *Latibulus* Gistel, 1848 (Hym., Ichneumonidae, Cryptinae, Cryptini) in Iran. *Journal of Insect Biodiversity and Systematics*, 6(2),: 157-165.
- Zardouei Heydari, M., Rakhshani, E., Mokhtari, A. and Schwarz, M. 2020d. A faunistic overview of *Temelucha* Förster (Hym., Ichneumonidae, Cremastinae) in Iran. *Turkish Journal of Zoology*, 44: 355-374.

اولین گزارش از زیرخانواده (Lycorininae (Hymenoptera: Ichneumonidae) از ایران

علی عامری^{۱*}، ابراهیم ابراهیمی^۱، علی اصغر طالبی^۲ و پاسکال روزه^۳

۱- بخش تحقیقات رده‌بندی حشرات، مؤسسه تحقیقات گیاه‌پزشکی، سازمان تحقیقات آموزش و ترویج کشاورزی، تهران، ایران.

۲- گروه حشره‌شناسی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران.

۳- Maison de la Bio, 55 chemin de la Castelette BP 11283, Avignon Cédex 9, France

پست الکترونیکی نویسنده مسئول مکاتبه: ameri@iripp.ir

دریافت: ۲۴ شهریور ۱۳۹۹؛ پذیرش: ۸ فروردین ۱۴۰۰

چکیده: مطالعه حاضر براساس نمونه‌های جمع‌آوری شده از استان آذربایجان شرقی (شمال غرب ایران) طی ماه‌های مرداد و شهریور سال ۱۳۹۸ انجام گردیده است. نمونه‌برداری با استفاده از تله مالیز در زیستگاه‌های مختلف انجام و گونه *Lycorina triangulifera* Holmgren, 1859 جمع‌آوری و شناسایی گردید. این اولین گزارش از زیرخانواده (Lycorininae (Hymenoptera: Ichneumonidae) از ایران می‌باشد. خصوصیات شکل‌شناسی به‌همراه جزئیات اشکال ارائه شده است.

واژگان کلیدی: Lycorininae، گزارش جدید، پراکنش، ایران