

Research Article

Contribution to the knowledge of Iranian Opiinae (Hymenoptera: Braconidae) with four new records

Sana Dolati¹, Ali Asghar Talebi¹ and Samira Farahani²

1. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, I. R. Iran.

2. Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

Abstract: A survey on the fauna of Opiinae (Hymenoptera: Braconidae) was conducted in the north-central region (2010-2011) and Khuzestan Province, southwestern Iran (2016–2017). Specimens were collected using Malaise traps, leading to the identification of 32 species. Among these, *Apodesmia ocellata* (Wesmael, 1835), *Opiostomus* (*Opiostomus*) *griffithsi* (Fischer, 1962), and *Xynobius* (*Xynobius*) *comatus* (Wesmael, 1835) and *Psytalia* (*Psytalia*) *romani* (Fahringer, 1935) are recorded for the first time in Iran. Furthermore, twenty species are reported as new provincial records. A brief diagnosis accompanied by illustrations is provided for these new records in Iran.

Keywords: Opiinae, parasitoid, taxonomy, new records, distribution, Iran

Introduction

The Opiinae Blanchard, 1845 is one of the largest and most complex subfamilies among braconid wasps. To date, approximately 39 genera and 2,060 species have been described within this subfamily (Yu *et al.*, 2016; Wu *et al.*, 2018). In the early twentieth century, most research on Opiinae relied on Gahan's conservative classification (Gahan, 1915), where most species were grouped under the genus *Opius* Wesmael.

A series of studies initiated by Fischer in 1957 significantly expanded our understanding of Opiinae diversity. His work culminated in three comprehensive monographs (Fischer, 1972, 1977, 1987), which provided a detailed classification of the subfamily Opiinae. Fischer's generic divisions were critically

revised by Wharton and Marsh (1978), Wharton (1987), and Wharton and van Achterberg (2000), leading to several modifications. Notably, Wharton elevated *Psytalia* Walker and *Utetes* Förster to generic rank in his revisions (Wharton, 1987). The genus *Opius* Wesmael remains the largest within Opiinae, comprising over 1,282 described species (Fisher 1972, 1977, 1987; Wharton, 1987, 1988; van Achterberg, 1997; Yu *et al.*, 2016).

Species within the subfamily Opiinae are endoparasitoids of cyclorapha (Diptera) larvae and play a crucial role in controlling dipterous pests such as fruit-flies (Tephritidae) and leaf-mining flies (Agromyzidae) (Li *et al.*, 2012).

Intensive studies on various braconid subfamilies outlined in Iran around since 2008 (Rakhshani *et al.*, 2008, 2012; Ameri *et al.*, 2013, 2014, 2015; Farahani *et al.*, 2014, 2015,

Handling Editor: Ehsan Rakhshani

*Corresponding authors: talebia@modares.ac.ir

Received: 01 February 2025, Accepted: 09 April 2025

Published online: 09 April 2025

2016; Zargar *et al.*, 2014, 2015, 2019 a-e, 2020; Ranjbar *et al.*, 2016; Ghafouri *et al.*, 2018; Abdoli *et al.*, 2019 a, b, 2024; Pourhaji *et al.*, 2022). This research forms part of ongoing publications by authors on Iranian Opiinae alongside other researchers (Ghahari and van Achterberg, 2016; Gadallah *et al.*, 2016; Safahani *et al.*, 2018; Peris-Felipo *et al.*, 2018; Dolati *et al.*, 2018, 2019, 2021). This study provides detailed distributional data for 32 species and diagnostic characters for newly recorded species in Iran.

Materials and Methods

This study was conducted in the North-central area of Iran encompassing Alborz, Guilan, Mazandaran, Qazvin and Tehran provinces from March to November 2010 and 2011. Additional sampling was conducted in southwestern Iran's, Khuzestan Province between March 2016 and July 2017. Specimens were collected using Malaise traps across various ecosystems. The specimens were preserved in 75% ethanol, processed according to van Achterberg (2009), and then mounted and labeled. Images were captured with an Olympus SZX9 stereomicroscope equipped with a Canon (EOS 550D) digital camera. A series of four to five pictures stacked into a single focused image using Helicon Focus 7 software. Morphological terminology follows Karlsson and Ronquist (2012) and Wharton (2006). Identification of the specimens was made using the keys provided by Fischer (1972), Papp (1978), Tobias and Jakimavicius (1986) and van Achterberg (1997). All specimens are deposited in the Insect Collection of the Department of Entomology, Tarbiat Modares University, Tehran, Iran (TMUC).

Results

Order Hymenoptera Linnaeus, 1758

Family Braconidae Nees von Esenbeck, 1811

Subfamily Opiinae Blanchard, 1845

Genus *Apodesmia* Foerster, 1863

***Apodesmia irregularis* (Wesmael, 1835)**

Material examined: Iran, Khuzestan province: Behbahan, Dodangeh, (30°42'08.38" N, 50°10'41.81" E, 300 m a.s.l.), 5.V.2017, 1 ♀; Lali, Taraz, (32°20'49.70" N, 49°05'11.31" E, 390 m a.s.l.), 5.III.2017, 3 ♀, leg.: M. Zargar.

Distribution in Iran: Lorestan (Farahani *et al.*, 2016), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Europe, Nearctic (Yu *et al.*, 2016).

Apodesmia ocellata (Wesmael, 1835)

Figs. 1(A-I), 2

Material examined: Iran, Mazandaran province: Joorband (36°26'17.28" N, 52°07'16.62" E, 272 m a. s. l.), 26.IX.2011, 1 ♀; Mazandaran province: Tangehvaz (36°21'55.02" N, 52°06'10.74" E, 692 m a. s. l.), 11.X.2011, 2 ♀, leg.: M. Khayrandish.

Diagnosis (Female): Antennae somewhat longer than the body, with 33-34 antennomeres; oral cavity developed, mouthparts light colored (Fig. 1B); mesonotum with medio-posterior mesoscutal depression (Fig. 1D); notauli not reaching to medio-posterior mesoscutal depression (Fig. 1D); scutellum smooth (Fig. 1D); mesepimeral sulcus on posterior margin of mesopleuron sculptured (Fig. 1E); precoxal sulcus sculptured (Fig. 1E); pterostigma narrow, cuneate, r vein originating distinctly anterior to its middle, 3RSa vein longer than 2RS vein, m-cu vein interstitial (Fig. 1H); legs monochromatic, yellow or dark brownish (Fig. 1G); propodeum coarsely rugose (Fig. 1D); second metasomal tergite smooth (Fig. 1F); body length 3.0-3.2 mm (Fig. 1I).

Distribution in Iran: Mazandaran (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016), Iran (**new record**) (Fig. 2).

Apodesmia saeva (Haliday, 1837)

Material examined: Iran, Guilan province: Ziaz (36°52'34.44" N, 50°13'17.40" E, 537 m a. s. l.), 10.V.2010, 1 ♀; Mazandaran province: Joorband (36°26'15.54" N, 52°07'13.50" E, 275 m a. s. l.), 12.IV.2011, 1 ♀, leg.: M. Khayrandish.




Figure 1 *Apodesmia ocellata* (Wesmael, 1835): A. Head, dorsal view; B. Head, frontal view; C. Head, lateral view; D. Mesosoma, dorsal view; E. Mesosoma, lateral view; F. Metasoma, dorsal view. G. Hind leg; H. Wings; I. Habitus, lateral view.



Figure 2 Global distribution map of *Apodesmia ocellata* (Wesmael, 1835) (Hymenoptera: Braconidae, Opiinae).

Distribution in Iran: Mazandaran (Dolati *et al.*, 2019; current study), Guilan (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

Apodesmia striatula (Fischer, 1957)

Material examined: Iran, Mazandaran province: Gaznasara (36°16'58.08" N, 52°10'55.62" E, 2013 m a. s. l.), 11.V.2011, 1♀; Guilan province: Orkom (36°45'44.34" N, 50°18'11.88" E, 1201 m a. s. l.), 17.V.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Guilan (Dolati *et al.*, 2021; current study), Mazandaran (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

Genus *Biosteres* Foerster, 1863

Biosteres haemorrhoeus (Haliday, 1837)

Material examined: Iran, Alborz province: Sarziarat (35°55'10.38" N, 51°06'51.24" E, 1980 m a. s. l.), 08.VI.2011, 1♀, leg.: M. Khayrandish.

Distribution in Iran: West Azarbaijan, Guilan (Farahani *et al.*, 2016), East Azarbaijan (Ghahari and van Achterberg, 2016), Alborz (current study).

General distribution: Eastern and Western Palaearctic, Europe, Nearctic (Yu *et al.*, 2016).

Genus *Opiostomus* Fischer, 1972

Opiostomus griffithsi (Fischer, 1962)

Figs 3(A-I), 4

Material examined: Iran, Qazvin province: Loshan (36°40'09.12" N, 49°25'37.74" E, 291 m a. s. l.), 07.VI.2011, 1♀, leg.: M. Khayrandish.

Diagnosis (Female): Temple somewhat shorter than longitudinal diameter of eye (Fig. 3A); oral cavity not developed (Fig. 3B); mandibles abruptly widened basally (Fig. 3B); mesonotum with medio-posterior mesoscutal depression (Fig. 3D); precoxal sulcus sculptured (Fig. 3E); pterostigma cuneate, r vein originating from its basal third, 3RSa vein longer than 2RS vein, 3RSb vein 2.0 × as long as 3RSa vein, 3RSb vein not reaching wing apex (Fig. 3H); length of hind femur 5.0 × its maximum width in middle part (Fig. 3G); propodeum softly rugose (Fig. 3D); first metasomal tergite as long as its width at apex (Fig. 3F); ovipositor short (Fig. 3F); body length 2.1 mm (Fig. 3I).

Distribution in Iran: Qazvin (current study).

General distribution: Europe, Western Palaearctic (Yu *et al.*, 2016), Iran (**new record**) (Fig. 4).

Genus *Opius* Wesmael, 1835

Opius bouceki Fischer, 1958

Material examined: Iran, Khuzestan province: Baghmalek, Ghale-Tol, (31°37'49.70" N, 49°52'53.35" E, 880 m a.s.l.), 5.III.2017, 2 ♀♀, leg.: M. Zargar.




Figure 3 *Opiostomus griffithsi* (Fischer, 1962): A. Head, dorsal view; B. Head, frontal view; C. Head, lateral view; D. Mesosoma, dorsal view; E. Mesosoma, lateral view; F. Metasoma, dorsal view. G. Hind leg; H. Wings; I. Habitus, lateral view.



Figure 4 Global distribution map of *Opiostomus griffithsi* (Fischer, 1962) (Hymenoptera: Braconidae, Opiinae).

Distribution in Iran: Sistan & Baluchestan (Khajeh *et al.*, 2014), Kerman (Safahani *et al.*, 2018), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Opius caricivora* Fischer, 1964**

Material examined: Iran, Khuzestan province: Dezful, Ghale-Rob, (32°17'27.94" N, 48°25'46.98" E, 97 m a.s.l.), 5.VI.2017, 2 ♀♀, leg.: M. Zargar.

Distribution in Iran: Sistan & Baluchestan (Khajeh *et al.*, 2014), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Oriental (Yu *et al.*, 2016).

***Opius flavipes* Szepligeti, 1898**

Material examined: Iran, Guilan province: Orkom (36°45'44.34" N, 50°18'11.88" E, 1201 m a. s. l.), 18.X.2010, 1 ♀, leg.: M. Khayrandish.

Distribution in Iran: Hormozgan (Ameri *et al.*, 2014), Kerman (Safahani *et al.*, 2018), Guilan (Dolati *et al.*, 2021; current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Opius gracilis* Fischer, 1957**

Material examined: Iran, Alborz province: Sarzariat (35°55'10.38" N, 51°06'51.24" E, 1980 m a. s. l.), 07.IX.2010, 1 ♀, leg.: M. Khayrandish.

Distribution in Iran: Golestan (Gadallah *et al.*, 2018), Kerman (Safahani *et al.*, 2018), Khuzestan

(Gadallah *et al.*, 2022), Alborz, Guilan and Tehran (Dolati *et al.*, 2019; current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius instabilis* Wesmael, 1835**

Material examined: Iran, Khuzestan province: Abadan, Naghsh-e-Hayer, (30°12'08.15" N, 48°26'19.39" E, 1 m a.s.l.), 21.IV.2017, 1 ♀; Khoramshahr, Shaneh, (30°25'32.24" N, 48°11'20.83" E, 2 m a.s.l.), 22.V.2017, 1 ♀, leg.: M. Zargar.

Distribution in Iran: Lorestan, Chaharmahal and Bakhtiary (Gadallah *et al.*, 2022), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius larissa* Fischer, 1968**

Material examined: Iran, Guilan province: Ziaz (36°52'27.18" N, 50°13'24.78" E, 490 m a. s. l.), 08.XI.2010, 1 ♀; Alborz province: Sarzariat (35°55'10.38" N, 51°06'51.24" E, 1980 m a. s. l.), 15.VI.2010, 3 ♀♀, leg.: M. Khayrandish, Khuzestan province: Behbahan, Dodangeh, (30°42'08.38" N, 50°10'41.81" E, 300 m a.s.l.), 5.V.2017, 1 ♀, leg.: M. Zargar.

Distribution in Iran: Alborz, Guilan (Dolati *et al.*, 2019; current study), Khorasan Razavi (Gadallah *et al.*, 2022), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius levis* Wesmael, 1835**

Material examined: Iran, Guilan province: Orkom (36°45'44.34" N, 50°18'11.88" E, 1201 m a. s. l.), 14.VI.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Fars (Lashkari Bod *et al.*, 2010, 2011), Kermanshah (Gadallah *et al.*, 2016), East Azarbaijan (Gadallah *et al.*, 2022), Guilan (Farahani *et al.*, 2016; current study), Sistan & Baluchestan (Khajeh *et al.*, 2014), Isfahan (Gadallah *et al.*, 2016), Mazandaran (Dolati *et al.*, 2021).

General distribution: Afrotropical, Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius lugens* Haliday, 1837**

Material examined: Iran, Guilan province: Orkom (36°45'44.34" N, 50°18'11.88" E, 1201 m a. s. l.), 28.VI.2010, 1♀; Qazvin province: Zereskh Road (36°21'39.72" N, 50°03'55.56" E, 1541 m a. s. l.), 13.IV.2011, 1♀, leg.: M. Khayrandish, Khuzestan province: Baghmalek, Ghale-Tol, (31°37'49.70" N, 49°52'53.35" E, 880 m a.s.l.), 5.III.2017, 3 ♀♀, leg.: M. Zargar.

Distribution in Iran: Kermanshah (Gadallah *et al.*, 2016), Kerman (Safahani *et al.*, 2018), Guilan (Dolati *et al.*, 2021; current study), Khuzestan and Qazvin (current study).

General distribution: Afrotropical, Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius mischa* Fischer, 1968**

Material examined: Iran, Qazvin province: Zereskh Road (36°21'43.02" N, 50°03'53.22" E, 1553 m a. s. l.), 13.IV.2011, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Guilan (Dolati *et al.*, 2021), Qazvin (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius nigricoloratus* Fischer, 1958**

Material examined: Iran, Khuzestan province: Andika, Dorab, (32°12'23.00" N, 49°26'37.00" E, 760 m a.s.l.), 4.V.2016, 2 ♀♀, leg.: M. Zargar.

Distribution in Iran: Guilan (Gadallah *et al.*, 2016), Semnan (Gadallah *et al.*, 2022), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

***Opius pallipes* Wesmael, 1835**

Material examined: Iran, Khuzestan province: Behbahan, Dodangeh, (30°42'08.38" N, 50°10'41.81" E, 300 m a.s.l.), 5.VI.2017, 1 ♀, leg.: M. Zargar.

Distribution in Iran: Guilan (Gadallah *et al.*, 2016), Sistan & Baluchestan (Khajeh *et al.*, 2014), Kerman (Ranjbar *et al.*, 2016, Safahani *et al.*, 2018), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Nearctic, Oriental (Yu *et al.*, 2016).

Genus *Phaetrotoma* Foerster, 1863***Phaetrotoma cingulatus* (Wesmael, 1835)**

Material examined: Iran, Guilan province: Orkom (36°45'44.34" N, 50°18'11.88" E, 1201 m a. s. l.), 27.IX.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Kermanshah (Gadallah *et al.*, 2016), Khuzestan, Mazandaran (Gadallah *et al.*, 2022), Guilan (current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Phaetrotoma depeculator* Förster, 1863**

Material examined: Iran, Mazandaran province: Tangehvaz (36°21'55.68" N, 52°06'10.32" E, 702 m a. s. l.), 03.X.2011, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Kermanshah (Gadallah *et al.*, 2016), Ardabil (Ghahari and van Achterberg, 2016), Guilan and Mazandaran (Dolati *et al.*, 2018; current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Phaetrotoma exigua* (Wesmael, 1835)**

Material examined: Iran, Mazandaran province: Gaznasara (36°16'58.08" N, 52°10'55.62" E, 2013 m a. s. l.), 27.V.2011, 2♀♀, leg.: M. Khayrandish, Khuzestan province: Behbahan, Dodangeh, (30°42'08.38" N, 50°10'41.81" E, 300 m a.s.l.), 21.IV.2017, 1 ♀; Dezful, Ghale-Rob, (32°17'27.94" N, 48°25'46.98" E, 97 m a.s.l.), 4.V.2016, 3 ♀♀; Ramhormoz, Gharabad, (30°59'37.73" N,

49°46'50.63" E, 126 m a.s.l.), 5.V.2017, 2 ♀♀, leg.: M. Zargar.

Distribution in Iran: Tehran (Fischer, 1990), Sistan & Baluchestan (Khajeh *et al.*, 2014), Lorestan (Farahani *et al.*, 2016), Kerman (Ranjbar *et al.*, 2016, Safahani *et al.*, 2018), Alborz, Guilan, Mazandaran, Qazvin and Tehran (Dolati *et al.*, 2018), Khuzestan and Mazandaran (current study).

General distribution: Afrotropical, Eastern and Western Palaearctic, Europe, Oriental (Yu *et al.*, 2016).

***Phaedrotoma nitidulator* (Nees, 1834)**

Material examined: Iran, Khuzestan province: Ramhormoz, Gharabad, (30°59'37.73" N, 49°46'50.63" E, 126 m a.s.l.), 5.V.2017, 2 ♀♀, leg.: M. Zargar.

Distribution in Iran: Ardabil province (Gadallah *et al.*, 2016), Khuzestan province (current study).

General distribution: Eastern and Western Palaearctic Nearctic (Yu *et al.*, 2016).

***Phaedrotoma ochrogaster* (Wesmael, 1835)**

Material examined: Iran, Guilan province: Ghazichak (36°45'57.54" N, 50°19'35.22" E, 1803 m a. s. l.), 10.V.2010, 1♀; Alborz province: Arangeh (35°55'07.20" N, 51°05'09.24" E, 1891 m a. s. l.), 29.VI.2010, 3♀♀; Mazandaran province: Tangehvaz (36°21'55.02" N, 52°06'10.74" E, 692 m a. s. l.), 11.X.2011, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Alborz, Guilan and Mazandaran (Dolati *et al.*, 2019; current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Phaedrotoma pseudonitida* (Fahringer, 1943)**

Material examined: Iran, Tehran province: Shahriar (35°40'03.06" N, 50°56'52.14" E, 1168 m a. s. l.), 25.V.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Tehran (Dolati *et al.*, 2021; current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

***Phaedrotoma pulchriceps* (Szépligeti, 1898)**

Material examined: Iran, Alborz province: Karaj (35°46'08.88" N, 50°56'55.20" E, 1277 m a. s. l.), 18.V.2010, 1♀, Arangeh (35°55'07.20" N, 51°05'09.24" E, 1891 m a. s. l.), 13.VII.2010,

1♀; Tehran province: Shahriar (35°40'08.10" N, 50°56'56.64" E, 1168 m a. s. l.), 07.IX.2010, 4♀♀, leg.: M. Khayrandish.

Distribution in Iran: East Azarbaijan (Gadallah *et al.*, 2022), Guilan (Farahani *et al.*, 2016, reported as *Opius* (*Phaedrotoma*) *pulchriventris* (Fischer, 1958)), Kerman (Madjdzadeh *et al.*, 2021); Alborz and Tehran (Dolati *et al.*, 2018, reported as *Phaedrotoma pulchriventris* (Fischer, 1958); current study).

General distribution: Eastern and Western Palaearctic, Europe, Nearctic, Oriental (Yu *et al.*, 2016).

***Phaedrotoma variegata* (Szépligeti, 1896)**

Material examined: Iran, Khuzestan province: Ramhormoz, Gharabad, (30°59'37.73" N, 49°46'50.63" E, 126 m a.s.l.), 5.V.2017, 3♀♀; Khoramshahr, Shaneh, (30°25'32.24" N, 48°11'20.83" E, 2 m a.s.l.), 21.VI.2016, 2♀♀; Andika, Chezi, (32°08'02.78" N, 49°38'30.56" E, 650 m a.s.l.), 5.V.2017, 2♀♀; Lali, Taraz, (32°20'49.70" N, 49°05'11.31" E, 390 m a.s.l.), 4.IV.2017, 2♀♀, leg.: M. Zargar.

Distribution in Iran: Tehran (Dolati *et al.*, 2018), Khuzestan (current study).

General distribution: Eastern and Western Palaearctic, Europe (Yu *et al.*, 2016).

Genus *Psytalia* Walker, 1860

***Psytalia romani* (Fahringer, 1935)**

Figs. 5(A-I), 6

Material examined: Iran, Khuzestan province: Dezful, Shahid Mohammad-e-Montazeri, (32°17'27.94" N, 48°25'46.98" E, 97 m a.s.l.), 21.V.2016, 2♀♀, leg.: M. Zargar.

Diagnosis (Female): Antenna as long as body, with 34 antennomeres; width of head in dorsal view 2.1 × as wide as length (Fig. 5A); length of eye in dorsal view as long as temple (Fig. 5A); oral cavity developed (Fig. 5B); mesosoma 1.3 × its maximum height (Fig. 5E); notauli present in anterior half; medio-posterior mesoscutal depression not developed (Fig. 5D); side of mesothorax smooth (Fig. 5E); precoxal sulcus sculptured (Fig. 5E); pterostigma triangular; m-cu vein antefurcal; vein 1cu-a postfurcal; 3RSa vein 1.2 × longer than 2RS vein (Fig. 5H); length of hind femur 3.0 × its

maximum width in middle part (Fig. 5G); propodeum with two lateral carinae or longitudinal ridge (Fig. 5F); ovipositor exerted, $0.9 \times$ length of metasoma (Fig. 5F). Body length 3.5 mm (Fig. 5I).

Distribution in Iran: Khuzestan (current study).

General distribution: Eastern Palaearctic (Russia Far East, China, Korea), (Wu et al., 2016; Yu et al., 2016), Iran (**new record**) (Fig. 6).




Figure 5 *Psyttalia romani* (Fahringer, 1935): A. Head, dorsal view; B. Head, frontal view; C. Head, lateral view; D. Mesosoma, dorsal view; E. Mesosoma, lateral view; F. Metasoma, dorsal view; G. Hind leg; H. Wings; I. Habitus, lateral view.



Figure 6 Global distribution of *Psyttalia romani* (Fahringer, 1935) (Hymenoptera: Braconidae, Opiinae).

Genus *Psyttoma* van Achterberg & Li, 2012

***Psyttoma wachsmanni* (Szépligeti, 1898)**

Material examined: Iran, Tehran province: Iran National Botanical Garden (35°44'19.91" N, 51°10'52.49" E, 1265 m a. s. l.), 18.V.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Lorestan (Farahani *et al.*, 2016), Alborz, Guilan and Qazvin (Dolati *et al.*, 2019), Tehran (current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

Genus *Utetes* Foerster, 1863

***Utetes rotundiventris* (Thomson, 1895)**

Material examined: Iran, Alborz province: Sarziarat (35°55'10.38" N, 51°06'51.24" E, 1980 m a. s. l.), 15.VI.2010, 1♀; Mazandaran province: Joorband (36°26'17.28" N, 52°07'16.62" E, 272 m a. s. l.), 26.IX.2011, 2♀♀, leg.: M. Khayrandish.

Distribution in Iran: Markazi (Gadallah *et al.*, 2022), Guilan (Dolati *et al.*, 2018), Mazandaran (Dolati *et al.*, 2018; current study), Alborz (current study).

General distribution: Eastern and Western Palaearctic (Yu *et al.*, 2016).

Genus *Xynobius* Foerster, 1863

***Xynobius caelatus* (Haliday, 1837)**

Material examined: Iran, Mazandaran province: Gaznasara (36°16'56.82" N, 52°10'58.50" E, 2032 m a. s. l.), 11.V.2011, 2♀♀; Mazandaran province: Tangehvaz

(36°21'55.68" N, 52°06'10.32" E, 702 m a. s. l.), 12.IV.2011, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Mazandaran and Qazvin (Dolati *et al.*, 2018; current study).

General distribution: Europe and Western Palaearctic (Yu *et al.*, 2016).

***Xynobius comatus* (Wesmael, 1835)**

Figs 7(A-I), 8

Material examined: Iran, Mazandaran province: Tangehvaz (36°21'55.02" N, 52°06'10.74" E, 692 m a. s. l.), 11.X.2011, 2♀♀, leg.: M. Khayrandish.

Diagnosis (Female): Antenna 1.4 × as long as body, with 23-24 antennomeres (Fig. 7I); width of head about 2.0 × its maximum height (Fig. 7A); oral cavity developed (Fig. 7B); mesosoma about 1.6 × its maximum height (Fig. 7E); mesonotum with medio-posterior mesoscutal depression (Fig. 7D); notauli reaching to medio-posterior mesoscutal depression (Fig. 7D); scutellum smooth (Fig. 7D); precoxal sulcus sculptured (Fig. 7E); pterostigma almost semi-elliptical, 3RSa vein 1.5 × as long as 2RS vein, 3RSb vein straight, about 2.0 × as long as 3RSa vein, m-cu vein interstitial to slightly antefurcal, wings hyaline transparent (Fig. 7H); legs long, thin and yellow, length of hind femur 5.0 × its maximum width in middle part (Fig. 7G); propodeum rugose, matte, lacking distinct longitudinal ridge (Fig. 7D); first metasomal tergite 1.3 × as long as its width at apex (Fig. 7F); ovipositor barely exerted (Fig. 7F); body dark colored, face and 2nd metasomal tergite

with somewhat developed red or yellowish pattern (Fig. 7I); body length 2.5 mm (Fig. 7I).

Distribution in Iran: Mazandaran (current study).

General distribution: Eastern and Western Palaearctic, Nearctic, Oriental (Yu *et al.*, 2016), Iran (**new record**) (Fig. 8).




Figure 7 *Xynobius comatus* (Wesmael, 1835): A. Head, dorsal view; B. Head, frontal view; C. Head, lateral view; D. Mesosoma, dorsal view; E. Mesosoma, lateral view; F. Metasoma, dorsal view. G. Hind leg; H. Wings; I. Habitus, dorsal view.



Figure 8 Global distribution of *Xynobius comatus* (Wesmael, 1835) (Hymenoptera: Braconidae, Opiinae).

***Xynobius curtifemur* (Fischer, 1961)**

Material examined: Iran, Guilan province: Ghazichak (36°45'52.62" N, 50°20'01.08" E, 1787 m a. s. l.), 03.V.2010, 3 ♀♀, leg.: M. Khayrandish.

Distribution in Iran: Guilan (Dolati *et al.*, 2018; current study).

General distribution: Europe and Western Palearctic (Yu *et al.*, 2016).

***Xynobius maculipes* (Wesmael, 1835)**

Material examined: Iran, Khuzestan province: Khoramshahr, Shaneh, (30°25'32.24" N, 48°11'20.83" E, 2 m a.s.l.), 4.IV.2017, 2 ♀♀; Baghmalek, Ghale-Tol, (31°37'49.70" N, 49°52'53.35" E, 880 m a.s.l.), 22.VI.2017, 1 ♀; Dezful, Ghale-Rob, (32°17'27.94" N, 48°25'46.98" E, 97 m a.s.l.), 5.VI.2017, 2 ♀♀, leg.: M. Zargar.

Distribution in Iran: Tehran (Fischer 1990), Guilan (Gadallah *et al.*, 2016), East Azerbaijan (Gadallah *et al.*, 2022), Mazandaran (Dolati *et al.*, 2019), Khuzestan (current study).

General distribution: Eastern and Western Palearctic, Oriental (Yu *et al.*, 2016).

Discussion

The results revealed the presence of 32 species from the subfamily Opiinae, with four species recorded for the first time in Iran. Our findings have increased the total number of Iranian Opiinae to 137 (Safahani *et al.*, 2018; Peris-Felipo *et al.*, 2018; Dolati *et al.*, 2018, 2019,

2021; Ameri *et al.*, 2020; current study) and the total number of Braconidae to 926 (Rakhshani *et al.*, 2008, 2012; Ameri *et al.*, 2013, 2014, 2015, 2020; Farahani *et al.*, 2014, 2015, 2016; Zargar *et al.*, 2014, 2015, 2019 a-e, 2020; Ranjbar *et al.*, 2016; Abdoli *et al.*, 2019 a, b).

This study provides the first report of four species from Iran at the border between Eastern and Western Palearctic regions. Three species (*Apodesmia ocellata*, *Opiostomus* (*Opiostomus*) *griffithsi*, *Psytalia* (*Psytalia*) *romani*) exhibit relatively uniform distribution across the Palearctic region. However, *Xynobius* (*Xynobius*) *comatus* is distributed across Palearctic, Nearctic and Oriental regions (Yu *et al.*, 2016). Most of the recorded species were found in Eastern and Western Palearctic regions as well as Europe (29 Eastern Palearctic species and both Western Palearctic & Europe having each about 31 species), while fewer are reported from other regions like Afrotropical (three species), Nearctic (six species) and Oriental (six species). In neighboring countries, the number of recorded species varies significantly. For example, Azerbaijan has three recorded species, Turkmenistan has ten, and Armenia has 23 (Yu *et al.*, 2016). These numbers stand in stark contrast to those of neighboring countries such as Russia, with 319 (Tobias 1986, 1988; Yu *et al.*, 2016), and Turkey, with 174 species (Beyarslan and Fischer, 2011, 2013; Yu *et al.*, 2016; Beyarslan 2020). Given these disparities, it is evident that further research on this

subfamily in Iran is necessary. Considering the high diversity of Opiinae and their crucial role in the biological control of phytophagous Diptera (Fischer and Koponen, 1999), further research focusing on taxonomy, faunistics, and host associations is essential.

Acknowledgments

We thank the Department of Entomology of Tarbiat Modares University for supporting this study financially. The authors wish to express their highest gratitude to Dr. Cornelis van Achterberg (Naturalis Biodiversity Center, P.O. 9517, 2300 RA Leiden, The Netherlands) and Dr. Peris-Felipo (Department of Agricultural Chemistry and Food Science, Autonomous University of Madrid, Spain) for their valuable guidance and supports. We cordially thank the Subject Editor and two anonymous reviewers for their valuable comments and recommendations on the earlier version of this paper.

References

- Abdoli, P., Talebi, A. A. and Farahani, S. 2019a. *Dolichogenidea fernandeztrianai* sp. nov. (Hymenoptera: Braconidae, Microgastrinae) from Iran. *Journal of Agricultural Science and Technology*, 21(3): 647-658.
- Abdoli, P., Talebi, A. A., Farahani, S. and Fernandez-Triana, J. 2019b. Three new species of the genus *Choeras* Mason, 1981 (Hymenoptera: Braconidae, Microgastrinae) from Iran. *Zootaxa*, 4545: 77-92. <https://doi.org/10.11646/zootaxa.4545.1.4>
- Abdoli, P., Talebi, A. A., Kavallieratos, N. G., Khosravi, R., and Bidari, F. 2024. Contribution to the phylogeny of Microgastrinae (Hymenoptera: Braconidae) based on mitochondrial COI and nuclear 28S rDNA genes, with comments on the identity of *Pholetesor circumscriptus* (Nees, 1834). *Journal of Insect Biodiversity and Systematics*, 10 (4): 965-981. <https://doi.org/10.61186/jibs.10.4.965>.
- Ameri, A., Talebi, A. A., Beyarslan, A., Kamali, K. and Rakhshani, E. 2013. Study of the genus *Bracon* Fabricius, 1804 (Hymenoptera: Braconidae) of southern Iran with description of a new species. *Zootaxa*, 3754: 353-380. <https://doi.org/10.11646/zootaxa.3754.4.2>.
- Ameri, A., Talebi, A. A., Rakhshani, E., Beyarslan, A. and Kamali, K. 2014. Study of the genus *Opius* Wesmael (Hymenoptera: Braconidae: Opiinae) in Southern Iran, with eleven new records. *Zootaxa*, 3884: 1-26. <https://doi.org/10.11646/zootaxa.3884.1.1>.
- Ameri, A., Talebi, A. A., Rakhshani, E. and Beyarslan, A. 2015. A Review of the Subfamily Hormiinae (Hymenoptera: Braconidae) from Iran. *Journal of Insect Biodiversity and Systematics*, 1(2): 111-123.
- Ameri, A., Ebrahimi, E. and Talebi, A. A. 2020. Additions to the fauna of Braconidae (Hym.: Ichneumonoidea) of Iran based on the specimens housed in Hayk Mirzayans Insect Museum with six new records for Iran. *Journal of Insect Biodiversity and Systematics*, 6(4): 353-364. <http://dx.doi.org/10.52547/jibs.6.4.353>.
- Beyarslan, A. and Fischer, M. 2011. Contributions to the Opiinae fauna of Turkey (Hym.: Braconidae). *Turkish Journal of Zoology*, 35(3): 293-305. <https://doi.org/10.3906/zoo-0903-20>.
- Beyarslan, A. and Fischer, M. 2013. Checklist of Turkish Opiinae (Hymenoptera: Braconidae). *Zootaxa*, 3721(5): 401-454. <http://dx.doi.org/10.11646/zootaxa.3721.5.1>.
- Beyarslan, A. 2020. Taxonomic studies on the Opiinae (Hymenoptera, Braconidae) fauna of the Turkish central part of Eastern Anatolia Region (Bingöl, Bitlis, Muş and Van). *Acta Biologica Turcica*, 33(1): 1-7.
- Dolati, S., Talebi, A. A., Farahani, S. and Khayrandish, M. 2018. New finding in Opiinae (Hymenoptera: Braconidae) from north of Iran. *Journal of Insect Biodiversity and Systematics*, 4(3): 163-182. <http://dx.doi.org/10.52547/jibs.4.3.163>.
- Dolati, S., Talebi, A. A., Farahani, S. and Khayrandish, M. 2019. New Data of the Genus *Opius* Wesmael (Hymenoptera: Braconidae, Opiinae) from Northern Iran.

- Journal of Agricultural Science and Technology, 21: 1871-1887.
- Dolati, S., Talebi A. A., Peris-Felipo, F. J., Farahani, S. and Khayrandish, M. 2021. New finding in Opiinae (Hymenoptera: Braconidae) from north of Iran. Zootaxa, 4903(3): 331-352. <https://doi.org/10.11646/zootaxa.4903.3.2>.
- Farahani, S., Talebi, A. A., Rakhshani, E., van Achterberg, C. and Sharkey, M. 2014. A contribution to the knowledge of Agathidinae (Hymenoptera: Braconidae) from Iran with description of a new species. Biologia, 69: 228-235. <https://doi.org/10.2478/s11756-013-0295-y>
- Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2015. A review of the subfamily Rogadinae (Hymenoptera: Braconidae) from Iran. Zootaxa, 3973: 227-250. <https://doi.org/10.11646/zootaxa.3973.2.2>.
- Farahani, S., Talebi, A. A. and Rakhshani, E. 2016. Iranian Braconidae (Insecta: Hymenoptera: Ichneumonoidea): diversity, distribution and host association. Journal of Insect Biodiversity and Systematics, 2: 1-92.
- Fischer, M. 1957. Die europäischen Arten der Gattung *Opius* Wesm. Teil IIa (Hymenoptera, Braconidae). Deutsche Entomologische Zeitschrift, 4(5): 332-358. <https://doi.org/10.1002/mmnd.4800040504>
- Fischer, M. 1972. Hymenoptera, Braconidae (Opiinae I). Das Tierreich, 91: 1-620.
- Fischer, M. 1977. Hymenoptera Braconidae (Opiinae II). (Amerika). Das Tierreich, 96: 1-1001.
- Fischer, M. 1987. Hymenoptera Braconidae (Opiinae III)-äthiopische, orientalische, australische und ozeanische Region. Das Tierreich, 104: 1-734.
- Fischer, M. 1990. Paläarktische Opiinae (Hymenoptera, Braconidae): Neue Arten und neue Funde aus dem Ungarischen Naturwissenschaftlichen Museum in Budapest. Annales Historico-Naturales Musei Nationalis Hungarici, 81: 205-238.
- Fischer, M. and Koponen, M. 1999. A survey of Opiinae (Hymenoptera: Braconidae) of Finland, part 2. Entomologica Fennica, 10(3): 129-160. <https://doi.org/10.33338/ef.84013>.
- Gahan, A. B. 1915. A revision of North American ichneumon-flies of the subfamily Opiinae. Proceedings of United States National Museum, 49: 63-95. <https://doi.org/10.5479/si.00963801.49-2095.63>.
- Gadallah, N. S., Ghahari, H., Peris-Filipo, F. J. and Fischer, M. 2016. Updated checklist of Iranian Opiinae (Hymenoptera: Braconidae). Zootaxa, 4066: 1-40. <https://doi.org/10.11646/zootaxa.4066.1.1>.
- Gadallah, N. S., Ghahari, H., Papp, J. and Beyarslan, A. 2018. New records of Braconidae (Hymenoptera) from Iran. Wuyi Science Journal, 34: 43-48.
- Gadallah, N. S., Ghahari, H. and Quicke, D. L. J. 2021. Further addition to the braconid fauna of Iran (Hymenoptera: Braconidae). Egyptian Journal of Biological Pest Control, 31: 1-9. <https://doi.org/10.1186/s41938-021-00376-8>.
- Gadallah, N. S., Peris Felipo, F. J., Ghahari, H., Quicke, D. J., Shaw, S. R., Fischer, M. 2022. Subfamily Opiinae Blanchard, 1845. In: Gadallah N. S., Shaw, S. R. and Ghahari, H. (Eds.), Braconidae of the Middle East (Hymenoptera): Taxonomy, Distribution, Biology, and Biocontrol Benefits of Parasitoid Wasps. United Kingdom: Elsevier Science, PP. 454-489. <https://doi.org/10.1016/B978-0-323-96099-1.00014-5>.
- Ghahari, H. and van Achterberg, C. 2016. A contribution to the study of subfamilies Microgastrinae and Opiinae (Hymenoptera: Braconidae) from the Arasbaran Biosphere Reserve and vicinity, Northwestern Iran. Natura Somogyiensis, 28: 23-32. <https://doi.org/10.24394/NatSom.2016.28.23>
- Ghafouri Moghaddam M., Rakhshani, E., van Achterberg, C. and Mokhtari, A. 2018. A study of the Iranian species of *Choeras* Mason (Hymenoptera: Braconidae: Microgastrinae), with the description of a new species. Zootaxa 4446(4): 455-476. <https://doi.org/10.11646/zootaxa.4446.4.3>.
- Karlsson, D. and Ronquist, F. 2012. Skeletal Morphology of *Opius dissitus* and *Biosteres carbonarius* (Hymenoptera: Braconidae), with

- a Discussion of Terminology. PLoS ONE, 7(4): e32573. <https://doi.org/10.1371/journal.pone.0032573>.
- Khajeh, N., Rakhshani, E., Peris-Felipo, F.J. and Zikic, V. 2014. Contributions to the Opiinae (Hymenoptera: Braconidae) of Eastern Iran with updated checklist of Iranian species. *Zootaxa*, 3784(2): 131-147. <https://doi.org/10.11646/zootaxa.3784.2.3>.
- Lashkari Bod, A., Rakhshani, E., Talebi, A. A. and Fischer, M. 2010. Identification and introduction of ten new records of Alysiinae (Hym.: Braconidae) from Iran. *Proceedings of the 19th Plant Protection Congress, Tehran, Iran*, p. 160.
- Lashkari-Bod, A., Rakhshani, E., Talebi, A. A., Lozan, A. and Žikic, V. 2011. A contribution to the knowledge of Braconidae (Hym., Ichneumonoidea) of Iran. *Biharean Biologist*, 5: 147-150.
- Li, X. Y., van Achterberg, C. and Tan, J. C. 2013. Revision of the subfamily Opiinae (Hymenoptera, Braconidae) from Hunan (China), including thirty-six new species and two new genera. *ZooKeys.*, 268: 1-168. <https://doi.org/10.3897/zookeys.268.4071>.
- Madjdzadeh, S. M., Purrezaali, M., Dolati, S. and Ghassemi-Khademi, T. 2021. New data on braconid wasps (Hymenoptera: Braconidae: Cheloninae, Opiinae, Rogadinae) of South-Eastern Iran. *Entomologie faunistique (Faunistic Entomology)*, 74: 1-7.
- Papp, J. 1978. Braconidae (Hymenoptera) from Korea, III. *Acta Zoologica Academiae Scientiarum Hungaricae*, 24: 133-148.
- Peris-Felipo, F. J., Belokobylskij, S. A., Derafshan, H. A. and Rakhshani, E. 2018. Revision of the genus *Hoplocrotaphus* Telenga, 1950 (Hymenoptera, Braconidae, Opiinae). *Journal of Hymenoptera Research*, 62: 55-72. <https://doi.org/10.3897/jhr.62.21858>
- Pourhaji, A., Lotfalizadeh, H., Kasebi, N., Gharali, B., Ameri, A. 2022. Associated *Bracon* Fabricius (Hymenoptera: Braconidae) with flower-heads of Asteraceae in Iran. *Journal of Insect Biodiversity and Systematics*, 8 (1): 35-47. <http://dx.doi.org/10.52547/jibs.8.1.35>.
- Rakhshani, E., Talebi, A. A., Stary, P., Tomanović, Ž., Manzari, S. and Kavallieratos, N. G. 2008. A review of *Aphidius* Nees (Hymenoptera, Braconidae, Aphidiinae) in Iran: host associations, distribution and taxonomic notes. *Zootaxa*, 1767: 37-54. <https://doi.org/10.5281/zenodo.182039>.
- Rakhshani, E. 2012. Aphid parasitoids (Hym., Braconidae, Aphidiinae) associated with pome and stone fruit trees in Iran. *Journal of Crop Protection*, 1: 81-95.
- Ranjbar, M., Majdzadeh, S. M., Peris-Felipo, F. J., Askari, M., Rakhshani, E. 2016. A Contribution to the Fauna of Opiinae (Hym.: Braconidae) of Kerman Province South-Eastern Iran. *Journal of Entomological Research Society*, 18: 19-26.
- Safahani, S., Madjdzadeh, S. M. and Peris-Felipo, F. J. 2018. Contribution to the fauna and phenological knowledge of high mountains Opiinae (Hymenoptera, Braconidae) in Kerman province (Iran). *Journal of Insect Biodiversity and Systematics*, 4: 73-83. <http://dx.doi.org/10.52547/jibs.4.2.73>.
- Tobias. V. I. and Jakimavicius, A. B. 1986. Otryad Hymenoptera - Pereponchatokrylye, Semestvo Braconidae, Podsemestvo Opiinae. In: Medvedev GS (Ed.) *Opredelitel' nasekomykh Evropekskosti SSSR*. Vol. 3. Hymenoptera. Part 5. Nauka, Leningrad, pp 7-100.
- van Achterberg, C. 1997. Revision of the Haliday collection of Braconidae (Hymenoptera). *Zoologische Verhandelingen Leiden*, 314: 1-115.
- van Achterberg, C. 2009. Can Townes type Malaise traps be improved? Some recent developments. *Entomologische Berichten*, 69 (4): 129-135.
- Wharton, R. A. 1987. Changes in nomenclature and classification of some opine Braconidae (Hymenoptera). *Proceedings of the Entomological Society of Washington*, 89: 61-73.
- Wharton, R. A. 1988. Classification of the braconid subfamily Opiinae (Hymenoptera).

- Canadian Entomologist, 120: 333-360. <https://doi.org/10.4039/Ent120333-4>.
- Wharton, R. A. 2006. The species of *Sternaulopius* Fischer (Hymenoptera: Braconidae, Opiinae) and the braconid *sternaulus*. Journal of Hymenoptera Research, 15(2): 317-347.
- Wharton, R. A. and Marsh, P. M. 1978. New World Opiinae (Hymenoptera: Braconidae) parasitic on Tephritidae (Diptera). Journal of the Washington Academy of Sciences, 68:147-167.
- Wharton, R. A. and van Achterberg, C. 2000. Family group names in Braconidae (Hymenoptera: Ichneumonoidea). Journal of Hymenoptera Research, 9: 254-270.
- Wu, Q., van Achterberg, C., Sheng, Y-Y. and Chen, X-X. 2018. A new genus from Vietnam (Hymenoptera, Braconidae, Opiinae), and the description of two new species. Journal of Hymenoptera Research, 15(2): 66: 1-13. <https://doi.org/10.3897/jhr.66.28881>.
- Yu, D. S., van Achterberg, C. and Horstmann, K. 2016. Taxapad 2016, Ichneumonoidea 2015. Database on flash-drive. <http://www.taxapad.com>, Nepean, Ontario, Canada.
- Zargar, M., Talebi, A. A., Hajiqaanbar, H., Farahani, S. and Ameri, A. 2014. A contribution to the knowledge of Braconinae (Hymenoptera: Braconidae) in some parts of northern and southern Iran. Journal of Crop Protection, 3(2): 233-243.
- Zargar, M., Talebi, A. A., Hajiqaanbar, H. and Papp, J. 2015. A study on the genus *Bracon* Fabricius (Hymenoptera: Braconidae) in north central Iran with four new records for Iranian fauna. Entomofauna, 36: 425-440.
- Zargar, M., Gupta, A., Talebi, A. A. and Farahani, S. 2019a. A review of the Iranian species of genus *Iconella* Mason (Hymenoptera: Braconidae: Microgastrinae) with description of two new species. Zootaxa, 4586 (3): 491-504. <https://doi.org/10.11646/zootaxa.4586.3.6>.
- Zargar, M., Gupta, A., Talebi, A. A. and Farahani, S. 2019b. Description of a new species of the genus *Deuterixys* (Hymenoptera: Braconidae: Microgastrinae) from Iran. Biologia, 75: 267-272. <https://doi.org/10.2478/s11756-019-00298-9>.
- Zargar, M., Gupta, A., Talebi, A. A. and Farahani, S. 2019c. Three new species and two new records of the genus *Cotesia* Cameron (Hymenoptera: Braconidae) from Iran. European Journal of Taxonomy, 571: 1-25. <https://doi.org/10.5852/ejt.2019.571>.
- Zargar, M., Talebi, A. A. and Farahani, S. 2019d. Faunistic study of the genus *Habrobracon* Ashmead (Hymenoptera: Braconidae) from Iran. Journal of Insect Biodiversity and Systematics, 05(3): 159-169. <https://doi.org/10.52547/jibs.5.3.159>.
- Zargar, M., Talebi, A. A. and Farahani, S. 2019e. Occurrence of *Ceratobracon stschegolevi* (Telenga, 1936) (Hymenoptera: Braconidae: Braconinae), a rare genus and species new for Iran. Biharean Biologist, 14 (1): 1-4.
- Zargar, M., Konstantin, S., Talebi, A. A. and Farahani, S. 2020. Study of the genus *Bracon* Fabricius 1804 (Hymenoptera: Braconidae) from Iran with an updated checklist. Zootaxa, 4758 (2): 201-230. <https://doi.org/10.11646/zootaxa.4758.2.1>.

اطلاعات تکمیلی در مورد زنبورهای پارازیتوئید زیرخانواده Opiinae (Hymenoptera: Braconidae) همراه با گزارش جدید چهار گونه در ایران

ثنا دولتی^۱، علی اصغر طالبی^{۱*} و سمیرا فراهانی^۲

- ۱ - گروه حشره شناسی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران.
 - ۲ - مؤسسه تحقیقات جنگل ها و مراتع، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران.
- پست الکترونیکی نویسنده مسئول مکاتبه: talebia@modares.ac.ir
دریافت: ۱۲ بهمن ۱۴۰۳؛ پذیرش: ۲۰ فروردین ۱۴۰۴

چکیده: در این مطالعه، فون زنبورهای زیرخانواده Opiinae در مناطق شمال مرکزی (۱۳۸۹-۱۳۹۱) و قسمت جنوب غربی ایران، استان خوزستان (۱۳۹۴-۱۳۹۵) مورد بررسی قرار گرفت. نمونه ها با استفاده از تله مالیز جمع آوری شدند. در نتیجه ۳۲ گونه شناسایی شدند که گونه های *Opiostomus (Opiostomus) griffithsi* (Fischer, ۱۸۳۵)، *Apodesmia ocellata* (Wesmael, ۱۸۳۵)، *Psytalia (Psytalia) romani* و *Xynobius (Xynobius) comatus* (Wesmael, ۱۸۳۵)، ۱۹۶۲ (Fahringer, ۱۹۳۵) برای اولین بار از ایران گزارش می شوند. علاوه بر این، ۲۰ گونه رکوردهای جدید استانی هستند. توصیف مختصر و عکس های مربوط به گونه هایی که برای اولین بار از ایران گزارش شده اند، تهیه شده است.

واژگان کلیدی: Opiinae، خوزستان، پارازیتوئید، گزارش جدید، پراکنش، ایران