

Research Article

Control of *Tetranychus urticae* by three predatory mites (Acari: Phytoseiidae) in a commercial greenhouse rose

Kamaloddin Ahmadi, Yaghoub Fathipour^{*} and Mohammad Bashiri

Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

Abstract: The two-spotted spider mite (TSSM), Tetranychus urticae Koch (Acari: Tetranychidae) is a serious pest of many greenhouse crops such as bean, cucumber, rose and other products. The control of this important pest has been mainly based on the use of acaricides. Phytoseiid mites are used mostly for biological control of pest mites. In this study, control of TSSM by phytoseiid predatory mites, Phytoseiulus persimilis Athias-Henriot, Amblyseius swirskii Athias-Henriot and Neoseiulus californicus McGregor was evaluated on four rose varieties including Avalanche, Dolcevita, Samurai and Sorbet in a commercial rose greenhouse. The total number of motile stages and eggs of each studied species on a leaf were counted weekly, through a zigzag sampling pattern and using a hand lens. The sampling was carried out randomly and the leaves of a rose plant were taken from the canopy base (shoots bent over beds), the middle (area between base and top) and the top of the canopy (flowering shoots), which added up to a total of 30 leaves. The highest population density of TSSM per leaf was recorded on Samurai (17.96 \pm 0.85) and the lowest population density was observed on Dolcevita (5.32 \pm 0.39). Based on population fluctuation data of TSSM and its predators on four rose varieties, it was found that the predatory mite P. persimilis had the ability to reduce the high TSSM density and N. californicus also continued to operate in low TSSM density, but A. swirskii did not have a clear impact on TSSM density reduction.

Keywords: biological control, integrated mite management, population fluctuation, *Tetranychus urticae*, phytoseiid mite

Introduction

Cut roses are one of the most popular ornamental plants in the world (Alipour *et al.*, 2016). The major cut roses produced commercially in Iran are greenhouse roses. The two-spotted spider mite (TSSM), *Tetranychus urticae* Koch (Acari: Tetranychidae) is a serious pest of many greenhouse crops such as bean, cucumber, rose and many others (Maleknia *et* *al.*, 2016). Damage of this pest on ornamental plants reduces the flower's beauty, so that damage symptoms on leaves and petals can reduce the price of the product.

The control of spider mites has been mainly based on the use of acaricides (Khajehali *et al.*, 2011). The wide use of acaricides has developed acaricide resistance in mites and is one of the reasons for environmental concern (Escudero and Ferragut, 2005; Attia *et al.*, 2013). Also the increase of consumer demand for healthy products has influenced the use of pesticides in crop production and many growers have tended to deployment of the integrated pest management (IPM) (De Vis and Barrera, 1999;

Handling Editor: Ali Asghar Talebi

^{*} **Corresponding authors**, e-mail: fathi@modares.ac.ir Received: 4 May 2019, Accepted: 10 November 2019 Published online: 4 December 2019

Nicetic et al., 2001; Opit et al., 2004; Casey et al., 2007; Holt et al., 2007). Biological control is an important part of IPM programs. For pest mites, the predators have been used more than other biological agents (Fathipour and Maleknia, 2016). It seems that the mite predators are impartible part of biological control programs in integrated mite management. Among these predators, phytoseiid mites are most used in the control of TSSM (Fathipour and Maleknia, 2016). The phytoseiid predator, Phytoseiulus persimilis Athias-Henriot is a specialist predator of TSSM which is often used as a periodic diffusion in biological control programs in greenhouse cut roses (Casey et al., 2007; McMurtry et al., 2013). Amblyseius swirskii Athias-Henriot is a generalist predator mainly used for augmentative biological control of mites, thrips and whiteflies for a range of greenhouse crop (Ragusa and Swirski, 1975; Buitenhuis et al., 2015). Neoseiulus californicus McGregor is one of the most important phytoseiid predators with a feeding preference for TSSM. These predators are used against TSSM in a range of greenhouse crops of temperate and subtropical regions around the world (Fathipour and Maleknia, 2016).

Although biological control is a safe and potentially effective method, it is usually not enough to protect the crop against pests alone (Gerson and Weintraub, 2007). One of the key elements for building a sustainable IPM program is the use of host plant resistance to pests (Khanamani et al., 2013), which can be used as a suitable replacement to broadspectrum insecticides (Fathipour and Mirhosseini, 2017). One of the efficient strategies to manage TSSM is combination of biological control agents and resistant plants (Khanamani et al., 2014).

In the present study, we aimed to determine population fluctuation of TSSM and its predators *P. persimilis*, *A. swirskii* and *N. californicus* on four rose varieties (Avalanche, Dolcevita, Samurai and Sorbet) under greenhouse conditions. Specifically, our findings could be used for the implementation of IPM programs against TSSM in the rose greenhouses.

Materials and Methods

This study was conducted in a commercial rose greenhouse with an area of ten thousand square meters at Karaj, which was located at altitude of 1300-1500 m above sea level. This greenhouse had been covered by glass, and all bush roses were cultivated on raised beds in a hydroponic cultivation system. Four rose varieties including Avalanche, Dolcevita, Samurai and Sorbet were cultivated in this greenhouse. The biological control program has been carried out based on the experimental plan suggested by the Koppert Biological Systems and we only recorded the results of the mentioned control program.

Treatments against TSSM included the use of *P. persimilis*, *A. swirskii* and *N. californicus* obtained from Koppert Biological Systems. The time and rate of releasing the phytoseiid predators during the project (for 31 weeks) is presented in Table 1. The release of the predators on each release date was applied equally regardless of plant variety in the greenhouse, except for the release of *P. persimilis* in the seventeenth week, which was applied as a hot-spot treatment.

Table 1Phytoseiidmitepredatorsusedingreenhouses rose.

Predatory	Application rate	Number	Release date
mite		per unit	(Week)
P. persimilis	40 bottles	2000	3/8/2016(1)
A. swirskii	2000 plus sachets	250	10/8/2016 (2)
P. persimilis	150 bottles	2000	23/8/2016 (3)
P. persimilis	70 bottles	2000	30/8/2016 (4)
P. persimilis	20 bottles	2000	6/9/2016 (5)
A. swirskii	5 bottles	5000	4/10/2016 (9)
N. californicus	4 bottles	25000	25/10/2016 (12)
P. persimilis	5 bottles	2000	30/11/2016 (17)
P. persimilis	23 bottles	2000	17/2/2017 (24)
P. persimilis	60 bottles	2000	2/3/2017 (26)
N. californicus	8 bottles	25000	2/3/2017 (26)

Abbreviations: P: Phytoseiulus; A: Amblyseius; N: Neoseiulus.

Downloaded from jcp.modares.ac.ir on 2025-05-21]

The release of the predators was done on the bent stems by sprinkling the contents of a bottle on the leaves while the bottle was rotated regularly for the uniform distribution of the predatory mites. This work was done from the end of each row to the beginning of the rows. Also slow-release sachet of *A. swirskii* was used by hanging on the bent stems of roses.

To provide sufficient moisture and for better development of the predatory mites in greenhouse, for ten days after each release period, water was sprayed on the bushes daily. During the study period, the pruned leaves were kept for one day on planting bed near the bushes, so that the predatory mites on them were returned to bushes, so we tried to prevent predators from leaving the greenhouse.

Leaves were considered as the sampling units. For collecting leaf samples, a zigzag pattern was used. In this way, the samplings of the mites were done in the aisles between the rows of plants, from both sides of the aisle. In each sampling, 30 rose leaves were sampled from 30 bushes. The sampling was carried out randomly and the leaves of a rose plant were taken from the canopy base (shoots bent over beds), the middle of the canopy (area between base and top) and the top (flowering shoots). Total motile stages (combined adults and juveniles) and eggs of TSSM and its predators on a leaf were counted weekly using a hand lens. All counts were performed in the morning. Density of these mites was determined from 6th September 2016 to 4th April 2017.

One-way ANOVA was used to compare the mean density of total life stages (egg, nymphs and adults) of TSSM and its predators among rose varieties within each sampling date. Differences among the means were determined at a significance level of 5% using Tukey's test. All data were tested for normality before analysis and data were analyzed using Minitab software.

Results

Population fluctuation of TSSM

The mean population density of TSSM per leaf of different rose varieties in each sampling date is shown in Table 2. The TSSM population density results showed a significant difference among different rose varieties at different dates. TSSM population density for the entire sampling period on each leaf for Avalanche, Dolcevita, Samurai and Sorbet varieties was record 14 ± 0.73 , 5.32 ± 0.39 , 17.96 ± 0.85 and 15.06 ± 0.90 , respectively (Table 2).

In general, it was monitored from the beginning of the sampling period, the population density of TSSM gradually decreased on all the four rose varieties, and for the relatively long time (from the 10th to the 24th week) it was fluctuating at very low level, but from the twenty-fourth week onward, the population density increased (Table 2).

Population fluctuation of P. persimilis

The population of P. persimilis was monitored from the beginning of the first release during the first week (Table 3). The highest population density was recorded on Avalanche in overall dates, followed by Samurai, Sorbet and Dolcevita $(0.92 \pm 0.07, 0.88 \pm 0.08, 0.56 \pm 0.06,$ and 0.25 ± 0.03 mite/leaf, respectively). Due to the subsequent releases of this predator during the first, third, fourth and fifth weeks (Table 1), its population density increased gradually on all the four rose varieties, and when the population density of the predator peaked, with a steep slope decreased to lower levels (Table 3). With the release of this predator in the seventeenth week (Table 1), no predator was observed in any of the varieties examined but by the re-release on other dates, the predator population density gradually increased (Table 3).

Population fluctuation of A. swirskii

The population density of *A. swirskii* per leaf on all rose varieties at different dates is shown in Table 4. Mean density of *A. swirskii* until last date of its observation in the greenhouse (twelfth week) on overall dates for Avalanche, Dolcevita, Samurai and Sorbet varieties was record 0.10 ± 0.007 , 0.05 ± 0.005 , 0.04 ± 0.005 , 0.07 ± 0.006 mite/leaf, respectively. Therefore, the highest population density of this predator per leaf was recorded on Avalanche (0.10 ± 0.007) and the lowest population density was observed on Samurai (0.04 ± 0.005).

Sampling	Ň	Б	đf	D				
date (week)	Avalanche	Dolcevita	Samurai	Sorbet	- г	ui	Р	
06/09/2016 (1)	52.2 ± 5.89^{a}	$21.13\pm2.83^{\mathrm{b}}$	52.8 ± 4.51^{a}	16.6 ± 1.98^{b}	54.87	3,119	0.001	
13/09/2016 (2)	37.77 ± 5.71^{a}	7.33 ± 2.13^{b}	47.6 ± 5.55^a	13.33 ± 1.49^{ab}	38.28	3,119	0.001	
20/09/2016 (3)	68.13 ± 5.38^a	21.7 ± 0.54^{b}	63.53 ± 5.01^{a}	26.67 ± 1.41^{ab}	45.21	3,119	0.001	
27/09/2016 (4)	33.93 ± 3.70^{ab}	$8.77 \pm 1.42^{\rm c}$	43.17 ± 4.26^a	16.97 ± 1.35^{b}	35.51	3,119	0.001	
04/10/2016 (5)	32.5 ± 2.89^a	$4.63 \pm 1.12^{\rm c}$	35.8 ± 4.10^{a}	8.4 ± 0.79^{b}	27.22	3,119	0.021	
11/10/2016 (6)	30.73 ± 3.54^{a}	0.07 ± 0.02^{b}	27.2 ± 1.42^{a}	$2.07 \pm 1.58^{\text{b}}$	74.68	3,119	0.001	
18/10/2016 (7)	30.63 ± 2.60^a	$0.33\pm0.11^{\rm c}$	26.97 ± 3.02^a	12.33 ± 2.52^{b}	31.10	3,119	0.001	
25/10/2016 (8)	$12.5 \pm 1.2^{\text{b}}$	$0.03\pm0.01^{\circ}$	27.27 ± 4.49^a	$0.77\pm0.32^{\rm c}$	9.12	3,119	0.014	
01/11/2016 (9)	9.57 ± 2.1^{a}	$0.03\pm0.01^{\text{b}}$	$10.7 \pm 1.85^{\rm a}$	0	8	3,119	0.001	
08/11/2016 (10)	$4.07\pm0.82^{\text{a}}$	0	$0.2\pm0.08^{\text{b}}$	0	11.51	3,119	0.001	
15/11/2016 (11)	6.97 ± 1.40^{a}	0	0	0	17	3,119	0.048	
22/11/2016 (12)	$0.6\pm0.35^{\text{a}}$	0	0	0	8	3,119	0.012	
29/11/2016 (13)	2.23 ± 0.81^{a}	0	0	0	11	3,119	0.037	
06/12/2016 (14)	$1.83\pm0.60^{\text{a}}$	0	0	0	6	3,119	0.041	
13/12/2016 (15)	0	0	0	0	-	-	-	
20/12/2016 (16)	0	0	0	0	-	-	-	
27/12/2016 (17)	0	0	0.27 ± 0.12^{a}	0.73 ± 0.43^{a}	5.61	3,119	0.001	
03/01/2017 (18)	$0.37\pm0.10^{\text{a}}$	$0.17\pm0.09^{\rm a}$	0.87 ± 0.56^{a}	$2.07 \pm 1.01^{\text{a}}$	12.30	3,119	0.001	
10/01/2017 (19)	0.63 ± 0.23^{b}	$4.8\pm0.93^{\text{a}}$	0.6 ± 0.47^{b}	3.53 ± 1.06^{a}	24.91	3,119	0.001	
17/01/2017 (20)	0.33 ± 0.13^{b}	0.30 ± 0.14^{b}	5.7 ± 1.20^{a}	$5.73 \pm 1.02^{\rm a}$	12.20	3,119	0.034	
24/01/2017 (21)	$4.17\pm0.60^{\text{a}}$	3.1 ± 0.60^{a}	0.77 ± 0.50^{b}	$3.4\pm1.5^{\rm a}$	9	3,119	0.001	
31/01/2017 (22)	0.07 ± 0.06^{b}	$0.03\pm0.03^{\text{b}}$	0	$3.6\pm1.8^{\rm a}$	11	3,119	0.001	
07/02/2017 (23)	8.70 ± 2.70^{a}	$0.23\pm0.23^{\rm c}$	2.67 ± 0.41^{b}	2.57 ± 0.43^{b}	5	3,119	0.011	
14/02/2017 (24)	2.93 ± 1.39^{b}	6.97 ± 1.68^{a}	0	6.73 ± 1.65^{a}	4	3,119	0.041	
21/02/2017 (25)	$3.23 \pm 1.2^{\rm c}$	$3.37\pm0.47^{\rm c}$	22 ± 1.01^{a}	$10.17 \pm 1.54^{\rm b}$	15	3,119	0.001	
28/02/2017 (26)	$5.27\pm0.99^{\circ}$	$15.67\pm1.95^{\text{b}}$	35.67 ± 2.37^a	32.53 ± 3.45^a	17.31	3,119	0.001	
07/03/2017 (27)	17.03 ± 4.07^{b}	13.87 ± 2.52^{b}	39.73 ± 3.03^a	33.37 ± 2.83^a	8	3,119	0.001	
14/03/2017 (28)	15.43 ± 5.37^{b}	$24.93 \pm 4.54^{\text{b}}$	39.57 ± 4.38^a	37.6 ± 3.46^{a}	12.11	3,119	0.048	
21/03/2017 (29)	$18.17\pm2.97^{\rm c}$	$16.4\pm2.87^{\rm c}$	40.37 ± 4.84^{b}	$54.87\pm5.37^{\mathrm{a}}$	10.68	3,119	0.001	
28/03/2017 (30)	21.9 ± 5.46^{b}	$10.67\pm2.18^{\rm c}$	15.27 ± 0.96^{b}	$69.2\pm3.10^{\rm a}$	5.62	3,119	0.001	
04/04/2017 (31)	$12.1\pm2.06^{\rm c}$	$0.37\pm0.19^{\rm d}$	18.17 ± 0.78^{b}	$103.43\pm2.58^{\rm a}$	14.22	3,119	0.001	
Overall dates	14 ± 0.73^{b}	$5.32\pm0.39^{\rm c}$	17.96 ± 0.85^{a}	15.06 ± 0.90^{b}	9.12	3,3479	0.001	

Table 2 Population density of *Tetranychus urticae* on four rose varieties.

 $\overline{\text{Means within a row followed by different letters are significantly different (Tukey test, P<0.05)}.$

Sampling	Numb	er of predatory n	Б	đf	D			
date (week)	Avalanche	Dolcevita	Samurai	Sorbet	- Г	di	Ľ	
06/09/2016 (1)	$0.23\pm0.10_a$	0.27 ± 0.12^{a}	0.23 ± 0.09^{a}	$0.1\pm0.06^{\rm a}$	8	3,119	0.011	
13/09/2016 (2)	0	0.07 ± 0.04^{a}	0.03 ± 0.03^{a}	0	9.11	3,119	0.035	
20/09/2016 (3)	0.13 ± 0.08^{ab}	0.07 ± 0.05^{b}	0.17 ± 0.09^{ab}	$0.47\pm0.17^{\rm a}$	6	3,119	0.00	
27/09/2016 (4)	$0.93\pm0.20^{\rm a}$	0.23 ± 0.12^{b}	0.33 ± 0.16^{ab}	0.73 ± 0.23^{ab}	4	3,119	0.00	
04/10/2016 (5)	1.20 ± 0.25^{ab}	0.47 ± 0.18^{b}	0.7 ± 0.24^{ab}	1.4 ± 0.31^{a}	8.12	3,119	0.013	
11/10/2016 (6)	$2.10\pm0.30^{\rm a}$	0.53 ± 0.15^{b}	0.43 ± 0.15^{b}	0.83 ± 0.19^{b}	14	3,119	0.00	
18/10/2016 (7)	$3.93\pm0.46^{\rm a}$	$0.83\pm0.22^{\rm c}$	2.1 ± 0.30^{b}	1.87 ± 0.40^{b}	12	3,119	0.00	
25/10/2016 (8)	$2.53\pm0.31^{\text{b}}$	0	$4.53\pm0.65^{\rm a}$	$1.6\pm0.32^{\rm c}$	5	3,119	0.00	
01/11/2016 (9)	2.37 ± 0.29^{b}	0	$5.67 \pm 1.05^{\rm a}$	$0.1\pm0.06^{\rm c}$	6.71	3,119	0.00	
08/11/2016 (10)	0.83 ± 0.22^{a}	0	$0.57\pm0.18^{\rm a}$	0	8.21	3,119	0.00	
15/11/2016 (11)	$0.3\pm0.15^{\rm a}$	0	0.17 ± 0.11^{a}	0	8.25	3,119	0.028	
22/11/2016 (12)	0.63 ± 0.22^{a}	0	0	0	7	3,119	0.016	
29/11/2016 (13)	$0.37\pm0.16^{\rm a}$	0	0	0	9.31	3,119	0.032	
06/12/2016 (14)	$0.27\pm0.15^{\rm a}$	0	0	0	4.96	3,119	0.041	
13/12/2016 (15)	0	0	0	0	-	-	-	
20/12/2016 (16)	0	0	0	0	-	-	-	
27/12/2016 (17)	0	0	0	0	-	-	-	
03/01/2017 (18)	0	0	0	0	-	-	-	
10/01/2017 (19)	0	0	0	0	-	-	-	
17/01/2017 (20)	0	0	0	0	-	-	-	
24/01/2017 (21)	0	0	0	0	-	-	-	
31/01/2017 (22)	0	0	0	0	-	-	-	
07/02/2017 (23)	0	0	0	0	-	-	-	
14/02/2017 (24)	0	0	0	0	-	-	-	
21/02/2017 (25)	$0.03\pm0.00^{\rm a}$	0	0	0	5	3,119	0.048	
28/02/2017 (26)	$0.17\pm0.00^{\rm a}$	0	0	0	5.81	3,119	0.033	
07/03/2017 (27)	0	0.2 ± 0.12^{a}	0	$0.13\pm0.09^{\rm a}$	9	3,119	0.010	
14/03/2017 (28)	0.23 ± 0.11^{ab}	0.43 ± 0.16^a	0.3 ± 0.15^{ab}	$0.07\pm0.05^{\rm b}$	12	3,119	0.036	
21/03/2017 (29)	2.1 ± 0.37^{a}	2.43 ± 0.38^a	1.53 ± 0.31^{ab}	$0.3\pm0.14^{\rm b}$	13.84	3,119	0.00	
28/03/2017 (30)	$7.8\pm0.85^{\rm a}$	$0.73\pm0.18^{\rm c}$	3.37 ± 0.36^{b}	$2.77\pm0.45^{\text{b}}$	7.62	3,119	0.00	
04/04/2017 (31)	2.43 ± 0.44^{b}	$1.5\pm0.38^{\rm c}$	7.2 ± 0.64^{a}	6.93 ± 0.81^{a}	14	3,119	0.00	
Overall dates	0.92 ± 0.07^{a}	0.25 ± 0.03^{c}	$0.88\pm0.08^{\rm a}$	0.56 ± 0.06^{b}	7.3	3,2519	0.00	

Table 3 Population density of Phytoseiulus persimilis on four rose varieties.

Means within a row followed by different letters are significantly different (Tukey test, P<0.05).

The population of *A. swirskii* was monitored from the beginning of the first release during the second week. The population density of this predator increased on all four rose varieties, but it was dramatically reduced after a few weeks. Finally, after the twelfth week, no density of this predator was observed on any of the studied varieties (Table 4).

Population fluctuation of N. californicus

The mean population density of *N. californicus* per leaf of different rose varieties on each sampling date is shown in Table 5. There was a significant difference in density of *N. californicus* among different rose varieties at different dates. The population density of this

predator during the overall date for Avalanche, Dolcevita, Samurai and Sorbet varieties was obtained 0.04 ± 0.007 , 0.07 ± 0.01 , 0.01 ± 0.004 , 0 and 0.05 ± 0.008 mite/leaf, respectively.

The N. californicus population was monitored from the beginning of its release during the twelfth week that after six weeks, the first individuals were observed in samurai variety. Subsequently, the predator population was observed very sparse and found at very low levels on different varieties (Table 5). After the second release of this predator in the twenty-sixth week (Table 1), there was an increasing trend but fluctuation was observed in the predator's population (Table 5).

Table 4 Population density of Amblyseius Swirskii on four rose varieties.

Sampling date	Numb	er of predatory m	Ē	đf	D		
(week)	Avalanche	Dolcevita	Samurai	Sorbet	Г	ui	г
06/09/2016 (2)	$0.067\pm0.05^{\rm a}$	0	0	0.03 ± 0.03^{a}	9.00	3,119	0.046
13/09/2016 (3)	0.13 ± 0.08^{a}	0.03 ± 0.03^{a}	$0.07\pm0.07^{\rm a}$	0.17 ± 0.08^{a}	13.20	3,119	0.034
20/09/2016 (4)	$0.13\pm0.08^{\rm a}$	$0.27\pm0.13^{\rm a}$	$0.07\pm0.05^{\rm a}$	$0.13\pm0.08^{\rm a}$	6.44	3,119	0.011
27/09/2016 (5)	$0.17\pm0.08^{\rm a}$	0.23 ± 0.1^{a}	0.23 ± 0.11^{a}	0.2 ± 0.14^{a}	8.10	3,119	0.00
04/10/2016 (6)	0.37 ± 0.11^{a}	0.03 ± 0.03^{b}	0	0.1 ± 0.07^{b}	7.98	3,119	0.014
11/10/2016 (7)	0.27 ± 0.11^{a}	0	0	0.067 ± 0.06^{a}	9.33	3,119	0.00
18/10/2016 (8)	0	0.03 ± 0.03^{a}	$0.1 \pm 0.07^{\mathrm{a}}$	0	5.55	3,119	0.029
25/10/2016 (9)	0	0	$0.03\pm0.03^{\rm a}$	0	6.39	3,119	0.017
01/11/2016 (10)	0	0	0	$0.03\pm0.03^{\rm a}$	7.14	3,119	0.031
08/11/2016 (11)	$0.03\pm0.03^{\rm a}$	0	0	0	8.00	3,119	0.024
15/11/2016 (12)	0	0	0	$0.07\pm0.06^{\rm a}$	5.21	3,119	0.042
Overall dates	0.10 ± 0.007^{a}	0.05 ± 0.005^{b}	0.04 ± 0.005^{b}	0.07 ± 0.006^{b}	12.06	3,1319	0.047

Means within a row followed by different letters are significantly different (Tukey test, P< 0.05).

Someling data (wash)	Numbe	er of predatory n	Б	ағ	D			
Sampling date (week)	Avalanche	Dolcevita	Samurai	Sorbet	- Г	ui	Г	
22/11/2016 (12)	0	0	0	0	-	-	-	
29/11/2016 (13)	0	0	0	0	-	-	-	
06/12/2016 (14)	0	0	0	0	-	-	-	
13/12/2016 (15)	0	0	0	0	-	-	-	
20/12/2016 (16)	0	0	0	0	-	-	-	
27/12/2016 (17)	0	0	0	0	-	-	-	
03/01/2017 (18)	0	0	$0.1\pm0.07^{\rm a}$	0	6.05	3,119	0.027	
10/01/2017 (19)	0	0.03 ± 0.01^{a}	0	0	10.44	3,119	0.012	
17/01/2017 (20)	0.07 ± 0.04^{a}	$0.07\pm0.02^{\rm a}$	0	0	6.12	3,119	0.023	
24/01/2017 (21)	0	0	0	0	-	-	-	
31/01/2017 (22)	0	0	0.03 ± 0.01^{a}	0	9.11	3,119	0.035	
07/02/2017 (23)	0	0	0	0	-	-	-	
14/02/2017 (24)	0	0	0	0.03 ± 0.01^{a}	5.08	3,119	0.039	
21/02/2017 (25)	0	0	0	0	-	-	-	
28/02/2017 (26)	0.07 ± 0.04^{a}	$0.07\pm0.07^{\rm a}$	0	0	7.81	3,119	0.042	
07/03/2017 (27)	0.03 ± 0.01^{b}	$0.53\pm0.16^{\rm a}$	0.03 ± 0.01^{b}	0.43 ± 0.15^{ab}	5.69	3,119	0.00	
14/03/2017 (28)	$0.23\pm0.10^{\rm a}$	0.13 ± 0.06^{ab}	0	0	13.04	3,119	0.00	
21/03/2017 (29)	0	$0.17\pm0.14^{\rm a}$	$0.03\pm0.03^{\rm a}$	0.1 ± 0.07^{a}	11.23	3,119	0.00	
28/03/2017 (30)	0.03 ± 0.01^{a}	$0.3\pm0.13^{\rm a}$	0	0.1 ± 0.07^{a}	9.00	3,119	0.00	
04/04/2017 (31)	0.33 ± 0.14^{a}	0.13 ± 0.08^{b}	$0.07\pm0.06^{\rm b}$	$0.43\pm0.15^{\rm a}$	14.21	3,119	0.00	
Overall dates	0.04 ± 0.007^{ab}	0.07 ± 0.01^{a}	0.01 ± 0.004^{b}	0.05 ± 0.008^{ab}	9.85	3,1319	0.038	

Table 5 I opulation density of <i>Neosetulus cultornicus</i> on four fose varietie	Tał	ble	5	Pop	pul	atio	n (density	of	1	Ve	ose	eiu	lus	са	lif	orn	icus	01	1 İ	four	rose	v	ariet	ies	
---	-----	-----	---	-----	-----	------	-----	---------	----	---	----	-----	-----	-----	----	-----	-----	------	----	-----	------	------	---	-------	-----	--

Means within a row followed by different letters are significantly different (Tukey test, P< 0.05).

Discussion

Different plant varieties have a different effect on the biological parameters of herbivorous species and their predators (Khanamani *et al.*, 2014). In this study, the differences observed in population density of TSSM and its predators are probably due to the leaf structure, growth characteristics, and the defense mechanisms of the rose varieties. This means that four rose varieties significantly affected the biology of TSSM. The effect of leaf structure or growth characteristics of the rose varieties on population density of TSSM and its predators was not examined in this study. Thus further work is necessary to determine whether the differences observed in population density of TSSM and its predators are due to the leaf structure or caused by growth characteristics of the plants. Skirvin and Williams (1999) reported that morphological structure of rose varieties can affect the population density of TSSM. Similar conclusions were also made by other researchers (Agrawal, 2000; Balkema-Boomstra *et al.*, 2003; Biswas *et al.*, 2004; Peralta and Tello, 2011; Schmidt, 2014).

The results showed that the population of *P*. persimilis was increased consistently in all four varieties with decreasing population of TSSM (Tables 2 and 3). A severe decline in predator density immediately after peak population was observed on four rose varieties, which indicates that there is not sufficient prey to feed the predator (Table 3). Due to the fact that P. persimilis is a specialized predator of TSSM, its density is reduced significantly in the absence of sufficient prey. These results were similar to results obtained by Blumel and Walzer (2002) on other varieties of this host plant. Also Gacheri et al. (2015) and Chacón-Hernández et al. (2016) reported that a severe decline occurs in density of predator by decreasing the density of TSSM. Hot-spot applications of *P. persimilis* in the seventeenth week was not effective in decreasing density of TSSM (Tables 1 and 3). Therefore, it is concluded that although P. persimilis has been successful in controlling the high density of TSSM, it does not have the ability to control the population of the pest in lower densities. This disagrees with that concluded by Gacheri et al. (2015) in rose farms in Kenya.

Generally, the population density of TSSM decreased from the beginning of sampling period and reached near zero per leaf in the tenth week and this trend continued for relatively long time (until the 24th week) (Table 2). The low density of the TSSM population from the 10th to the 24th week could mainly be due to the N. californicus activity in this period (Tables 2 and 5). Due to the prey preferences of N. californicus on TSSM (Fathipour and Maleknia, 2016) and according to Tables 2 and 5, it seems that the low density of TSSM in different varieties from the 10th to the 24th week has prevented its population growth. Therefore, it is a good option for TSSM control in low density, and seems that its release is very suitable. Blumel and Walzer (2002) reported that N. californicus has the ability to survive in low prey density, but the lack of prey prevents its population density growth.

We did not find a definite cause for reduction of *A. swirskii* population (Table 4). It

is probable that the *A. swirskii* has not been able to compete with the *P. persimilis* and it is possible that the feeding patch has left. This is a general predator and may have changed the feeding patch in search for another food source, such as thrips and whitefly (Messelink and

Janssen, 2008; van Maanen *et al.*, 2012). In conclusion, the obtained results showed that the predatory mite *P. persimilis* had the ability to reduce the high TSSM density and *N. californicus* also continued to operate in low TSSM density, but *A. swirskii* did not have a clear impact on reducing TSSM density.

Acknowledgments

The support of this research by the Department of Entomology, Tarbiat Modares University, is greatly appreciated. The authors also thank Seyed Javad Hajimir manager of the Yaran greenhouse for providing the study site.

References

- Agrawal, A. A. 2000. Host-range evolution: adaptation and tradeoffs in fitness of mites on alternative hosts. Ecology, 81: 500-508.
- Alipour, Z., Fathipour, Y. and Farazmand, A. 2016. Age-stage predation capacity of *Phytoseiulus persimilis* and *Amblyseius swirskii* (Acari: Phytoseiidae) on susceptible and resistant rose cultivars, International Journal of Acarology, 42: 224-228.
- Attia, S., Grissa, K. L., Lognay, G., Bitume, E., Hance, T. and Mailleux, C. 2013. A review of the major biological approaches to control the worldwide pest *Tetranychus urticae* (Acari: Tetranychidae) with special reference to natural pesticides. Journal of Pest Science, 86 (3): 361-386.
- Balkema-Boomstra, A. G., Zijlstra, S., Verstappen, F. W. A., Inggamer, H., Mercke, P. E., Jongsma, M. A. and Bouwmeester, H. J. 2003. Role of Cucurbitacin C in resistance to spider mite (*Tetranychus urticae*) in cucumber (*Cucumis* sativus L.). Journal of Chemical Ecology, 29: 225-235.

- Biswas, G. C., Islam W., Haque M. M., Saha R.
 K., Hoque K. M. F., Islam M. S. and Haque
 M. E. 2004. Some biological aspects of Carmine Mite, *Tetranychus cinnabarinus* Boisd. (Acari: Tetranychidae) infesting eggplant from Rajshahi. Journal of Biological Sciences, 4: 588-591.
- Blumel, S. and Walzer, A. 2002. Efficacy of different release strategies of *Neoseiulus californicus* McGregor and *Phytoseiulus persimilis* Athias Henriot (Acari: Phytoseiidae) for the control of two-spotted spider mite (*Tetranychus urticae* Koch) on greenhouse cut roses. Systematic and Applied Acarology, 7: 35-48.
- Buitenhuis, R., Murphy G., Shipp, L. and Scott-Dupree, C. 2015. *Amblyseius swirskii* in greenhouse production systems: a floricultural perspective. Experimental and Applied Acarology, 65: 451-464.
- Casey, C., Newman, J., Robb, K., Tjosvold, S., Macdonald, J. and Parrella, M. 2007.
 IPM program successful in California greenhouse cut roses. California Agriculture, 61 (2): 71-78.
- Chacón-Hernández, J., Cerna-Chávez, E., Ochoa-Fuentes, Y., Hernández-Juárez, A., Palacios-Bazaldúa, L. and Landeros-Flores J. 2016. Spatial distribution and population fluctuation of *Tetranychus urticae* (Koch) and *Phytoseiulus persimilis* (Athias-Henriot) in rose crops. Phytology, 85: 236-245.
- De Vis, R. and Barrera, J. 1999. Use of two predators *Phytoseiulus persimilis* Athias-Henriot (Acari: Phytoseiidae) and *Neoseiulus californicus* (McGregor) (Acari: Phytoseiidae) for the biological control of *Tetranychus urticae* Koch (Acari: Tetranychidae) in roses in the Bogota´ plateau. Acta Horticulture, 482: 259-268.
- Escudero, L.A. and Ferragut, F. 2005. Life-history of the predatory mites *Neoseiuluscalifornicus* and *Phytoseiulus persimilis* (Acari: Phytoseiidae) on four spider mite species and prey, with special reference to *Tetranychus evansi* (Acari: Tetranychidae). Biological Control, 32: 378-384.

- Fathipour, Y. and Maleknia, B. 2016. Mite predators. In: Omkar (Ed.), Ecofriendly Pest Management for Food Security. Elsevier, San Diego, USA, pp: 329-366.
- Fathipour, Y. and Mirhosseini, M. A. 2017.
 Diamondback Moth (*Plutella xylostella*) Management. In: Reddy, G. V. P. (Ed.), Integrated Management of Insect Pests on Canola and Other Brassica Oilseed Crops.CABI, Croydon, UK, PP: 13-43.
- Gacheri, C., Kigen, T. and Sigsgaard, L. 2015. Hot-spot application of biocontrol agents to replace pesticides in large scale commercial rose farms in Kenya. BioControl, 60: 795-803.
- Gerson, U. and Weintraub, P. G. 2007. Mites for the control of pests in protected cultivation. Pest Management Science, 63: 658-676.
- Holt, K. M., Opit, G. P., Nechols, J. R., Margolies, D. C. and Williams, K. A. 2007. Comparing chemical and biological control strategies for twospotted spider mites on mixed production of ivy geranium and impatiens. HortTechnology, 17: 322-327.
- Khajehali, J., Nieuwenhuyse, P. V., Demaeght, P., Tirry, L. and Leeuwen, T. V. 2011. Acaricide resistance and resistance mechanisms in *Tetranychus urticae* populations from rose greenhouses in the Netherlands. Pest Management Science, 67: 1424-1433.
- Khanamani, M., Fathipour, Y. and Hajiqanbar, H. 2013. Population growth response of *Tetranychus urticae* to eggplant quality: application of female age-specific and agestage, two-sex life tables. International Journal of Acarology, 39: 638-648.
- Khanamani, M., Fathipour, Y., Hajiqanbar, H. and Sedaratian, A. 2014. Two-spotted spider mite reared on resistant eggplant affects consumption rate and life table parameters of its predator, *Typhlodromus bagdasarjani* (Acari: Phytoseiidae). Experimental and Applied Acarology, 63: 241-252.
- Maleknia, B., Fathipour, Y. and Soufbaf, M. 2016. How greenhouse cucumber cultivars affect population growth and two-sex life table parameters of *Tetranychus urticae*

(Acari: Tetranychidae). International Journal of Acarology, 42: 70-78.

- McMurtry, J. A., De Moraes, G. J. and Sourassou, N. F. 2013. Revision of the lifestyles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. Systematic and Applied Acarology, 18: 297-320.
- Messelink, G. J., Maanen, R., van Steenpaal, S. E. F. and Janssen, A. 2008. Biological control of thrips and whiteflies by a shared predator: two pests are better than one. Biological Control, 44: 372-379.
- Nicetic, O., Watson, D. M., Beattie, G. A. C., Meats, A. and Zheng, J. 2001. Integrated pest management of two-spotted mite *Tetranychus urticae* on greenhouse roses using petroleum spray oil and the predatory mite *Phytoseiulus persimilis*. Experimental and Applied Acarology, 25: 37-53.
- Opit, G. P., Nechols, J. R. and Margolies, D. C. 2004. Biological control of two spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae), using *Phytoseiulus persimilis* (Acari: Phytoseidae) on ivy geranium: assessment of predator release ratios. Biological Control, 29: 445-452.

- Peralta, O. and Tello, V. 2011. Tabla de vida de *Tetranychus cinnabarinus* (Acari: Tetranychidae) sobre tres variedades de melón. Cucumis melo. Revista Colombiana de Entomología, 37: 21-26.
- Ragusa, S. and Swirski, E. 1975. Feeding habits, development and oviposition of the predacious mite *Amblyseius swirskii* Athias-Henriot (Acarina: Phytoseiidae) on pollen of various weeds. Israel Journal of Entomology, 10: 93-103.
- Schmidt, A. 2014. Leaf structures affect predatory mites (Acari: Phytoseiidae) and biological control: a review. Experimental and Applied Acarology, 62: 1-17.
- Skirvin, D. J. and Williams, M. 1999.
 Differential effects of plant species on a mite pest (*Tetranychus Urticae*) and its predator (*Phytoseiulus Persimilis*):
 Implications for biological control. Experimental and Applied Acarology, 23: 497-512.
- Van Maanen, R., Broufas, G., Oveja, M. F., Sabelis, M. W. and Janssen, A. 2012. Intraguild predation among plant pests: western flower thrips larvae feed on whitefly crawlers. BioControl, 57: 533-539.

کنترل Tetranychus urticae با استفاده از سه گونه شکارگر فیتوزیید در یک گلخانه تجاری رز

كمال الدين احمدي بيداخويدي، يعقوب فتحي پور* و محمد بشيري

گروه حشرهشناسی کشاورزی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران. پست الکترونیکی نویسندگان مسئول مکاتبه: fathi@modares.ac.ir دریافت: ۱۴ اردیبهشت ۱۳۹۸؛ پذیرش: ۱۹ آبان ۱۳۹۸

چکیدہ: کنه تارتن دولکهای (Tetranychus urticae Koch (Acari: Tetranychidae) یک آفت جهانی برای بسیاری از گونههای گیاهی مانند خیار، سویا، رز و غیره میباشد. کنترل کنههای تارتن عمدتاً بر اساس استفاده از آفت کشهای شیمیایی صورت می گیرد. شکار گرهای فیتوزیید برای کنترل کنههای آفت و هم-چنین برخی از حشرات استفاده شدهاند. در این مطالعه، کنترل کنه تارتن دولکهای با استفاده از کنههای شكارگر فيتوزييد Amblyseius swirskii Athias-Henriot ، Phytoseiulus persimilis Athias-Henriot و Neoseiulus californicus McGregor روى چهار رقم رز شامل Samurai ،Dolcevita ،Avalanche و ارزیابی شد. مجموع تعداد تخم و کل مراحل متحرک هر یک از کنههای مورد مطالعه با استفاده از یک لنز دستی در واحد برگ و بهصورت هفتگی شمارش شد. نمونهبرداری بهصورت تصادفی انجام گرفت و برگهای یک بوته رز از تاج پوشش تحتانی (شاخههای خم شده)، تاج پوششمیانی (بین شاخههای پایینی و بالایی) و تاج پوشش بالایی (شاخههای گلدهنده) انتخاب شدند. در مجموع تعداد ۳۰ نمونه برگ در هر بار نمونه-برداری جمع آوری شد. بیشترین میانگین جمعیت کنه تارتن دولکهای بهازای هر برگ روی رقم Samurai (۱۷/۹۶ ± ۱۷/۹۶) و کمترین تراکم آن روی رقم Dolcevita (۵/۳۹ ± ۱۷/۹۶) ثبت گردید. بر اساس دادههای حاصل از تغییرات جمعیت کنه تارتن دولکهای و شکارگرهای آن روی چهار رقم رز، شکارگر P. persimilis به-عنوان یک عامل بیولوژیک مؤثر در کاهش تراکم بالای کنه تارتن دولکهای شناخته شد. همچنین شکارگر .N californicus توانایی حفظ دوام خود در جمعیت پایین کنه تارتن دولکهای را داشت اما شکارگر A. swirskii تأثير آشکاری بر کنترل کنه تارتن دولکهای نداشت.

واژگان كليدى: شكارگرهاى فيتوزييد، كنه تارتن دولكهاى، پويايى جمعيت، مديريت تلفيقى كنه آفت