

Research Article

Characterization and genetic variability of coat protein gene of *Apple chlorotic leaf spot virus* isolates from apple in Kurdistan and western Azarbaijan Provinces, Iran

Kamal Shujaei, Mahdi Azaryar and Mohammad Hajizadeh*

Department of Plant Protection, Faculty of Agriculture, University of Kurdistan, Sanandaj, Iran.

Abstract: In this study, 174 leaf samples from main apple growing areas in Kurdistan and western Azarbaijan Provinces were examined for infection by *Apple chlorotic leaf spot virus* (ACLSV). The result of RT-PCR showed that 17 samples (9.7%) were infected by ACLSV in both provinces. Subsequently, the genetic structure of ACLSV was determined based on the full coat protein (CP) sequences of eight isolates from this study and those recently published from Iran and worldwide. In phylogenetic analysis, ACLSV isolates from apple clustered into three known phylogenetic types; B6, P205, and SHZ, which are characterized by three amino acids at positions of 40, 75 and 79. The new Iranian ACLSV isolates showed the CP amino acid sequence of Ser⁴⁰-Tyr⁷⁵-Glu⁷⁹ which placed them with the B6 type. Further analyses showed that the Iranian ACLSV isolates (the new and previously published isolates) had lower genetic variability compared to isolates from other countries except for Brazilian population. Most of the CP codons were under negative selection ($dN/dS < 1$) except for one codon at position 92 which was under positive selection. This study suggests that negative selection was most important evolutionary factor driving the genetic evolution of ACLSV.

Keywords: apple, negative selection, phylogenetic analysis, ACLSV

Introduction

Apple is commercially the most important fruit crop and has the most exporting value in Iran. Iran stands at 4th and 7th positions in orchard area and apple production with 0.24 mHa and 2.45 mt, respectively (FAOSTAT, 2016). Viruses as biotic factor are the main agent in low productivity of apple worldwide (Watpade *et al.*, 2012). Among various viral diseases, ACLSV is widely distributed in Rosaceae fruit trees such as apple, apricot, cherry and pear, usually causing a latent

infection in most commercial cultivars (Watpade *et al.*, 2012). However, some apple cultivars are susceptible and may develop severe symptoms. The severity of symptoms induced by ACLSV depends on host plant species and virus strain (Nemeth, 1986). infection rates of up to 80-100% with yield losses in the order of 30-40% by thid virus have been reported (Nemchinov *et al.*, 1995; Wu *et al.*, 1998; Cembali *et al.*, 2003).

ACLSV is the type member of the genus *Trichovirus*, subfamily *Trivirinae*, family *Betaflexiviridae*, within the order *Tymovirales* (Adams *et al.*, 2016). ACLSV has a filamentous particle, composed of a positive-sense, single-stranded RNA molecule of about 7.5 kb excluding the polyadenylated tail (Yoshikawa, 1988; Yoshikawa *et al.*, 1992). The genome of ACLSV

Handling Editor: Masoud Shams-bakhsh

*Corresponding authors, e-mail: m.hajizadeh@uok.ac.ir
Received: 4 August 2019, Accepted: 27 October 2019
Published online: 11 November 2019

contains three overlapping open reading frames (ORF), which encode a 216-kDa replication-associated protein, a 50-kDa movement protein which overlaps both replicase and CP coding regions, and a 22-kDa coat protein (CP), respectively (Sato *et al.*, 1993; Niu *et al.*, 2012).

Molecular variation and characterization were reported for the ACLSV isolates from India, China and Japan (Al Rwahnih *et al.*, 2004; Yaegashi *et al.*, 2007; Rana *et al.*, 2010; Chen *et al.*, 2014). Considering that Iran is the 7th largest producer of apple around the world, a greater understanding of ACLSV variability is necessary, in particular with regard to isolates from apple. In Iran, ACLSV has been reported from several provinces (Derakhshian *et al.*, 2010; Farmahini *et al.*, 2014; Keshavarz and Shams-Bakhsh, 2015; Alemzadeh *et al.*, 2016; Abtahi *et al.*, 2019) and previous surveys have indicated that 2.67% of pear (Farmahini *et al.*, 2014) and 18.55% of apple (Keshavarz and Shams-Bakhsh, 2015) trees are infected with ACLSV. Recently, Abtahi *et al.* (2019) showed that Iranian ACLSV isolates belonged to B6 and P205 phylogenetic groups based on partial CP gene sequences. So far, all Iranian isolates of ACLSV CP have been partially sequenced (Keshavarz and Shams-Bakhsh, 2015; Naderpour *et al.*, 2016; Abtahi *et al.*, 2019) and complete CP gene sequence of ACLSV has not been previously determined. Also, despite of the detection of ACLSV in some provinces of Iran, no information is available on presence of ACLSV in Kurdistan Province. In this work, we performed an updated analysis of the genetic variation and phylogenetic relationship of Iranian ACLSV isolates by analysis of the recently published coat protein gene sequences (Abtahi *et al.*, 2019) and those investigated in the present study.

Materials and Methods

Source of isolates

During July, August, and September 2017 and 2018, 174 symptomatic and non-symptomatic apple leaf samples were collected from several locations of Kurdistan and Western Azarbaijan Provinces of Iran.

RT-PCR detection of ACLSV

Total nucleic acid (TNA) was extracted from one leaf per sample based on the method described by Foissac *et al.* (2000) with minor modifications. Reverse transcription (RT) reactions were performed by a cDNA synthesis kit (HyperScriptTM Reverse Transcriptase, GeneAll, South Korea) using random hexamer primer. The synthesised cDNA was subjected to PCR amplification in a 12.5 µl PCR master mix (GeneAll, South Korea) and 10 pmol of each primer, ACLS-MF: 5' GAGGATCCATGGC GGCAG' and ACLSV-MR: 5' TACTCG AGGTAAATGCAAAGATCAG 3' (designed in this study), In PCR master mix. PCR products were electrophoresed on a 1.2% (w/v) agarose gel (containing 0.5 µg/ml EtBr) in 1X TAE buffer.

Cloning and sequencing of RT-PCR products

RT-PCR products of eight apple trees were purified using a Nucleic Acid Purification Kit (GeneAll, South Korea), ligated into the pTG19 cloning vector (Vivantis, Malaysia), and the ligation mix was transformed into *Escherichia coli* strain DH5α as described by Chung *et al.* (1989). The recombinant plasmids were confirmed by restriction analysis and a purified clone from each isolate was subjected to sequencing by Macrogen Inc. (Seoul, South Korea).

Multiple sequence alignment and phylogenetic analysis

Multiple sequence alignment of ACLSV CP sequences, including eight sequences of ACLSV-CP obtained in this study and additional sequences retrieved from GenBank was performed separately using Clustal W (Thompson *et al.*, 1997) implemented in MEGA X (Kumar *et al.*, 2018) with default parameters. To investigate the evolutionary relationship and genetic diversity of these isolates, phylogenetic analysis was done by neighbor joining (NJ) method implemented in MEGA X and branch support was computed with 1000 bootstrap replicates.

Population genetic parameters and checking for recombination

Estimate number of haplotypes (H), polymorphic sites (S), average pairwise nucleotide diversity (π), average number of nucleotide differences (K), and ratio of non-synonymous to synonymous nucleotide diversity (dN/dS) were obtained using DnaSP V.5.10.01 (Rozas, 2009). To identify individual codon positions evolving under negative (purifying) and positive selections, three different codon-based maximum-likelihood algorithms; single likelihood ancestor counting (SLAC), fixed effects likelihood (FEL), internal fixed effects likelihood (IFEL), fast and unconstrained bayesian approximation (FUBAR) within the HyPhy software package (Pond *et al.*, 2005) were used with significance level set at P-value < 0.1.

Recombination detection program RDP4 (Martin *et al.*, 2015) was applied to detect and analyze recombination signal in the CP gene sequences of ACLSV using several methods (Chimaera, Geneconv, MaxChi, RDP, Siscan,

LARD and 3Seq), with default setting, and a P-value threshold of 0.05.

Results

Detection of ACLSV by RT-PCR and sequencing

RT-PCR assays were performed to detect ACLSV in 174 samples. The results showed that 17 samples (9.7%) of both provinces were infected by ACLSV (Fig. 1). Eight isolates chosen from positive samples based on the geographic regions and symptoms were cloned and sequenced. Sequences were submitted to GenBank (accession numbers MK354022-MK354029; Table 1). Alleight ACLSV-CP sequences contained 582 nucleotides (nt) with 193 deduced amino acid residues and had a TAG at the end as stop codon. Identity of ACLSV-CP isolates ranged from 91 to 99.3% at the nucleotide level and 96.8-99.5% at the amino acid level.




Figure 1 Electrophoresis of PCR products amplified by ACLSV-MF and ACLSV-MR primers. M: Marker (Gene Ruler™ DNA Ladder Mix 1Kb), 1-3: Coat protein gene of ACLSV amplified from apple samples. 4: Negative control. The picture of leaf sample infected by ACLSV (Line 3) is shown on the right.

Table 1 Data on full coat protein gene sequences of Iranian isolates of ACLSV characterized in this study.

Collection site	Virus species	Isolate name	Host	Symptoms ³	Accession numbers
Kamyaran ¹	ACLSV	KSS	Apple	LD, Y	MK354022
Khoye ²	ACLSV	X2	Apple	Y	MK354023
Khoye ²	ACLSV	X	Apple	Y	MK354024
Divandareh ¹	ACLSV	D4	Apple	LD, Y	MK354025
Sanandaj ¹	ACLSV	SSa	Apple	LD, M	MK354026
Saqquez ¹	ACLSV	Sq	Apple	M	MK354027
Kamyaran ¹	ACLSV	KS	Apple	LD, Mo	MK354028
Marivan ¹	ACLSV	MS	Apple	NS	MK354029

¹ Kurdistan Province, Iran, ² Western Azarbaijan, Iran.

³ LD; leaf deformation, Y; yellowing, M; mosaic, mo; mottle, NS; no symptoms, BLM; burnt leaf margin, VC; vein clearing, LC; leaf curl. These symptoms may not necessarily refer to ACLSV and might be caused by other agents.

Phylogenetic analysis

Phylogenetic analysis based on the full-length CP sequences of the eight new ACLSV apple isolates in this study and 40 analogues of apple isolates from Brazil, China, Japan, Germany, India, Latvia, and Lithuania were divided into three major phylogroups; B6 (Yaegashi *et al.*, 2007), SHZ (Chen *et al.*, 2014), and P205 (Yaegashi *et al.*, 2007) (Fig. 2). Phylogroup I (B6 type) with the CP amino acid sequence of Ser⁴⁰-Tyr⁷⁵-Glu⁷⁹ was clustered into two subgroups; subgroup I was composed of the 22 isolates, including the all eight isolates described in this study and subgroup II contained 3 isolates from Brazil with one difference in amino acid at position 32. Phylogroup II (P205 type) with the CP amino acid sequence of Ala⁴⁰-Phe⁷⁵-Glu⁷⁹ was clustered into two subgroups; each subgroup included 6 isolates with two differences in amino acids at positions 72 and 97. Phylogroup III (SHZ type) included 11 isolates mainly from China containing the CP amino acid sequences of Ser⁴⁰-Tyr⁷⁵-Ser⁷⁹ with one exception in the isolate GZL13 with alanine (Ala) at position 40.

Coat protein variability of the Iranian ACLSV sequences

The genetic diversity of ACLSV isolates was studied by sequencing and aligning full-length CP sequences of cDNA clones in west and northwest Iranian isolates. The haplotype diversity for the new Iranian isolates of ACLSV was 1.000, whereas nucleotide diversity was 0.065 ± 0.007 . Nucleotide diversity value was lower than 0.1, indicating a low genetic diversity within Iranian ACLSV isolates. dN/dS ratio was lower than 1 (0.037), indicating that CP gene of Iranian ACLSV isolates is under negative selection.

The ACLSV-CP showed the greatest nucleotide diversity at position 351, while showing lower nucleotide diversity close to its 5'-terminal region (Fig. 3). We also found a decrease in the π value from position 494 towards the 3'-terminal region. Accordingly, lowest variability of amino acids was detected in the C and N-terminal regions of the CP (amino acids 2 to 58 and 127 to 193, respectively) with two amino acid changes in positions 40 and 170, while the middle region was substantially more variable (nine amino acid variations).




Figure 2 Neighbor-Joining phylogenetic tree constructed from the nucleotide alignment of coat protein sequences (582 nt) from the *Apple chlorotic leaf spot virus* isolates of this study (8 isolates) and 40 isolates retrieved from GenBank. Numbers over the branches indicate the bootstrap support. Amino acid abbreviations: A, alanine; S, serine; Y, tyrosine; E, glutamate; V, valine; F, phenylalanine; I, isoleucine.




Figure 3 Trend of polymorphism along the coat protein gene sequences in the Iranian ACLSV population, comprising of the new (this study) and published Iranian isolates (Abtahi et al., 2019). The curves were generated by sliding windows with 50 and 10 as the window and step sizes, respectively.

[DOR: 20.1001.1.22519041.2019.8.4.11.7] [Downloaded from jcp.modares.ac.ir on 2025-04-11]

Identifying individual codon positions evolving under negative and positive selections

Most of the codons in CP gene sequences of ACLSV isolates were under negative selection ($d_N/d_S < 1$) except for one codon, which was significantly under positive selection at positions 92, detected by SLAC and FEL methods (Table 2). In comparison with positive selection site, the sites under negative selection sites were very high, indicating strong negative selection was accrued on the CP gene.

Table 2 Analysis of selection for the CP of ACLSV Iranian isolates using the in Datamonkey server.

Methods	ACLSV	
	PS	NS
SLAC ¹	1 (92)	116
FEL ²	1 (92)	120
IFEL ³	0	112
FUBAR ⁴	0	140

PS: sites under positive selection, NS: sites under negative selection.

Codon-based maximum-likelihood algorithms.

¹ Single likelihood ancestor counting (SLAC).

² Fixed effects likelihood (FEL).

³ Internal Fixed effects likelihood (IFEL).

⁴ Fast, Unconstrained Bayesian Approximation (FUBAR).

Discussion

In this study, the infection, genetic variability and phylogenetic relationship of ACLSV from apple in Kurdistan and western Azarbaijan Provinces were studied. The infection level with ACLSV in apple trees in these two provinces was low, while a high infection level with ACLSV in apple cultivars and rootstocks was reported in different countries; up to 86% in Moldova (Verderevskaia and Marinescu, 1985), 73.8% in the Lithuania (Mazeikiene *et al.*, 2018), 70% in the Czech Republic (Grimova' *et al.*, 2016), 66% in Greece (Mathioudakis *et al.*, 2010) and 60% in the USA (Waterworth, 1993).

In phylogenetic analysis, ACLSV isolates from apple divided into three known phylogroups (B6, P205 and SHZ) by combinations of the three amino acids at positions: 40 (serine or alanine), 75 (tyrosine or

phenylalanine), and 79 (glutamate or serine) (Chen *et al.*, 2014) and the new Iranian ACLSV isolates were clustered in the B6 type (Fig. 2). We found that B6 and P205 phylogroups divided into two subgroups, I and II (Fig. 2). Two subgroups in the B6 and P205 had one and two differences in amino acid alignment, respectively. Subgroup I in B6 had alanine instead of valine at position 32 and subgroup I in p205 had valine and serine instead of isoleucine and asparagine at positions 72 and 97, respectively. Although the CP region in ACLSV isolates is relatively conserved, co-variations has been reported among phylogroups (Yaegashi *et al.*, 2007; Chen *et al.*, 2014; Niu *et al.*, 2012). Based on the combinations in the five amino acids at positions 40, 59, 75, 130, and 184, ACLSV isolates from Japan were clustered into the two major phylogroups B6 and P205 (Yaegashi *et al.*, 2007). Similarly, ACLSV isolates from apple samples worldwide have been tentatively classified into the three types among a new cluster (SHZ) of China based on the amino acids sequences at three positions 40, 75, 79 (Chen *et al.*, 2014). In the same way, ACLSV isolates from peach samples have been tentatively clustered into Z1 and Ta Tao 5 types (Niu *et al.*, 2012). To understand the role of these amino acid variations in pathogenicity and life cycle needs further data from more isolates, mutation study and ACLSV infection clones of these isolates. The long terminal branches of the isolates belonging to subgroup B of the B6 type (Fig. 1) may suggest that they have accumulated mutations over a long time.

To better characterize the genetic variability of the different Iranian isolates of ACLSV analyzed in this study and those published by Abtahi *et al.*, (2019), the descriptors of variability were determined and compared with the isolates from other countries (Nickel *et al.*, 2018; Table 2). Lower genetic variability in ACLSV-CP (0.0952 ± 0.006) was observed in the Iranian isolates than the isolates from other countries except for Brazilian population (0.0769 ± 0.005). The possible explanation for this result could be that the Iranian isolates originated from

one host (apple) (Table 1; Abtahi *et al.*, 2019) and therefore host-adaptation as a means of the heterogeneous nature of virus populations (Garcia-Arenal *et al.*, 2003) had not occurred.

The greatest nucleotide variation in Iranian isolates was found at the middle portion of ACLSV-CP (Fig. 3) similar to that reported from China (Chen *et al.*, 2014), Japan (Nickel *et al.*, 2018) and India (Rana *et al.*, 2010). In contrast, the Brazilian isolates showed conservation at the middle of CP (Nickel *et al.*, 2018). However, in the Iranian isolates the lowest nucleotide variation was observed in close to its 5'-terminal region similar to the Chinese, Brazilian, Japanese, and Indian isolates (Rana *et al.*, 2010; Nickel *et al.*, 2018).

The low genetic diversity between the Iranian ACLSV-CP sequences suggests that negative selection restricted the number of molecular variants and that virus populations are stable. Further analysis showed that the d_N/d_S value for the ACLSV Iranian and worldwide (among our isolates) isolates were 0.047 and 0.066, respectively. This result was also confirmed by evaluation of selective constraints across codon sites by comparing rates of synonymous to non-synonymous substitutions. One positively-selected codon in the ACLSV-CP was detected. In contrast, 112-140 negatively selected sites were found in ACLSV-CP by four methods (Table 2). These data suggest that strong negative selection is acting on the preservation of the ACLSV genome. Due to multiple functions of CP in viruses including genome encapsidation and protection, transmission between plants, host and/or vector interactions, cell to cell movement, suppression of gene silencing, negative selection pressures play a critical role in genome stability. Chare and Holmes (2004) analyzed selection pressures in the capsid genes of plant RNA viruses and found that vector borne viruses are subjected to a greater selection than non-vector-borne viruses. The d_N/d_S value for the 10 non-vector-borne viruses was > 0.093 whereas, for ACLSV-CP worldwide isolates was 0.066 (Nickel *et al.*, 2018) suggesting that the value is not absolute and

depends on the virus species and virus-host interaction.

In conclusion, the present analysis thus confirms the existence of a new cluster (BR) in phylogeny among ACLSV-CP isolates and a new amino acid sequence combination was suggested for the phylogroup differentiation in this virus. This study also confirms that negative selections are likely the potential drivers of the dynamics of ACLSV molecular evolution.

References

- Abtahi, F., Shams-Bakhsh, M., Safaie, N., Azizi, A., Autonell, C. R. and Ratti, C. 2019. Incidence and genetic diversity of apple chlorotic leaf spot virus in Iran. *Journal of Plant Pathology*, 101: 513-519.
- Adams, M. J., Lefkowitz, E. J. and King, A. M. Q. 2016. Ratification vote on taxonomic proposals to the International Committee on Taxonomy of Viruses. *Archives of Virology*, 161: 2921-2949.
- Al Rwahnih, M., Turturo, C., Minafra, A., Saldarelli, P., Myrta, A., Palla's, V. and Savino, V. 2004. Molecular variability of *Apple chlorotic leaf spot virus* in different hosts and geographical regions. *Journal of Plant Pathology*, 86: 117-122.
- Alemzadeh, E., Katsiani, A. T., Efthimiou, K. and Katis, N. I. 2016. Occurrence of apple chlorotic leaf spot virus in apple and quince in southern Iran. *Journal of Plant Pathology*, 98: 177.
- Cembali, T., Folwell, R. J., Wandschneider, P., Eastwell, K. C. and Howell, W. 2003. Economic implications of a virus prevention program in deciduous tree fruits in the US. *Crop Protection*, 22: 1149-1156.
- Chare, E. R. and Holmes, E. C. 2004. Selection pressures in the capsid genes of plant RNA viruses reflect mode of transmission. *Journal of General Virology*, 85: 3149-3157.
- Chen, S., Zhou, Y., Ye, T., Hao, L., Guo, L., Fan, Z., Li, S. and Zhou, T. 2014. Genetic variation analysis of apple chlorotic leaf spot virus coat protein reveals a new phylogenetic

- type and two recombinants in China. Archives of Virology, 159: 1431-1438.
- Chung, C. T., Niemela, S. N. and Muller, R. H. 1989. One step preparation of competent *Escherichia coli*: transformation and storage of bacterial cells in the same solution. Proceedings of the National Academy of Sciences of the United States of America, 86: 2172-2175.
- Derakhshian, M., Massah, A. and Bahar, M. 2010. Molecular identification of *Apple chlorotic leaf spot virus* in Isfahan and Western Azarbaijan provinces. 19th Iranian Plant Protection Congress, Tehran, Iran, 683.
- FAOSTAT. 2016. FAOSTAT Database. Food and Agriculture Organization of the United Nations.
- Farmahini, M., Pourrahim, R., Elahinia, A., Rouhibakhsh, A. and Farzadfar, S. 2014. An Investigation of *Apple chlorotic leaf spot* (ACLSV) and Tomato Ring Spot (ToRSV) in Some Iranian Pear Gardens. Iranian Journal of Virology, 8: 25-32.
- Foissac, X., Savalle-Dumas, L., Gentit, P., Dulucq, M. J. and Candresse, T. 2000. Polyvalent detection of fruit tree Tricho, Capillo and Faveaviruses by nested RT-PCR using degenerated and inosine containing primers (PDO RT-PCR). Acta Horticulturae, 357: 52-59.
- Garcia-Arenal, F., Fraile, A. and Malpica, J. M. 2003. Variation and evolution of plant virus populations. International Microbiology, 6: 225-232.
- Grimova, L., Winkowska, L., Zika, L. and Rysanek, P. 2016. Distribution of viruses in old commercial and abandoned orchards and wild apple trees. Journal of Plant Pathology, 98: 549-554.
- Keshavarz, T. and Shams-Bakhsh, M. 2015. Incidence and distribution of *Apple chlorotic leaf spot virus* in the main fruit growing areas of Iran. Archives of Phytopathology and Plant Protection, 48: 306-312.
- Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Molecular Biology and Evolution, 35: 1547-1549.
- Martin, D. P., Murrell, B., Golden, M., Khoosal, A. and Muhire, B. 2015. RDP4: Detection and analysis of recombination patterns in virus genomes. Virus Evolution, 1: vev003.
- Mathioudakis, M. M., Maliogka, V. I., Katsiani, A. T. and Katis, N. I. 2010. Incidence and molecular variability of apple stem pitting and apple chlorotic leaf spot viruses in apple and pear orchards in Greece. Journal of Plant Pathology, 92: 139-147.
- Mazeikiene, I., Siksnianiene, J. B., Gelvonauskiene, D., Bendokas, V. and Stanys, V. 2018. Prevalence and molecular variability of *Apple chlorotic leaf spot virus* capsid protein genes in Lithuania. Journal of Plant Disease and Protection, 125: 389-396.
- Naderpour, M., Shahbazi, R. and Sadeghi, L. 2016. Detection of some pome and stone fruit viruses in pome fruit orchards of West-Azarbaijan and Khorasan-e-Razavi provinces. Proceeding of 22th Iranian Plant protection congress, 27-30 August, University of Tehran, Karaj, Iran.
- Nemchinov, L., Hadidi, A., Foster, J. A., Candresse, T. and Verderevskya, T. 1995. Sensitive detection of *Apple chlorotic leaf spot virus* from infected apple or peach tissue using RT-PCR, IC-RT-PCR, or multiplex IC-RT-PCR. Acta Horticulturae, 386: 51-62.
- Németh, M. 1986. Virus, mycoplasma and rickettsia diseases of fruit trees. Martinus Nijhoff Publishers. Dordrecht. The Netherlands.
- Nickel, O., Silva, F. N., Fajardo, T. V. M. and Gorayeb, E. S. 2018. Characterization and genetic variability of coat protein genes of *Apple chlorotic leaf spot virus* isolates from southern Brazil. Tropical Plant Pathology, 43: 109-116.
- Niu, F. Q., Pan, S., Wu, Z. J., Jiang, D. M. and Li, S. F. 2012. Complete nucleotide sequences of the genomes of two isolates of apple chlorotic leaf spot virus from peach (*Prunus persica*) in China. Archives of Virology, 157: 783-786.

- Pond, S. L. K., Frost, S. D. W. and Muse, S. V. 2005. HyPhy: hypothesis testing using phylogenies. *Bioinformatics*, 21: 676-679.
- Rana, T., Chandel, V., Kumar, Y., Ram, R., Hallan, V. and Zaidi, A. A. 2010. Molecular variability analyses of *Apple chlorotic leaf spot virus* capsid protein. *Journal of Biosciences*, 35: 605-615.
- Rozas, J. 2009. DNA Sequence Polymorphism Analysis using DnaSP. In: Posada D (editor), *Bioinformatics for DNA sequence analysis; methods in molecular biology series*. Humana Press, NJ, USA, pp. 337-350.
- Sato, K., Yoshikawa, N., Takahashi, T. 1993. Complete nucleotide sequence of the genome of an apple isolate of *Apple chlorotic leaf spot virus*. *Journal of General Virology*, 74: 1927-1931.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G. 1997. The Clustal_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, 25: 4876-4882.
- Verderevskaia, T. D. and Marinescu, V. G. 1985. Virus and mycoplasma diseases of fruit cultures and grapevine. Kishinev, Moldova, pp. 117-127.
- Waterworth, H. 1993. Processing foreign plant germplasm at the national plant germplasm quarantine center. *Plant Disease*, 76: 854-860.
- Watpade, S., Raigond, B., Thakur, P. D., Handa, A., Pramanick, K. K., Sharma, Y. P. and Tomar, M. 2012. Molecular detection of latent *Apple chlorotic leaf spot virus* in Elite mother plants of apple. *Archives of Virology*, 23: 359-363.
- Wu, Y. Q., Zhang, D. M., Chen, S. Y., Wang, X. F. and Wang, W. H. 1998. Comparison of three ELISA methods for the detection of *Apple chlorotic leaf spot virus* and *Apple stem grooving virus*. *Acta Horticulturae*, 472: 55-60.
- Yaegashi, H., Isogai, M., Tajima, H., Sano, T. and Yoshikawa, N. 2007. Combinations of two amino acids (Ala⁴⁰ and Phe⁷⁵ or Ser⁴⁰ and Tyr⁷⁵) in the coat protein of apple chlorotic leaf spot virus are crucial for infectivity. *Journal of General Virology*, 88: 2611-2618.
- Yoshikawa, N. and Takahashi, T. 1988. Properties of RNAs and proteins of Apple stem grooving and Apple chlorotic leaf spot viruses. *Journal of General Virology*, 69: 241-245.
- Yoshikawa, N., Sasaki, E., Kato, M. and Takahashi, T. 1992. The nucleotide sequence of apple stem grooving capillovirus genome. *Virology*, 19: 98-105.

تعیین مشخصات و تنوع ژنتیکی ژن پروتئین پوششی جدایه‌های ویروس لکه‌برگی کلروتیک سیب جدا شده از سیب باغات استان‌های آذربایجان غربی و کردستان

کمال شجاعی، مهدی آذریار و محمد حاجی‌زاده*

گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه کردستان، سنندج، ایران.

پست الکترونیکی نویسندگان مسئول مکاتبه: m.hajizadeh@uok.ac.ir

دریافت: ۱۳ مرداد ۱۳۹۸؛ پذیرش: ۵ آبان ۱۳۹۸

چکیده: در این پژوهش، ۱۷۴ نمونه برگی سیب از مناطق عمده کاشت درختان سیب در غرب (استان کردستان) و شمال غرب (استان آذربایجان غربی) برای آلودگی به ویروس لکه‌برگی کلروتیک سیب مورد بررسی قرار گرفتند. نتایج RT-PCR حاکی از آلودگی ۱۷ نمونه (۹/۷ درصد از نمونه‌ها) به ACLSV از هر دو استان بود. سپس، بررسی تنوع ژنتیکی و آنالیز تبارزایی این ویروس با تعیین توالی ژن کامل پروتئین پوششی از هشت جدایه ویروس در این پژوهش و جدایه‌های تعیین توالی شده از ایران و موجود در بانک ژن انجام شد. در بررسی تبارزایی، جدایه‌های ACLSV به سه گروه شناخته شده B6، P205 و SHZ براساس ترکیب آمینواسیدی در موقعیت‌های ۴۰، ۷۵ و ۷۹ تفکیک شدند. جدایه‌های این پژوهش با ترکیب آمینواسیدی سرین^{۴۰}، تیروزین^{۷۵} و اسید گلوتامیک^{۷۹} در گروه تبارزایی شناخته شده B6 قرار گرفتند. میزان تنوع ژنتیکی جدایه‌های ایرانی حاکی از تنوع ژنتیکی کم این ویروس در مقایسه با سایر کشورها به جز جمعیت برزیلی این ویروس بود. اغلب کدون‌های پروتئین پوششی تحت گزینش منفی بودند به جز کدون ۹۲ که تحت فشار گزینش مثبت بوده است. این پژوهش نشان داد که فشار انتخاب منفی بیش‌ترین تأثیر را در تکامل ژن این ویروس داشته است.

واژگان کلیدی: سیب، فشار انتخاب منفی، آنالیز تبارزایی، ACLSV