

Research Article

Bioactivity of medicinal plant extracts as toxicants and enzyme inhibitors against insect pests of stored commodities

Kazam Ali^{1*}, Muhammad Sagheer², Mansoor ul Hasan², Abdul Rashid³ and Muhammad Shahid⁴

1. Centre for Agriculture and Bioscience International, Central and West Asia, Rawalpindi, Pakistan.

2. Department of Entomology, University of Agriculture, Faisalabad, Pakistan.

3. Department of Plant Pathology, University of Agriculture, Faisalabad, Pakistan.

4. Department of Bio-Chemistry, University of Agriculture, Faisalabad, Pakistan.

Abstract: The present research was performed to evaluate the bioactivity of *Citrullus colocynthis* (L.) and *Melia azedarach* L. extracts against three major stored grain insect pests including *Tribolium castaneum* (Herbst), *Trogoderma granarium* Everts, and *Sitophilus granaries* (L.). Toxicity and enzyme inhibition activity of acetylcholinesterase (AChE), α -carboxylesterase (α -CE), β -carboxylesterase (β -CE), acid phosphatases (ACP) and alkaline phosphatases (ALP) in three insect species induced by both plant extracts were evaluated at four different dose rates viz., 5, 10, 15 and 20%. Results showed maximum mortality (34.29%) in *S. granarius* with *M. azedarach* at maximum interaction of time and dilution level. In *T. castaneum* and *T. granarium* maximum recorded values for mortality were 30.87% and 18.95%, respectively, with extract of *M. azedarach*. Plant extract of *C. colocynthis* reported a maximum mortality of 21.92%, 19.18% and 16.89% in *T. castaneum*, *S. granarius* and *T. granarium*, respectively. Findings proved that both plant extracts had decent lethal impacts on tested insect species. Exposure of studied insects to plants extracts also resulted in significant inhibition of AChE, α -CE, β -CE, ACP and ALP. All tested enzymes in three insects were maximally inhibited by plant extract of *M. azedarach* except α -CE which was slightly more inhibited in *S. granarius* and ACP which was highly inhibited in *T. granarium* and *S. granarius*, by plant extract of *C. colocynthis*. Outcomes exhibit that plant based extract of *M. azedarach* is more pronounced in stored grain insect pests and propose the capability of using these plant extracts for safety of stored commodities as a safe substitute for insecticides.

Keywords: stored product pests, enzyme inhibition, lethal effects, toxicity

Introduction

Stored grains and their products are at great risk to the infestation by insect pests (Ukeh *et al.*, 2012). Stored food commodities are infested by hundreds of hexapods and other arthropods

among which about 600 species belongs to order Coleoptera (Rajendran and Sriranjini, 2008). In Pakistan, Red rust flour beetle *Tribolium castaneum*, Khapra beetle *Trogoderma granarium*, Angumois grain moth *Sitotroga cerealella*, Lesser grain borer *Rhyzopertha dominica*, Rice weevil *Sitophilus oryzae* and Grain weevil *Sitophilus granarius* are reported to be the most injurious insect pests of stored commodities (Iqbal *et al.*, 1992). *T. castaneum* is a secondary pest of stored

Handling Editor: Saeid Moharramipour

*Corresponding author: k.ali@cabi.org

Received: 02 June 2020, Accepted: 10 October 2020

Published online: 27 October 2020

commodities, both larvae and adults feed on grain dust and broken grain and spend entire life cycle outside the grain kernels (Karunakaran *et al.*, 2004). In severe infestation, the flour turns grayish and has a pungent smell (benzoquinone), disagreeable odour making it unfit for human consumption. This insect causes substantial loss in storage because of its high reproductive potential (Prakash *et al.*, 1987). Primary stored grain pest *T. granarium* has been nominated as one of the 100 worst invasive species worldwide. It is a serious pest of stored products under hot, dry conditions (Lowe *et al.*, 2000). Generally, young larvae of *T. granarium* feed on damaged seed, while older larvae feed on whole grains. Larvae attack the embryo point or a weak place in the pericarp of grain or seed. The khapra beetle can cause significant weight loss (weight loss between 5-30%, extreme cases of 70%) when left undisturbed in stored grain. Granary weevil (*S. granarius*) is another important stored grain pest which causes significant losses during storage (White and Leesch, 1996). The weevils bore into the kernels, insert their eggs within the endosperm of the grain and develop inside whole grain kernels as small, white, wrinkled, grub-like larvae (CABI, 2007), meanwhile their damage remains un-noticed until the weevils emerge from the seed with an exit hole (Niewiada *et al.*, 2005).

Recently, a great attention has been paid to the use of *Citrullus colocynthis* and *Melia azedarach* extract as natural insecticides. The biological activity of these plants has been investigated against many insect pests (Soam *et al.*, 2013). These plants are known to have a range of compounds, which show insecticidal, antibacterial, larvicidal, deterrent, antifeedant, growth regulating, antifertility, metamorphosis and reproduction disturbance effects (Pravin *et al.*, 2013; Soam *et al.*, 2013). Aqueous and methanolic extracts of plant, *C. colocynthis* demonstrated high antimicrobial activity against some bacteria and fungi. *C. colocynthis* can be used medically as an abortifacient, cathartic, purgative and vermifuge, as well as for the treatment of fever, cancer, amenorrhea,

jaundice, leukemia, rheumatism, tumour, and as an insect repellent (Soam *et al.*, 2013). El-Naggar *et al.* (1989) noted the impact of colocynthin and hydrated colocynthin isolated from *C. citrullus* against on seven insect species and reported effectual outcomes. Chemicals isolated from *Melia azedarach* L. species Meliaceae family have gained a particular attention from applied entomologists because of their excellent properties as insect control agents (Luo *et al.*, 1995). *M. azedarach* is native to Iran, India, and China (Hong and Ellis, 1998). The plant has become the object of studies to evaluate properties from different plant structures, in particular insecticidal, antiviral, antioxidant, bactericide, and antiparasitic activities (Ahmed *et al.*, 2008). The insecticidal activity of *M. azedarach* is found in leaves, fruits, and seeds, and is due to a group of biologically active triterpenoids they have antifeeding effects (Isman, 2006). Generally, extracts from green fruits and leaves have been those most efficacious because of their antifeedant effect, mainly on beetles and lepidopterans (Carpinella *et al.*, 2003, 2005; Nathan and Kim, 2005; Defago *et al.*, 2006).

Enzymes are main group of proteins and biomolecules which enhance metabolic activities in the organisms. Esterases and phosphatases are important components for the proper functioning of several important physiological processes in insects (Lassiter *et al.*, 1995; Shanmugavelu *et al.*, 2000). A little change in the levels of enzymes affects the metabolic process in organism (Roy, 2002). In insects, resistance against insecticides arises by the change of metabolic enzymes or increased detoxification (Parakrama Karunaratne, 1998). Due to the change in the target site in the insect, insecticides cannot bind to that site and insect behaves normally (Damayanthi and Karunaratne, 2005). By checking the change in metabolism of insects, the activity of the enzymes released in their plasma due to cell disturbance can be calculated (Coppo *et al.*, 2002). So inhibition of enzymes is a reliable method to assess the pressure on the insects by pollutants. Esterases and phosphatases are considered as reliable bio-

markers for assessing the toxic effects of numerous insecticidal compounds on the physiology of targeted insects (Srinivas *et al.*, 2004). This study was designed to investigate the toxic and enzyme inhibitory effects of *C. colocynthis* and *M. azedarach* extracts against *T. castaneum*, *T. granarium* and *S. granarius*.

Materials and Methods

Collection and rearing of insects

The mixed populations of stored grain insect pests were collected from grain market and flour grinding mills, and were brought into the Grain Research Training and Storage Management Cell, Department of Entomology, University of Agriculture Faisalabad, Pakistan. In the laboratory, the target insects, red flour beetle *T. castaneum*, Khapra beetle *T. granarium* and Granary weevil *S. granarius* were separated and reared for homogeneous population.

Populations of the three coleopteran species were cultured in sterilized plastic jars (1.0 kg capacity) and 50 adults of each species were released separately into their favorite diet, these adults were allowed for copulation and were removed after 5 days. Wheat flour was used for rearing of *T. castaneum* (Sagheer *et al.*, 2014), while wheat grains were used as culture medium for *T. granarium* and *S. granarius*. After removal of adults, sieved flour and grains containing eggs laid by these insects were again put into the jars and placed in incubators for getting same age population. Rearing jars of *T. castaneum* and *T. granarium* were placed in incubators (SANYO) at 30 ± 2 °C and $65 \pm 5\%$ R.H., while *S. granarius* was cultured at 27 ± 2 °C and $70 \pm 5\%$ R. H.

Collection of plant materials

Plant materials, such as fruits of *C. colocynthis* (Tuma) were collected from district Layyah and leaves of *M. azedarach* (Darek) were collected from the fields of University of Agriculture, Faisalabad (UAF).

Preparation of plant extracts

The fruits of *C. colocynthis* and leaves of *M.*

azedarach were washed with sterilized water before placing in shade for drying. Once the plant materials were dried they were ground in electrical grinder and were brought into fine powder. The plant extracts were obtained by adding 50.0g plant powder in 100 ml acetone and solution was set on a Rotary Shaker (IRMECO, OS-10) (Sagheer *et al.*, 2014) at 220 rpm for a period of 24 hours. After 24 hours rotation the solution obtained was filtered and placed on rotary evaporator to remove the extra solvent (acetone). Thus, the plant extracts obtained after evaporation were considered as stock solution and stored at 4.0 °C. Four dilution levels (5.0, 10.0, 15.0 and 20.0%) of each plant extract were prepared from the stock solutions, using acetone as solvent.

Toxic effect of plant extracts

Different dilutions (5.0, 10.0, 15.0, 20.0%) of acetone extract of *C. colocynthis* and *M. azedarach* were applied on favorite diet of each tested insect species i.e., wheat flour for *T. castaneum*, and wheat grains for *T. granarium* and *S. granarius*. After evaporation of acetone the dried weighted wheat flour (40.0 g) and wheat grains (40.0 g) were put into treatment jars separately. Fifty adults of *T. castaneum* and *S. granarius*, fifty second instar larvae of *T. granarium* were released separately into experimental jars containing treated wheat and flour. Each treatment was replicated three times and experimental jars were placed in SANYO incubator under optimum conditions as discussed above. Observations for lethal impact of *C. colocynthis* and *M. azedarach* were made after 2, 4, 6, 8 and 10 days of experiment. The alive larvae of *T. granarium* and alive adults of *T. castaneum* and *S. granarius* were stored in phosphate buffer solution to evaluate the enzymes activity.

Mortality was calculated using Abbott's (1925) formula;

$$\text{Corrected Mortality (\%)} = \frac{(\text{Mo} - \text{Mc})}{100 - \text{Mc}} \times 100$$

Where,

Mo = Mortality observed in treatments

Mc = Mortality observed in control

Inhibition of esterases and phosphatases in survivors of toxicity assay

Preparation of whole body homogenate

The survived specimen (adults of *T. castaneum* and *S. granarius*, larvae of *T. granarium*) in toxicity experiment which were stored in buffer solution were rinsed with clean water, and the adhering water was entirely removed from insect body by blotting with tissue paper. In ice-cold sodium phosphate buffer (20 mM, pH 7.0), the survivors (larvae and adults) of test insects were homogenized separately using a teflon hand homogenizer for eventual estimation of esterases and phosphatases inhibition. For biochemical analyses, clear supernatants were used obtained by centrifuging the whole body homogenates at 8000 rpm for 20 minutes. Before using all the solutions for homogenization, glassware were stored at 4.0 °C and homogenates were kept on ice until used for different assays.

Determination of acetylcholinesterase (AChE)

Acetylcholinesterase (AChE) activity in the whole body homogenates of *T. castaneum*, *T. granarium* and *S. granarius* were measured spectrophotometrically according to Ellman *et al.* (1961) with slight modification of using acetylthiocholine iodide instead of acetylthiocholine chloride as substrate. 50 µl of acetylthiocholine iodide (2.6×10^{-3} M) as a substrate and 1 ml of sodium phosphate buffer (20 mM, pH 7.0) was added to 100 µl of enzyme solution taken from whole body homogenates and incubated at 25 °C for 5 minutes. Then 400 µl of freshly prepared 0.3% Fast blue B salt in 3.3% SDS (sodium dodecyl sulphate) was added to stop reaction. Sample was run through spectrophotometer and optical density was recorded at 405 nm.

Determination of α and β -carboxylesterase (α -CE & β -CE)

The activity of α - and β -carboxylesterase in the whole body homogenates of test insects was measured by the method of van Asperen (1962). The α -carboxylesterase activity was recorded

by using α -naphthylacetate as substrate. For this purpose 50 µl of α -naphthylacetate (250 µM) and 1 ml of sodium phosphate buffer (20 mM, pH-7.0) was added in 50 µl whole body homogenates. This solution was incubated at 30 °C for 20 minutes. 400 µl of freshly prepared 0.3% Fast blue B salt in 3.3% SDS, was added to stop the reaction. Sample was run on spectrophotometer and optical density was noted at 430 nm.

Same procedure was followed for β -carboxylesterase activity, except that β -naphthylacetate was used as substrate and optical density was measured at 590nm.

Determination of acid and alkaline phosphatases (ACP & ALP)

The level of acid phosphatases (ACP) and alkaline phosphatases (ALP) were calculated by following the Asakura (1978) method. P-nitrophenyl phosphate was used as substrate for the estimation of phosphatases. For acid phosphatase (ACP), 100 µl of 20mM p-nitrophenyl phosphate (substrate) and 450µl sodium acetate buffer (50 mM, pH 4.6) were added in 50 µl enzyme solution. The solution was incubated at 37 °C for 15 mins. Then, 100 µl of 0.5N NaOH was added to stop reaction. Optical density of sample was recorded at 405 nm.

Same procedure was followed for alkaline phosphatases (ALP) except that 450 µl Tris HCl (50 mM, pH-8) was used in place of sodium acetate buffer.

Enzyme inhibitions (%) of test enzymes were computed using the formula given by Wang *et al.*, (2014).

$$\text{Enzyme Inhibition (\%)} = \frac{(\text{OD}_b - \text{OD}_o)}{\text{OD}_b} \times 100$$

OD_b = Optical density of blank (control treatment)

OD_o = Optical density of treatments

Statistical analyses

Separate two-way factorial ANOVA were performed for both *C. colocynthis* and *M. azedarach* plant extracts against each tested insect species. The differences and alterations

in the levels of different enzymes in *Tribolium castaneum*, *Trogoderma granarium* and *Sitophilus granarius*, were computed by using mean difference Tukey-HSD test (Statistica-

8.1). The means sharing similar letters within column are statistically same in Table 1. The level of acceptance for significant difference was $p \leq 0.05$ in all cases.

Table 1 Comparison of mortality in three stored grains insect species by acetone extracts of *Citrullus colocynthis* and *Melia azedarach*.

Time (Days)	Conc. (%)	Mean Mortality \pm S.E (%)					
		<i>Citrullus colocynthis</i>			<i>Melia azedarach</i>		
		<i>T. castaneum</i>	<i>T. granarium</i>	<i>S. granarius</i>	<i>T. castaneum</i>	<i>T. granarium</i>	<i>S. granarius</i>
2	5	3.33 \pm 0.67 j	1.34 \pm 0.34 f	3.38 \pm 0.68 i	3.33 \pm 0.89 j	0.67 \pm 0.33 j	4.05 \pm 1.18 l
	10	6.00 \pm 0.67 hij	2.68 \pm 1.34 ef	4.73 \pm 1.17 hi	5.33 \pm 1.34 ij	2.01 \pm 1.34 ij	6.08 \pm 0.68 l
	15	8.00 \pm 1.15 fi	4.02 \pm 0.67 def	6.08 \pm 0.69 ghi	8.67 \pm 0.67 fj	2.68 \pm 0.67 hij	10.14 \pm 1.49 k
	20	8.00 \pm 0.35 fi	7.38 \pm 2.01 b-f	7.44 \pm 1.86 fi	12.00 \pm 2.01 d-g	4.69 \pm 2.01 fj	11.49 \pm 1.34 k
4	5	5.41 \pm 0.68 ij	2.68 \pm 0.67 ef	6.12 \pm 0.68 ghi	6.08 \pm 0.67 hij	3.35 \pm 0.67 g-j	7.24 \pm 1.17 kl
	10	8.11 \pm 0.68 fi	5.37 \pm 1.16 def	6.80 \pm 1.18 fi	6.76 \pm 1.16 g-j	5.37 \pm 1.16 e-j	12.93 \pm 1.67 jk
	15	8.79 \pm 0.59 fi	6.04 \pm 0.67 c-f	9.52 \pm 2.07 d-h	12.84 \pm 0.67 def	6.71 \pm 0.67 e-j	15.65 \pm 1.34 ij
	20	10.14 \pm 0.68 efg	9.39 \pm 2.32 a-e	10.20 \pm 1.17 d-g	15.54 \pm 2.32 d	9.39 \pm 2.32 c-g	19.05 \pm 1.98 hi
6	5	6.80 \pm 0.34 g-j	5.37 \pm 1.16 def	8.16 \pm 1.78 e-i	6.80 \pm 1.16 g-j	4.69 \pm 1.16 fj	15.65 \pm 2.34 ij
	10	9.52 \pm 1.34 e-h	6.71 \pm 0.67 b-f	8.84 \pm 1.51 d-h	11.56 \pm 0.67 d-h	7.38 \pm 0.67 e-i	21.09 \pm 2.11 gh
	15	12.93 \pm 0.68 de	9.39 \pm 2.32 a-e	11.56 \pm 1.18 c-f	16.33 \pm 2.32 d	10.74 \pm 2.32 b-f	24.49 \pm 1.67 fg
	20	15.65 \pm 1.35 cd	10.74 \pm 1.78 a-d	12.93 \pm 0.68 b-e	23.81 \pm 1.78 bc	14.09 \pm 1.78 bcd	28.57 \pm 1.39 de
8	5	10.27 \pm 1.37 efg	5.41 \pm 0.68 def	9.59 \pm 1.06 d-h	9.59 \pm 0.68 e-i	5.41 \pm 0.68 e-j	19.86 \pm 1.34 h
	10	11.64 \pm 0.36 ef	8.11 \pm 0.68 b-f	13.70 \pm 1.79 bcd	14.38 \pm 0.68 de	8.79 \pm 0.68 c-h	26.02 \pm 2.17 ef
	15	17.81 \pm 1.19 bc	11.49 \pm 1.79 a-d	15.75 \pm 1.88 abc	21.92 \pm 1.79 c	11.49 \pm 1.79 b-e	31.05 \pm 3.11 bcd
	20	19.18 \pm 1.68 abc	14.19 \pm 2.44 ab	17.12 \pm 1.45 ab	27.76 \pm 2.44 ab	16.89 \pm 2.44 ab	33.56 \pm 1.68 abc
10	5	13.01 \pm 1.34 de	7.44 \pm 0.68 b-f	13.01 \pm 0.67 b-e	11.64 \pm 0.68 d-g	8.11 \pm 0.68 d-i	24.65 \pm 1.58 f
	10	15.75 \pm 1.59 cd	8.79 \pm 1.17 b-f	16.44 \pm 2.05 abc	15.07 \pm 1.17 de	10.81 \pm 1.17 b-f	29.13 \pm 2.87 cd
	15	20.55 \pm 0.68 ab	13.52 \pm 0.68 abc	17.81 \pm 1.45 ab	24.65 \pm 0.68 bc	14.87 \pm 0.68 abc	33.93 \pm 1.62 ab
	20	21.92 \pm 1.03 a	16.89 \pm 2.34 a	19.18 \pm 1.86 a	30.87 \pm 2.34 a	18.95 \pm 2.34 a	34.29 \pm 2.67 a

Separate two ways factorial ANOVA's were performed for test insects against each acetone based plant extract. The means were computed using Tukey-HSD test, similar letters within treatments (column) are not statistically different ($p < 0.05$).

Results

Toxicity experiments were performed to evaluate the lethal effects of plant extracts of *C. colocynthis* and *M. azedarach* against *T. castaneum*, *T. granarium* and *S. granarius*. Inhibition of various enzymes, acetylcholinesterase (AChE), α -carboxylesterase (α -CE), β -carboxylesterase (β -CE), acid phosphatases (ACP) and alkaline phosphatases (ALP) were studied in the survivors of the toxicity experiment. The means comparison for mortality at different interactions of time interval (2, 4, 6, 8 and 10 days) and concentrations

(5, 10, 15 and 20%) induced by acetone extract of *C. colocynthis* and *M. azedarach* in *T. castaneum*, *T. granarium* and *S. granarius* are given in Table 1. Results indicated that *M. azedarach* forced a maximum 34.29% mortality in *S. granarius*, while maximum mortality evidenced in other two insect species *T. castaneum* and *T. granarium* were 30.87% and 18.95% respectively, which were also noted with *M. azedarach* at 20% dilution level after 10 days interval. The result reveals that plant extracts of *C. colocynthis* showed slightly less toxicity than *M. azedarach* except the three different tested organisms and recorded a

maximum mortality rate of 21.92% in *T. castaneum*, 16.89% in *T. granarium* and 19.18% in *S. granarius* at 20% concentration after 10 days of application. The impact of time and concentration proves directly proportional to mortality of tested insects. Initially *C. colocynthis* was established as stronger insecticide than *M. azedarach* against *T. granarium* as it reported 7.38% mortality which was only 4.69% with *M. azedarach* at 20% concentration after 2 days exposure, but as time went on *M. azedarach* became more lethal and showed higher values for mortality in the three insect species (Table 1).

Results evidenced that *M. azedarach* plant extract showed high inhibition of AChE (acetylcholinesterase) activity for all three test insects. Maximum AChE inhibition 37.14% was assessed in *S. granarius* at 20% concentration, while at the same dilution level 33.33% and 31.48% were the maximum inhibition values for *T. castaneum* and *T. granarium* by *M. azedarach*. Plant extract of *C. colocynthis* showed maximum inhibition (27.79%) of AChE activity in *T. castaneum* at 15% concentration, while at 20% concentration maximum inhibition (22.23% and 25.58%) of AChE activity were noted against *T. granarium* and *S. granarius*, respectively. Lower concentration (5%) showed

minimum inhibition effect for AChE activity as it noted 25%, 16.07%, 14.67% inhibition by *C. colocynthis* plant extract; while 24.16%, 25.77%, 25.92% inhibition of AChE activity was forced by *M. azedarach* in *T. castaneum*, *T. granarium* and *S. granarius*, respectively (Fig. 1).

The inhibition level of α -CE (α -carboxylesterase) activity significantly increased at various concentrations, and it reached the highest level at 20% concentration in three insects when exposed to the both plant extracts ($p < 0.05$). This enzyme inhibition level steadily decreased in *T. castaneum* from 10% concentration (17.06% inhibition) to 15% concentration (13.98% inhibition) with *C. colocynthis* extract while from 5% concentration (13.94% inhibition) to 10% concentration (10.33% inhibition) with *M. azedarach* plant extract. Maximum α -CE inhibition level in *T. castaneum*, *T. granarium* and *S. granarius* noted at highest concentration (20%) were 19.67%, 17.63% and 22.34% induced by *C. colocynthis*. *M. azedarach* force a maximum 24.98% and 22.36% inhibition of α -CE activity in *T. castaneum* and *T. granarium* at 20% concentration respectively, while in *S. granarius* highest (16.79%) inhibition of α -CE activity was noted at 15% concentration (Fig. 2).




Figure 1 Impact of *Citrullus colocynthis* and *Melia azedarach* extracts at different concentrations on acetylcholinesterase (AChE) activity in three stored grains insect species.




Figure 2 Impact of *Citrullus colocynthis* and *Melia azedarach* extracts at different concentrations on α -carboxylesterase (α -CE) activity in three stored grain insect species.

The exposure of the target insects to both plant extracts had great impact on the inhibition level of β -CE (β -carboxylesterase) activity. From lower (5%) to higher (20%) concentration the inhibition level of β -CE activity steadily increased from 9.14% to 17.72% in *T. castaneum*, 12.72% to 21.71% in *T. granarium* and 14.67% to 25.58% in *S. granarius* when exposed to plant extract of *C. colocynthis* ($p < 0.05$). *M. azedarach* extract showed significant effect on percent inhibition of β -CE (β -carboxylesterase) activity in *T. castaneum*, *T. granarium* and *S. granarius* ($p < 0.01$). At 20% concentration, 41.67%, 38.21% and 30.84% inhibition of β -CE activity was observed in *T. castaneum*, *T. granarium* and *S. granarius*, respectively. The effect of various concentrations of *M. azedarach* on inhibition of β -CE activity against *T. granarium* and *S. granarius* was in the following order: 20% > 15% > 10% > 5%. Minimum percent inhibition of β -CE activities noted in *T. castaneum*, *T. granarium* and *S. granarius*, were 20.83%, 30.49% and 18.47%, respectively, at 5% concentration (Fig. 3).

Inhibition of acid phosphatase (ACP)

activity slightly increased from 5 to 20% dilution level of *C. colocynthis* and *M. azedarach* in three coleopteran species (Fig. 4). Maximum ACP inhibition (38.57%) was assessed in *T. castaneum* with *M. azedarach* at 20% concentration, while these values were 15.38% and 13.41% for *T. granarium* and *S. granarius* at maximum dilution level of *M. azedarach* respectively. Exposure of the test insects to *C. colocynthis* treated diet significantly increased the inhibition of ACP activity ($p < 0.05$) with increase in dilution level and resulted in maximum of 29.92%, 18.65% and 15.21% ACP inhibition against *T. castaneum*, *T. granarium* and *S. granarius*, respectively at 20% concentration (Fig. 4).

Results illustrated that inhibition of alkaline phosphatase activity gradually increased from lower (5%) to higher (20%) concentration in three test insects exposed to the plant extracts ($p < 0.05$). At 20% dilution, this enzyme level was significantly inhibited ($p < 0.01$). Maximum ACP inhibition (29.44%) was noted in *T. granarium* with *M. azedarach* at 20% concentration, while extract of *C. colocynthis* forced 13.73% inhibition in *T. granarium*

which was the lowest inhibition value in three stored grain species at the highest dose rate (20%). Exposure to *C. colocynthis* extract (20%) resulted in 17.34% and 21.70% inhibition of ACP activity in *T. castaneum* and

S. granarius. Whereas *M. azedarach* extract affected slightly higher than *C. colocynthis* and evidenced 18.41% and 23.09% inhibition of ACP enzyme activity in *T. castaneum* and *S. granarius* respectively (Fig. 5).




Figure 3 Impact of *Citrullus colocynthis* and *Melia azedarach* extracts at different concentrations on β -carboxylesterase (β -CE) activity in three stored grains insect species.




Figure 4 Impact of *Citrullus colocynthis* and *Melia azedarach* extracts at different concentrations on acid phosphatases (ACP) activity in three stored grains insect species.




Figure 5 Impact of *Citrullus colocynthis* and *Melia azedarach* extracts at different concentrations on alkaline phosphatases (ALP) activity in three stored grains insect species.

Discussions

The findings for insect mortality evidenced that lethal impact of *C. colocynthis* and *M. azedarach* was directly related to exposure periods and concentrations. Benzi *et al.* (2009) reported similar results showing the toxic efficacy of extracts from some medicinal plants against mites and insects. Similarly, mortality of adults of *T. castaneum* increased with increase in concentration at maximum exposure period (Bibi *et al.*, 2008). The insecticidal activity of *M. azedarach* is found in leaves, fruits, seeds, and is due to a group of biologically active triterpenoids that have anti-feeding effects (Valladares *et al.*, 1997; Isman, 2006). *C. colocynthis* has a large range of insecticidal, antibacterial, larvicidal, deterrent, antifeedant, growth regulating and antifertility compounds (Pravin *et al.*, 2013; Soam *et al.*, 2013). Ali *et al.* (2017) verified same results under similar laboratory conditions that exposure of *T. castaneum*, *T. granarium* and *S. granarius* to plant extracts of *Azadirachta indica* and *Datura innoxia*, cause significant mortality in the test populations after 10 days

intervals at 20.0% concentrations. Our results testify, plant extract of *M. azedarach* at tested concentrations proved more valuable as they forced higher mortality. These outcomes are supported by Anwar *et al.* (2005) who checked the neem (*A. indica*) oil in a warehouse, against four important stored grain insect pests *Rhyzopertha dominica*, *S. granarius*, *T. castaneum* and *T. granarium* at various dilution levels (5%, 10%, 15% and 20%) in natural conditions at three time periods (30, 60, and 90 days). They observed increased mortality with the increase in dose rate of the spray material.

Both toxicant, *C. colocynthis* and *M. azedarach*, showed decent anti-enzymatic activities in tested insect species, *T. castaneum*, *T. granarium* and *S. granarius*. Different insecticides targeted the AChE (Abdelgaleil *et al.*, 2009; Kang *et al.*, 2013), carboxylesterases also verified as principal enzyme in many tissues of a number of insects (Park and Kamble, 1999). Based on our findings, AChE activity was inhibited by both plants (*M. azedarach* and *C. colocynthis*) extracts but at minimum concentration (5%) enzyme inhibition was also low and AChE inhibition

increases at lateral dilution level (20%). Enzyme inhibition results are supported by Wang and his co-workers (2014), as they noted the impact of *Citrus limonum*, *Litsea cubeba*, *Cinnamomum cassia* and *Allium sativum* against *Alphitobius diaperinus* (darkling beetle) pest of stored poultry feed. They found that essential oils of tested plants significantly inhibited the levels of AChE activity and *A. sativum* resulted in highest inhibition (> 80%) of AChE activity. Our results are in agreement with their findings as they verified that inhibition of AChE activity increased with exposure time. Kim *et al.* (2013) reported some compounds from plant oils of apiaceae family, as inhibitor of AChE activity against *S. oryzae*, some other scientists also noted same results in number of insect pests (Breuer *et al.*, 2003; Nathan *et al.*, 2008). Our findings also demonstrated that *C. colocynthis* and *M. azedarach* extracts caused significant inhibition of α -carboxylesterase (α -CE) and β -carboxylesterase (β -CE) activities in three insect species. Outcomes indicated maximum inhibition of α -CE and β -CE activity in *T. castaneum*, *T. granarium* and *S. granarius* at 20% concentration. Dose dependent reactions of α -CE and β -CE activities were also noted in the larvae of *Choristoneura rosaceana* exposed to *M. azedarach* oil (Smirle *et al.*, 1996). Mujeeb and Shakoori (2012) evidenced that Fury (synthetic pyrethroid) inhibits the carboxylesterase (CE) activity in all life stages of red flour beetle, *T. castaneum*. Koodalingam *et al.* (2011) proved that when the larvae of *Aedes aegypti* were released to extract of soapnut, *Sapindus emarginatus*, it significantly reduced the activities of AChE and β -CE, while no changes were observed in the level of α -CE activity.

Most of the physiological processes are phosphatases dependent as they play a vital role in the completion of their normal functions (Majerus *et al.*, 1999). Our results showed that higher inhibition (38.57%) of acid phosphatases (ACP) activity was noted in *S. granarius* exposed to 20% solution of *M. azedarach*, whereas 18.65% and 15.21% maximum

inhibition of ACP was checked in *T. granarium* and *T. castaneum* feed on *C. colocynthis* treated diet. In case of alkaline phosphatases (ALP) maximum inhibited values 29.44%, 23.09% and 18.41% were recorded for *T. granarium*, *S. granarius* and *T. castaneum* respectively, by plant extract of *M. azedarach*. Phosphatases are deemed as reliable tools to assess the toxic impacts of various chemicals on physiological status of insects (Srinivas *et al.*, 2004). These results are justified by Nathan *et al.* (2005) as they tested and concluded that the exposure of larvae of *Cnaphalocrocis medinalis* and *Spodoptera litura* to azadirachtin, resulted in significant inhibition of acid and alkaline phosphatases. Similarly acetone and ethanol based plant extracts of *A. indica* and *Datura innoxia* (Ali *et al.*, 2017), *Artemisia annua* (Shekari *et al.*, 2008), *Teucrium royleanum* (Ahmad *et al.*, 2007a), *Andrachne cordifolia* (Ahmad *et al.*, 2007b), *Cassia obtusifolia* (Kim *et al.*, 2007), *Gloriosa superba* (Khan *et al.*, 2007), *Paeonia emodi* (Khan *et al.*, 2005) and *Corydalis incise* (Kim, 2002) were shown to have significant impact on the inhibition of AChE, ALP, urease, lipoxxygenase, and amino transferase of various hexapods.

Conclusions

All the reported outcomes taken together indicate that both plant extracts *M. azedarach* and *C. colocynthis* release their anti-insecticidal activity by various modes of action as evidenced from mortality and numerous adverse changes examined in various important enzymes, including AChE, α -CE, β -CE, ACP and ALP of *T. castaneum*, *T. granarium* and *S. granarius*. Results revealed distinct differences in alterations of biochemical characteristics in the three stored grain insect species exposed to the botanical biocide tested. It also verified from the findings of the experiment that *T. castaneum* and *S. granarius* were more susceptible, while *T. granarium* was slightly tolerant to both toxicants. Conclusions suggest the use of these botanicals as a substitute for chemical insecticides for storage of wheat

grains, flour and their byproducts.

Conflict of interests

The Authors have no conflict of interest.

Author's contribution

Kazam Ali: Collected experimental material, performed the experiments, analyzed the data and wrote the article

Muhammad Sagheer, Mansoor ul Hasan and Abdul Rashid: Developed protocols for plants extracts and their lethal impacts and also supervised the studies.

Muhammad Shahid: Developed protocols and supervised the enzyme studies of insects.

Acknowledgments

We acknowledge the contribution of the anonymous reviewers for their comments and suggestions for the improvement of this manuscript. The authors are also thankful to the Higher Education Commission (HEC) of Pakistan for providing financial assistance as a PhD scholarship to the first author, the research article is extracted from PhD thesis.

References

- Abbott, W. S. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18: 265-267.
- Abdelgaleil, S. A. M., Mohamed, M. I. E., Badawy, M. E. I. and El-arami, S. A. A. 2009. Fumigant and contact toxicities of monoterpenes to *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst) and their inhibitory effects on acetylcholinesterase activity. *Journal of Chemical Ecology*, 35: 518-525.
- Ahmad, B., Mukarram, S. M., Khan, H., Hassan, M. and Shan, J. 2007a. Enzyme inhibition activities of *Teucrium royleanum*. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 22: 730-732.
- Ahmad, B., Shan, S. M., Bashir, S. and Shan, J. 2007b. Enzyme inhibition activities of *Andrachne cordifolia* Mull. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 22: 235-238.
- Ahmed, M., Ahmed, M., Thayyil, H., Zameeruddin, K. and Ibrahim, M. 2008. Antioxidative activity of *Melia azedarach* Linn leaf extract. *Iranian Journal of Pharmacology and Therapeutics*, 7: 31-34.
- Ali, K., Sagheer, M., Hasan, M. and Rashid, A. 2017. Impact of extracts of *Azadirachta indica* and *Datura innoxia* on the esterases and phosphatases of three stored grains insect pests of economic importance. *Pakistan Journal of Agricultural Sciences*, 54: 71-81.
- Anwar, M., Ashfaq, M., Hasan, M. and Anjum, F. M. 2005. Efficacy of *Azadirachta indica* oil on bagging material against some insect pests of wheat stored in warehouses at Faisalabad. *Pakistan Entomologist*, 27: 1-5.
- Breuer, M., Hoste, B., Loof, A. D. and Naqvi, S. N. H. 2003. Effect of *Melia azedarach* extract on the activity of NADPH-cytochrome c reductase and cholinesterase in insects. *Pesticide Biochemistry and Physiology*, 76: 99-103.
- CABI, 2007. *Tribolium castaneum* (red flour beetle) Datasheet. *Crop Protection Compendium*, CAB International Publishing, Wallingford, UK.
- Carpinella, C., Ferrayoli, C. and Palacios, M. 2005. Antifungal synergistic effect of scopoletin, a hydroxycoumarin isolated from *Melia azedarach* L. fruits. *Journal of Agricultural and Food Chemistry*, 53: 2922-2927.
- Carpinella, M. C., Defago, M. T., Valladares, G. and Palacios, S. M. 2003. Antifeedant and insecticide properties of a limonoid from *Melia azedarach* (Meliaceae) with potential use for pest management. *Journal of Agricultural and Food Chemistry*, 15: 369-674.
- Coppo, J. A., Mussart, N. B. and Fioranelli, S. A. 2002. Physiological variations of enzymatic activities in blood of Bullfrog, *Rana catesbeina* (Shaw, 1802). *Revista Veterinaria*, 12: 22-27.
- Damayanthi, B. T. and Karunaratne, S. H. P. P.

2005. Biochemical characterization of insecticide resistance in insect pests of vegetables and predatory ladybird beetle. Journal of the National Science Foundation of Sri Lanka, 33: 115-122.
- Defago, M., Valladares, G., Banchio, E., Carpinella, C. and Palacios, S. 2006. Insecticide and antifeedant activity of different plant parts of *Melia azedarach* on *Xanthogaleruca luteola*. Fitoterapia, 77: 500-505.
- El-Naggar, M. E., Abdel-Sattar, M. M. and Mosallam, S. S. 1989. Toxicity of colocynithin and hydrated colocynithin from alcoholic extract of *Citrullus colocynthis* pulp. Journal of the Egyptian Society of Parasitology, 19: 179-185.
- Hong, T. and Ellis, R. 1998. Contrasting seed storage behaviour among different species of Meliaceae. Seed Science and Technology, 26: 77-95.
- Iqbal, J., Irshad, M. and Baloch, U. K. 1992. Insects of stored cereals and their ecology. Proc. FAO Training of Trainers Course on Integrated Pest Management in Food grains, 29 May to 10 June 1989, Islamabad. Pp. 23-28.
- Isman, M. B. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. Annual Review of Entomology, 51: 45-66.
- Kang, J. S., Kim, E., Lee, S. H. and Park, I. K. 2013. Inhibition of acetylcholinesterases of the pinewood nematode, *Bursaphelenchus xylophilus*, by phytochemicals from plant essential oils. Pesticide Biochemistry and Physiology, 105: 50-56.
- Karunakaran, C., Jayas, D. S. and White, N. D. G. 2004. Mass determination of wheat kernels from X-ray images. Biosystems Engineering, 87(3): 267-274.
- Parakrama Karunaratne, S. H. P. 1998. Insecticide resistance in insects. Ceylon Journal of Science Biological Sciences, 25: 72-99.
- Khan, H., Khan, M. A. and Hussan, I. 2007. Enzyme inhibition activities of the extracts from rhizomes of *Gloriosa superba* Linn. (Colchicaceae). Journal of Enzyme Inhibition and Medicinal Chemistry, 22: 722-725.
- Khan, T., Ahmad, M., Nisar, M., Ahmad, M., Lodhi, M. A. and Choudhary, M. I. 2005. Enzyme inhibition and radical scavenging activities of aerial parts of *Paeonia emodi* Wall (Paeoniaceae). Journal of Enzyme Inhibition and Medicinal Chemistry, 20: 245-249.
- Kim, D. H., Yoon, B. H., Kim, Y. W., Lee, S., Shin, B. Y., Jung, J. W., Kim, H. J., Lee, Y. S., Choi, J. S., Kim, S. Y., Lee, K. T. and Ryu, J. H. 2007. The seed extract of *Cassia obtusifolia ameliorates* learning and memory impairments induced by scopolamine or transient cerebral hypoperfusion in mice. Journal of Pharmacological Sciences, 105: 82-93.
- Kim, D. K. 2002. Inhibitory effect of corynoline isolated from the aerial parts of *Corydalis incise* on the acetylcholinesterase. Archives of Pharmacal Research, 25: 817-819.
- Kim, S. W., Kang, J. and Park, I. K. 2013. Fumigant toxicity of Apiaceae essential oils and their constituents against *Sitophilus oryzae* and their acetylcholinesterase inhibitory activity. Journal of Asia-Pacific Entomology, 16: 443-448.
- Koodalingam, A., Mullainadhan, P. and Arumugam, M. 2011. Effects of extract of soapnut *Sapindus emarginatus* on esterases and phosphatases of the vector mosquito, *Aedes aegypti* (Diptera: Culicidae). Acta Tropica, 118: 27-36.
- Lassiter, M. T., Apperson, C. S. and Roe, R. M. 1995. Juvenile hormone metabolism during the fourth stadium and pupal stage of the southern house mosquito, *Culex quinquefasciatus* Say. Journal of Insect Physiology, 41: 869-876.
- Lowe, S. M., Browne, S., Boudjelas S. and DePoorter, M. 2000. 100 of the World's Worst Invasive Alien Species: A selection from the Global Invasive Species Database. Invasive Species Specialist Group, World Conservation Union (IUCN). Available on line at issg.Org/booklet.

- Luo, L., Loon, J. J. A. V. and Schoonhoven, L. M. 1995. Behavioural and sensory responses to some neem compounds by *Pieris brassicae* larvae. *Physiological Entomology*, 20: 134-140.
- Majerus, P. W., Kisseleva, M. V. and Norris, F. A. 1999. The role of phosphatases in inositol signaling reactions. *The Journal of Biological Chemistry*, 274: 10669-10672.
- Mujeeb, K. A. and Shakoori, A. R. 2012. Effect of Fury, a synthetic pyrethroid on esterases of different developmental stages of stored grain pest, Red Flour Beetle, *Tribolium castaneum* (Herbst.) - Spectrophotometric Analysis. *Pakistan Journal of Zoology*, 44: 601-613.
- Nathan, S. S. and Kim, S. 2005. Effects of *Melia azedarach* L. extract on the teak defoliator *Hyblaea puera* Cramer (Lepidoptera: Hyblaeidae). *Crop Protection*, 25: 287-291.
- Nathan, S. S., Choi, M. Y., Seo, H. Y., Paik, C. H., Kalivani, K. and Kim, J. D. 2008. Effect of azadirachtin on acetylcholinesterase (AChE) activity and histology of the brown plant hopper *Nilaparvata lugens* (Stal). *Ecotoxicology and Environmental Safety*, 70: 244-250.
- Nathan, S. S., Kalaivani, K., Murugan, K. and Chung, P. G. 2005. The toxicity and physiological effect of neem limnoids on *Cnaphalocrocis medinalis* (Guenee) the rice leaf folder. *Pesticide Biochemistry and Physiology*, 81: 113-122.
- Niewiada, A., Nawrot, J., Szafranek, J., Szafranek, B., Synak, E., Jelen H. and Wasowicz, E. 2005. Some factors affecting egg-laying of the granary weevil (*Sitophilus granarius* L.). *Journal of Stored Product Research*, 41: 544-555.
- Park, N. J. and Kamble, S. T. 1999. Distribution and inhibition of esterases in body tissues of susceptible and resistant German cockroaches (Dictyoptera: Blattellidae). *Annals of Entomological Society of America*, 92: 556-562.
- Prakash, A. J., Rao, I. C. and Mathur, K. C. 1987. Rice Storage and insect pests management. BR Publishing Corporation, New Delhi, pp, 337.
- Pravin, B., Tushar, D., Vijay, P. and Kishanchnad, K. 2013. Review on *Citrullus colocynthis*. *International Journal of Research in Pharmacy and Chemistry*, 3: 46-53.
- Rajendran, S. and Sriranjini, V. 2008. Plant products as fumigants for stored-product insect. *Journal of Stored Product Research*, 44: 126-135.
- Roy, S. S. 2002. Some toxicological aspects of chlorpyrifos to the intertidal fish, *Boleophthalmus dussumieri*. PhD Thesis University of Mumbai India, pp. 52-71.
- Sagheer, M., Hasan, M., Rashid, A., Ali, K., Majid, A. and Khan, F. Z. A. 2014. Growth regulatory activities of indigenous plant extracts against red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Pakistan Journal of Agricultural Sciences*, 51: 991-995.
- Shanmugavelu, M., Baytan, A. R., Chesnut, J. D. and Bonning B. C. 2000. A novel protein that binds juvenile hormone esterase in fat body tissues and pericardial cells of the tobacco horn worm, *Manduca sexta* L. *The Journal of Biological Chemistry*, 275: 1802-1806.
- Shekari, M., Sendi, J. J., Etebari, K., Zibaee, A. and Shadparvar, A. 2008. Effects of *Artemisia annua* L. (Asteracea) on nutritional physiology and enzyme activities of elm leaf beetle, *Xanthogaleruca luteola* Mull. (Coleoptera: Chrysomellidae). *Pesticide Biochemistry and Physiology*, 91: 66-74.
- Smirle, M. J., Lowery, T. and Zurowski, C. L. 1996. Influence of neem oil on detoxification enzyme activity in the oblique banded leaf roller, *Choristoneura rosaceana*. *Pesticide Biochemistry and Physiology*, 56: 220-230.
- Soam, P. S., Singh, T. and Vijayvergia, R. 2013. *Citrullus colocynthis* (Linn.) and *Luffa acutangula* (L.) roxb, schrad source of bioinsecticides and their contribution in managing climate change. *International Journal of Applied Biology and Pharmaceutical Technology*, 4: 7-9.

- Srinivas, R., Udikeri, S. S., Jayalakshmi, S. K. and Sreeramulu, K. 2004. Identification of factors responsible for insecticide resistance in *Helicoverpa armigera*. *Comparative Biochemistry and Physiology*, 137: 61-269.
- Ukeh, D. A., Oku, E. E., Udo, I. A., Nta, A. I. and Ukeh, J. A. 2012. Insecticidal Effect of Fruit Extracts from *Xylopia aethiopica* and *Dennettia tripetala* (Annonaceae) against *Sitophilus oryzae* (Coleoptera: Curculionidae). *Chilean Journal of Agricultural Research*, 72: 195-200.
- Valladares, G., Defago, M. T., Palacios, S. M. and Carpinella, M. C. 1997. Laboratory evaluation of *Melia azedarach* (Meliaceae) extracts against the elm leaf beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology*, 90: 747-750.
- Wang, X., Li, Q., Shen, L., Yang, J., Cheng, H., Jiang, S., Jiang, C. and Wang, H. 2014. Fumigant, contact, and repellent activities of essential oils against the darkling beetle, *Alphitobius diaperinus*. *Journal of Insect Science*, 14: 1-11.
- White, N. D. G. and Leesch, J. G. 1996. Chemical control. In: Subramanyam, B. and Hagstrum, D. W. (Eds.), *Integrated Management of Insects in Stored Products*. Marcel Dekker, New York-Basel-Hong Kong, pp. 287-330.

فعالیت زیستی عصاره‌های گیاهان دارویی به‌عنوان سموم بازدارنده آنزیمی در برابر حشرات آفات انباری

کاظم علی^{۱*}، محمد ساغیر^۲، منصور الحسن^۲، عبدالرشید^۳ و محمد شهید^۴

۱- مرکز بین‌المللی کشاورزی و علوم زیستی، آسیای مرکزی و غربی، راولپندی، پاکستان.

۲- گروه حشره‌شناسی، دانشگاه کشاورزی، فیصل‌آباد، پاکستان.

۳- گروه بیماری‌شناسی گیاهی، دانشگاه کشاورزی، فیصل‌آباد، پاکستان.

۴- گروه بیوشیمی، دانشگاه کشاورزی، فیصل‌آباد، پاکستان.

پست الکترونیکی نویسنده مسئول مکاتبه: k.ali@cabi.org

دریافت: ۱۳ خرداد ۱۳۹۹؛ پذیرش: ۱۹ مهر ۱۳۹۹

چکیده: پژوهش حاضر برای ارزیابی فعالیت زیستی عصاره گیاه هندوانه ابوجهل *Citrullus colocynthis* و زیتون تلخ *Melia azedarach* علیه سه حشره آفت انباری مهم شامل شپشه آرد *Tribolium castaneum*، لمبه گندم *Trogoderma granarium* و شپشه برنج *Sitophilus granarius* انجام شد. سمیت و فعالیت مهار آنزیم استیل کولین استراز (AChE)، آلفاکربوکسی استراز (α -CE)، بتاکربوکسیل استراز (β -CE)، اسید فسفاتازها (ACP) و آلکالین فسفاتازها (ALP) در سه گونه حشره نسبت به دو عصاره گیاه در چهار غلظت ۵، ۱۰، ۱۵ و ۲۰ درصد ارزیابی شد. حداکثر مرگ‌ومیر در شپشه برنج (۳۴/۲۹ درصد) توسط زیتون تلخ مشاهده شد. در شپشه آرد و لمبه گندم حداکثر مقادیر ثبت شده برای مرگ‌ومیر به ترتیب برابر با ۳۰/۸۷ و ۱۸/۹۵ درصد توسط عصاره زیتون تلخ بود. عصاره گیاه هندوانه ابوجهل بیش‌ترین میزان مرگ‌ومیر را به ترتیب در شپشه آرد، شپشه برنج و لمبه گندم به ترتیب برابر با ۲۱/۹۲، ۱۹/۱۸ و ۱۶/۸۹ درصد بود. یافته‌ها ثابت کرد که هر دو عصاره گیاه اثرات کشنده مناسبی روی گونه‌های حشرات آزمایش شده داشتند. قرار گرفتن حشرات در معرض عصاره‌های گیاهان نیز منجر به مهار معنی‌دار آنزیم‌های AChE، α -CE، β -CE، ACP و ALP شد. همه آنزیم‌های آزمایش شده در سه حشره توسط عصاره گیاه زیتون تلخ حداکثر مهارکنندگی را داشتند، به‌جز عصاره گیاه هندوانه ابوجهل که α -CE در شپشه برنج و ACP در لمبه گندم و شپشه برنج را بیشتر مهار کردند. نتایج نشان می‌دهد که عصاره گیاه زیتون تلخ روی آفات انباری تأثیر بیش‌تری داشت. بنابراین، عصاره‌های گیاهی مطالعه شده به‌عنوان یک جایگزین مطمئن به‌جای سموم شیمیایی برای حفاظت محصولات انباری در برابر حشرات آفت پیشنهاد می‌شوند.

واژگان کلیدی: آفات انباری، مهار آنزیم، اثرات کشنده، سمیت