

Phytoseiid mites (Acari: Mesostigmata: Phytoseiidae) in some regions of western and north western Iran

Bahman Asali Fayaz and Mohammad Khanjani*

Department of Plant Protection, College of Agriculture, Bu Ali-Sina University, Hamadan, I. R. Iran.

Abstract: Study of phytoseiid mites in some regions of western and north-western Iran were carried out during 2008–2011. In this study, 21 species belonging to the six genera were collected and identified. They were associated with aerial parts, soil and litter under cultivated, uncultivated plants and some phytophagous mites and insects.

Keywords: mite, predator, phytophage, fauna, Iran

Introduction

The members of phytoseiid mites are well known predatory mites in the subclass Acari and already some of them are introduced as biological control agents of some phytophagous mites e. g. two spotted spider mites, eriophyid mites and small insect pests such as thrips and whiteflies (Muma, 1971; Meyerdirk and Coudriet, 1986; Sabelis, 1996; Gerson *et al.*, 2003; Nomikou *et al.*, 2003; Asali Fayaz *et al.*, 2011a). The family Phytoseiidae has worldwide distribution and its members are found from the all seven biogeographic regions: Nearctic, Neotropical, Ethiopian, West Palearctic, East Palearctic, Oriental and Australian (Tixier *et al.*, 2008). More than 2280 species were reported (Chant and McMurtry, 2007) and only 75 species were recorded from Iran (Faraji *et al.*, 2007, 2008; Ueckermann *et al.*, 2009; Jafari *et al.*, 2011; Shirdel *et al.*, 2008, 2009, Hajizadeh *et al.*, 2002, 2010; Daneshvar and Denmark, 1982; Daneshvar, 1980, 1987; Khalil-Manesh, 1973; McMurtry, 1977). In this study, 21 species were

collected and identified from some regions of western and northwestern Iran.

Material and Methods

This study was carried out in order to collecting and identifying phytoseiid mites in some regions of western and north-western Iran (Hamedan, Kurdistan, Kermanshah and Ardebil provinces) during 2008–2011. The collected samples were transferred to Acarology laboratory for processing. The mites of the foliage samples were obtained by shaking method and Berlese funnel was used to extract the litters and soil specimens. All specimens were directly mounted on microscopic slides using Hoyer's medium (Walter and Krantz, 2009). All specimens were collected by B. Asali Fayaz. The slides were examined under a Nomarski Olympus BX51 microscope. The classification systems used follows that of Chant and McMurtry (2007) and Denmark (1992). List of hosts, localities and map of distribution of the collected specimens are indicated in Table 1 and Figure 1, respectively. The base of map (Fig. 1) is modified from Google earth software (© 2011 Europe Technologies). All specimens are deposited in the Collection of the Acarology Laboratory, University of Bu-Ali Sina, Hamedan, Iran.

Handling Editor: Dr. Hamidreza Hajiqanbar

* **Corresponding author**, e-mail: mkhanjani@gmail.com
Received: 1 June 2012, Accepted: 9 July 2012

Table 1 Localities status of collected specimens in this study.

Locality	Province	Geographical status	a.s.l. (m)
Fandoghlu forest	Ardabil	38° 23' N, 48° 32' E	1360
Heyran defile	//	38° 26' N, 48° 35' E	1474
Amzajerd greenhouses complex	Hamedan	35° 00' N, 48° 34' E	1669
Asad Abad vicinity	//	34° 46' N, 48° 06' E	1566
Bahar vicinity	//	34° 55' N, 48° 27' E	1709
Faculty of Agriculture, Bu–Ali Sina University	//	34° 48' N, 48° 29' E	1810
Ganjnameh region (Hamedan vicinity)	//	34° 45' N, 48° 26' E	2147
Ghorveh Dar Jazin region (Razan vicinity)	//	35° 21' N, 49° 06' E	1810
Haji Abad village (Famenin vicinity)	//	35° 05' N, 48° 57' E	1624
Hamedan vicinity	//	34° 45' N, 48° 31' E	2015
Heydareh Ghazi Khan village (Bahar vicinity)	//	34° 52' N, 48° 19' E	1874
Heydareh village (Hamedan vicinity)	//	34° 48' N, 48° 28' E	1830
Joraghan village (Hamedan vicinity)	//	34° 54' N, 48° 32' E	1715
Kabodarahangh	//	35° 13' N, 48° 45' E	1663
Malham Abad village (Asad Abad vicinity)	//	34° 49' N, 48° 08' E	1822
Maryanj vicinity	//	34° 49' N, 48° 24' E	1994
Saleh Abad in (Bahar vicinity)	//	34° 55' N, 48° 19' E	1767
Sarabe Gamasiab (Nahavand vicinity)	//	34° 02' N, 48° 22' E	1822
Shahrestaneh region (Toyserkan vicinity)	//	34° 41' N, 48° 21' E	2075
Sanghestan village (Hamedan vicinity)	//	34° 47' N, 48° 35' E	1858
Yekn Abad village (Bahar vicinity)	//	34° 51' N, 48° 27' E	1740
Dalaho vicinity	Kermanshah	34° 33' N, 45° 58' E	1570
Rijab region	//	34° 28' N, 45° 59' E	950
Sahneh vicinity	//	34° 28' N, 47° 41' E	1345
25 Km Sanandaj–Kamyaran	Kurdistan	35° 05' N, 46° 55' E	1309
30 Km Sanandaj–Kamyaran	//	35° 05' N, 46° 42' E	1309
Km 40 Sanandaj–Marivan	//	35° 24' N, 46° 53' E	1705
Baneh vicinity	//	36° 11' N, 46° 13' E	1498
Chenu region (Sanandaj vicinity)	//	35° 08' N, 46° 57' E	1328
Divandareh vicinity	//	35° 54' N, 47° 01' E	1821
Kamyaran vicinity	//	34° 51' N, 46° 56' E	1628
Karvandan village (Dehgolan vicinity)	//	35° 18' N, 47° 22' E	1813
Marivan	//	35° 26' N, 46° 13' E	1320
Naran region (Sanandaj vicinity)	//	35° 00' N, 46° 58' E	1404
Sannadaj vicinity	//	35° 05' N, 46° 45' E	1309
Sarv Abad of (Marivan vicinity)	//	35° 27' N, 46° 13' E	1284

Results

Subfamily Amblyseiinae Muma

Tribe Amblyseiini Muma

Subtribe Amblyseiina

Genus *Amblyseius* Berlese, 1914

Amblyseius obtusus (Koch, 1839)

Zercon obtusus Koch, 1839

World distribution: Armenia, Azerbaijan, Canada, Czech Republic, England, France, Germany (First time from Germany? in gardens, meadows and bank of stream and pool), Greece, Hungary, Italy, Moldova, Morocco, New Zealand, Norway, Poland, Russia, Spain, Sweden, Turkey, Ukraine, USA, Venezuela (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Australian, Holarctic.

Material examined: Sarabe Gamasiab, 24 ix 2009, 2 (♀♀), soil and litter of alfalfa; Marivan, 19 xi 2009, 1 (♀), soil and litter of apple.

Amblyseius rademacheri (Dosse, 1958)

Typhlodromips rademacheri Dosse, 1958: 44.

World distribution: Armenia, Azerbaijan, China, Denmark, Georgia, Germany (First time, on apple), Hungary, Iran, Italy, Japan, Moldova, Netherlands, Russia, Slovakia, South Korea, Switzerland, Ukraine (Moraes *et al.*, 2004) and new record for the collected region of this survey.

Regional distribution: Palearctic, Oriental.

Material examined: Heyran defile, 29 ix 2008, 1 (♀), aerial part of Lady Fern bushes, *Athyrium filix-femina* (L.) Roth (Dryopteridaceae).

Subtribe Proprioseiopsina Chant & McMurtry

Genus *Proprioseiopsis* Muma, 1961

Proprioseiopsis messor (Wainstein, 1960)

Typhlodromus messor Wainstein, 1960: 688.

World distribution: Algeria, Armenia, Australia, Azerbaijan, France, Gaza Strip, Georgia (First time, on grass), Germany, Greece, Israel, Italy, Morocco, New Zealand, South Africa, Spain, Turkmenistan, Ukraine (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007; Nourbakhsh and Kamali, 1995) and new record for the collected region.




Figure 1 Geographic distribution of phytoseiid species in some regions of western and north western Iran (Google earth © 2011 Europe Technologies).

- A.** Hamedan province
B. Kurdistan province
C. Kermanshah province
D. Ardebil province

List of species mentioned at Fig. 1.

N.	Species	N.	Species
1	<i>Amblyseius obtusus</i>	12	<i>Phytoseius ciliatus</i>
2	<i>Amblyseius rademacheri</i>	13	<i>Phytoseius plumifer</i>
3	<i>Proprioseiopsis messor</i>	14	<i>Typhlodromus (Anthoseius) bagdasarjani</i>
4	<i>Neoseiulus agrestis</i>	15	<i>Typhlodromus (Anthoseius) khosrovensis</i>
5	<i>Neoseiulus barkeri</i>	16	<i>Typhlodromus (Anthoseius) rodriguezii</i>
6	<i>Neoseiulus bicaudus</i>	17	<i>Typhlodromus (Anthoseius) rhenanus</i>
7	<i>Neoseiulus marginatus</i>	18	<i>Typhlodromus (Anthoseius) tamaricis</i>
8	<i>Neoseiulus sugonjaevi</i>	19	<i>Typhlodromus (Typhlodromus) athiasae</i>
9	<i>Neoseiulus tauricus</i>	20	<i>Typhlodromus (Typhlodromus) leptodactylus</i>
10	<i>Neoseiulus zwoelferi</i>	21	<i>Typhlodromus (Typhlodromus) tubifer</i>
11	<i>Paragigagnathus insuetus</i>		

Regional distribution: Afrotropical, Australian, Palearctic.

Material examined: Yekn Abad village, 28 v 2010, 1 (♀), aerial part of wall barley, *Hordeum murinum* L. (Poaceae); Divandareh vicinity, 18 v 2011, aerial part of Betony, *Stachys lavandifolia* Vahl (Lamiaceae); Sahneh vicinity, 15 xi 2009, 1 (♀), aerial part of grape, *Vitis vinifera* L. (Vitaceae).

Tribe Neoseiulini Chant & McMurtry, 2003

Genus Neoseiulus Huges, 1948: 141

***Neoseiulus agrestis* (Karg, 1960)**

Typhlodromus agrestis Karg, 1960: 449.

World distribution: Czech Republic, France, Germany (First time, in soil), Hungary, Kazakhstan, Moldova, Netherlands, Russia, Spain, Switzerland, Turkey, Ukraine, USA (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Holarctic

Material examined: Maryanaj vicinity, 22 v 2010, 2 (♀♀), bark of apple tree; Bahar vicinity, 15 vii 2010, 1 (♀), tubers of potato, *Solanum tuberosum* L. (Solanaceae), infested by golden nematode, *Globodera rostochiensis* (Wollenweber) (Heteroderidae).

***Neoseiulus barkeri* Hughes, 1948**

World distribution: Algeria, Australia, Brazil, Canary Islands, Cape Verde, China, England (First time, on germinating barley), Finland, France, Georgia, Germany, Ghana, Greece, Guinea, Israel, Italy, Japan, Jordan, Netherlands, Nigeria, Norway, Reunion Island, Russia, South Africa, South Korea, Spain, Sweden, Turkey, Ukraine, West Bank, Yemen (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Afrotropical, Australian, Neotropical, Oriental, Palearctic.

Material examined: Amzajerj greenhouses complex, 22 v 2010, 1 (♀), aerial part of cucumber shrub, *Cucumis sativus* L. (Cucurbitaceae); 30 Km Sanandaj–Kamyaran, 06 xi 2009, 1 (♀), aerial part of strawberry.

***Neoseiulus bicaudus* (Wainstein, 1962)**

Amblyseius bicaudus Wainstein, 1962: 146.

World Distribution: Armenia, Azerbaijan, Caucasus Region, France, Georgia, Greece, Hungary, Israel, Italy, Kazakhstan (First time on grass), Moldova, Norway, Russia, Spain, Switzerland, Tajikistan, Turkey, Ukraine, USA (Moraes *et al.*, 2004), Japan (Ehara and Amano, 2004), Iran (Asali Fayaz *et al.*, 2011a,b; Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Neotropical, Palearctic.

Material examined: Ganjnameh region, 28 x 2009, 10 (♀♀) and 2 (♂♂), aerial part of Chicory, *Cichorium intybus* L. (Asteraceae), infested by two spotted spider mite (TSSM), *Tetranychus urticae* Koch (Tetranychidae); Heydareh village 16 viii 2010, 2 (♀♀), aerial part of Bermuda grass, *Cynodon dactylon* (L.) Pers. (Poaceae); Saleh Abad, 03,12 xi 2009, 3 (♀♀), soil and litter of Willow, *Salix alba* L. (Salicaceae), and 12 xi 2009, 2 (♀♀), soil and litter of gum bushes, *Astragalus verus* Olivier: Fabaceae; Sarabe Gamasiab, 17 i 2010, 1 (♀), soil and litter of Bermuda grass, *Cynodon dactylon* (L.) Pers. (Poaceae); and 1 (♀), aerial part of *Rosa persica* J. F. Gmel. (Rosaceae); Faculty of Agriculture, Bu–Ali Sina University, 21 v 2010, 1 (♀), aerial part of Yarrow, *Achillea millefolium* L. (Asteraceae), infested by TSSM; 25 Km Sanandaj–Kamyaran, 6 xi 2009, 1 (♀), soil and litter of strawberry shrubs; Naran region **Sanandaj vicinity** 13 xi 2009, 3 (♀♀), soil and litter of raspberry bushes, *Rubus hyrcanus* Juz. (Rosaceae); Fandoghlu forest, 31 vi 2010, 3 (♀♀), soil and litter of Hazelnut trees, *Corylus avellanae* L.

***Neoseiulus marginatus* (Wainstein, 1961)**

Typhlodromus marginatus Wainstein, 1961: 158.

World distribution: Algeria, Armenia, Azerbaijan, France, Georgia, Greece, Hungary, Kazakhstan (First time on herb), Kenya, Moldova, Russia, Turkmenistan, Ukraine (Moraes *et al.*, 2004) Iran (Faraji *et al.*, 2007; Hajizadeh, 2007; Nourbakhsh and Kamali, 1995) and new record for the collected regions of this survey.

Regional distribution: Palearctic

Material examined: Faculty of Agriculture, Bu–Ali–Sina University, 14 vi 2009, 1 (♀), aerial part of white Marshmallow, *Althaea officinalis* L.: Malvaceae, infested by TSSM and 1 (♀), aerial part of yellow melilot, *Melilotus officinalis* (L.) Pall. (Fabaceae), and 15 v 2009, 2 (♀♀), aerial part of Hairy vetch, *Vicia villosa* Roth (Fabaceae), infested by TSSM and 1 (♀), aerial part of Italian bugloss, *Echium italicum* L. (Boraginaceae) infested by TSSM and 21 iv 2010, 1 (♀), aerial part of sickle weed, *Falcaria sioides* (Wibel) Aschers (Apiaceae); Heydareh village, 10 ix 2010, 1 (♀), soil and litter of walnut, *Juglans regia* L. (Juglandaceae), and 11 xi 2009 and 10 ix 2010, 4 (♀♀), soil and litter of peach tree, *Prunus persica* (L.) Batsch (Rosaceae), and 16 xi 2009, 5 (♀♀), soil and litter of apple trees; Joraghan village, 16 iii 2010, 1 (♀), soil and litter of alfalfa; Sanghestan village, 26 iv 2010, 1 (♀), aerial part of Broad leaf plantain, *Plantago major* L. (Plantaginaceae); Kabodarahangh vicinity, 14 xii 2010, 1 (♀), aerial part of Russian olive, *Elaeagnus angustifolia* L. (Elaeagnaceae); Sarabe Gamasiab, 28 ix 2009 and 17 x 2009, 2 (♀♀), aerial part of Syrian mesquite, *Prosopis stephaniana* (M. Bieb.) (Fabaceae), and 1 (♀), soil and litter under apple tree; Kamyaran vicinity, 06 xi 2010, 1 (♀), soil and litter of peach trees, *Prunus persica* (L.) Batsch (Rosaceae); Sarv Abad, 20 v 2009, 1 (♀), soil and litter of apple tree; Kamyaran vicinity, 21 vi 2009, 1 (♀), aerial part of Horse Mint, *Mentha longifolia* (L.) Huds. (Lamiaceae); Sannadaj vicinity, 18 xi 2009, 1 (♀), soil and litter of apricot trees, *Prunus armenica* L. (Rosaceae); Naran village, 18 xi 2009, 1 (♀), aerial part of Field Bindweed bushes, *Convolvulus arvensis* L. (Convolvulaceae); Sahneh vicinity, 20 xi 2009, 1 (♀), soil and litter of alfalfa; Rijab region 16 iv 2010, 1 (♀), aerial part of Common mallow, *Malva neglecta* Wallr. (Malvaceae); Heyran defile, 09x 2008, 1 (♀), aerial part of Lady Fern bushes, *Athyrium filix-femina* (L.) Roth (Dryopteridaceae).

***Neoseiulus sugonjaevi* (Wainstein & Abbasova, 1974)**

Amblyseius sugonjaevi Wainstein & Abbasova, 1974: 796.

World distribution: Azerbaijan (First time in rodent nest), Uzbekistan (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions.

Regional distribution: Palearctic.

Material examined: Sarabe Gamasiab, 23 ix 2009, 1 (♀), soil and litter under wild almond, *Amygdalus scoparia* Spach. (Rosaceae); Asad Abad vicinity, 10 x 2009, 1 (♀), aerial part of alfalfa; Shahrestaneh region, 12 x 2009, 1 (♀), aerial part of walnut trees, *Juglans regia* L. (Juglandaceae); Saleh Abad vicinity, 11 xi 2009, 1 (♀), aerial part of gum. Sanandaj vicinity, 08 x 2009, 1 (♀), aerial part of onion, *Allium cepa* L. (Amaryllidaceae) and 08 x 2009 and 06 xi 2009, 6 (♀♀), soil and litter of tomato, *Lycopersicum esculentum* Miller (Solanaceae), and 05 xi 2009, 1 (♀), soil and litter under Sweet cherry tree, *Prunus avium* L. (Rosaceae), and 2 (♀♀), soil and litter of Plum trees, *Prunus cerasifera* Ehrh. (Rosaceae); Naran region, 07 xi 2009, 2 (♀♀), aerial part, soil and litter of alfalfa, *Medicago sativa* L. (Fabaceae), and 11 xi 2009, 1 (♀), soil and litter of apple trees; Chenu region, 11 xi 2009, 12 (♀♀), aerial part and apple fruits under trees and also 3 (♀♀), soil and litter of tomato. Sahneh vicinity, 07 viii 2009 and 09 xi 2009, 4 (♀♀), aerial part of alfalfa. Heyran region, 27 ix 2008, 1 (♀) and aerial part of Lady Fern bushes, *Athyrium filix-femina* (L.) Roth (Dryopteridaceae).

***Neoseiulus tauricus* (Livshitz & Kuznetsov, 1972)**

Amblyseius tauricus Livshitz & Kuznetsov, 1972: 24.

World distribution: Armenia, Azerbaijan, China, France, Greece, Ukraine (First time on unspecified substrate) (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected region of this survey.

Regional distribution: Palearctic.

Material examined: Faculty of Agriculture, Bu–Ali–Sina University, 21 ix 2011, 2 (♀♀) and aerial part of Yarrow, *Achillea millefolium* L. (Asteraceae).

***Neoseiulus zwoelferi* (Dosse, 1957)**

Typhlodromus zwoelferi Dosse, 1957: 301.

World distribution: Azerbaijan, Finland, Germany (First time on apple), Iran, Israel, Kazakhstan, Montenegro, Norway, Russia, Sweden, Switzerland, Turkey, Ukraine, USA (Moraes *et al.*, 2004), Greece (Papadoulis *et al.*, 2009) and new record for the collected regions of this survey.

Regional distribution: Palearctic

Material examined: Kabodarahangh vicinity, 24 iv 2010, 2 (♀♀), aerial part of alfalfa, *Medicago sativa* L. (Fabaceae), and 20 iii 2010, 1 (♀), soil and litter of alfalfa, 3 (♀♀) soil and litter of Woodland sage plants, *Salvia nemorosa* L. (Lamiaceae); Joraghan village, 08 viii 2009, 2 (♀♀), soil and litter of alfalfa; Heydareh village, 17 ix 2009, 1 (♀), soil and litter of walnut trees, *Juglans regia* L. (Juglandaceae), and 12 xi 2009, 2 (♀♀), soil and litter of apple; Sarabe Gamasiab region, 28 ix 2009, 11 (♀♀) soil and litter of Plane tree, *Platanus orientalis* L. (Platanaceae), and 18 i 2010, 2 (♀♀), soil and litter under Plane tree, *Platanus orientalis* L. (Platanaceae), and 1 (♀), soil and litter under gum bushes, *Astragalus verus* Olivier (Fabaceae); Campus of agricultural faculty, 21 iv 2010, 1 (♀), aerial part of Corn Buttercup plants, *Ranunculus arvensis* L. (Ranunculaceae); Sanghestan village, 26 iv 2010, 4 (♀♀), aerial part of Red clover, *Trifolium pretense* L. (Fabaceae) and 4 (♀♀), aerial part of Yarrow, *Achillea millefolium* L. (Asteraceae); Yekn Abad village, 05 iv 2010, 1 (♀), aerial part of Wall barley, *Hordeum murinum* L. (Poaceae); Hamedan vicinity, 15 viii 2010, 2 (♀♀), soil under gum bushes, *Astragalus verus* Olivier (Fabaceae); Haji Abad village, 2 (♀♀) aerial part of alfalfa; Heydareh Ghazi Khan village, 3 (♀♀), unknown host (soil); Baneh vicinity, 15 v 2010, 3 (♀♀), soil and litter of Sweet cherry tree, *Prunus avium* L. (Rosaceae); Veinesar region of Ghorveh vicinity, 09 viii 2009, 2 (♀♀) aerial part of alfalfa; Sanandaj vicinity, 09 viii 2009, 1 (♀), aerial part of

alfalfa; Karvandan village, 11 viii 2009 and 02 ix 2009, 8 (♀♀), aerial part of alfalfa; Kamyaran vicinity, 07 x 2009, 1 (♀) soil and litter under plum tree, *Prunus cerasifera* Ehrh. (Rosaceae); and 1 (♀), soil and litter of tomato; Karvandan village, 24 x 2009, 3 (♀♀), aerial part of alfalfa; Km 40 Sanandaj–Marivan, 24 x 2009, 1 (♀), soil and litter of alfalfa, *Medicago sativa* L. (Fabaceae), and 2 (♀♀), soil and litter under tomato; Sanandaj vicinity, 07 xi 2009, 2 (♀♀), soil and litter under apricot trees, *Prunus armenica* L. (Rosaceae), and 2 (♀♀), soil and litter under sour cherry, *Prunus cerasus* L. (Rosaceae); Naran region, 13 xi 2009, 1 (♀), aerial part of Horse Mint, *Mentha longifolia* (L.) Huds. (Lamiaceae); and aerial part of alfalfa; Chenu region, 19 xi 2009, 13 (♀♀), aerial part of apple trees, *Mallus domestica* Moller (Rosaceae), infested by Woolly apple aphid, *Eriosoma lanigerum* (Hausmann) (Aphididae) and 25 (♀♀), apple fruit under trees and 2 (♀♀), soil and litter of tomato, *Lycopersicon esculentum* Miller (Solanaceae), and 1 (♀), aerial part of raspberry bushes, *Rubus hyrcanus* Juz. (Rosaceae); Kamyaran vicinity, 02 i 2010, 1 (♀), aerial part of Cherry plum; Dalaho vicinity, 1 (♀) unknown host (soil).

Remarks: This species was observed more than other species which were associated with low-growth plants e. g. herbs and also soil and litter of plants.

Genus *Paragigagnathus* Amitai & Grinberg, 1971

***Paragigagnathus insuetus* (Livshitz & Kuznetsov, 1972)**

Amblyseius insuetus Livshitz & Kuznetsov, 1972: 27.

World distribution: Greece, Turkmenistan, Ukraine (First time on *Tamarix* sp.) (Moraes *et al.*, 2004), Iran (Hajizadeh *et al.*, 2010) and new for the collected regions of this survey.

Region distribution: Palearctic.

Material examined: Heyran defile, 06 xii 2008, 1 (♀), aerial part of Salt cedar tree, *Tamarix gallica* L. (Tamaricaceae).

Subfamily Phytoseiinae Berlese

Genus *Phytoseius* Ribaga, 1904***Phytoseius ciliatus* Wainstein, 1975**

Phytoseius (Dubininellus) ciliatus Wainstein, 1975: 921.

World distribution: Russia (First time on unspecified substrate), Iran (Moraes *et al.*, 2004) and new for the collected region of this survey.

Region distribution: Palearctic.

Material examined: Fandoghlu forest, 04 xi 2009 and 17 x 2009, 2 (♀♀), aerial part of Hazelnut trees, *Corylus avellanae* L. (Corylaceae).

***Phytoseius plumifer* (Canestrini & Fanzago, 1876)**

Gamasus plumifer Canestrini & Fanzago, 1876: 130.

World distribution: Algeria, Armenia, Azerbaijan, Egypt, France, Georgia, Hungary, Israel, Italy (First time on nettle), Jordan, Kazakhstan, Lebanon, Portugal, Ukraine, USA (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Holarctic.

Material examined: Sarabe Gamasiab region, 28 ix 2009, 4 (♀♀), aerial part of Yellow goats beard, *Tragopogon major* Jacq (Asteraceae) and 17 x 2010, 1 (♀), aerial part of Chicory, *Cichorium intybus* L. (Asteraceae); Heydareh village, 06 xi 2009, 3 (♀♀), aerial part of apple tree; Hamedan vicinity, 24 vii 2010, 1 (♀), aerial part of Quince trees, *Cydonia oblonga* Mill. (Rosaceae); Sanandaj vicinity, 09 x 2009, 4 (♀♀), aerial part of raspberry bushes, *Rubus hyrcanus* Juz. (Rosaceae); Fandoghlu forest, 05 x 2008, 19 (♀♀), leaf of Fig, *Ficus carica* L. (Moraceae), and 05 x 2008, 4 (♀♀), aerial part of Lady Fern bushes, *Athyrium filix-femina* (L.) Roth (Dryopteridaceae), and 05 x 2008, 1 (♀), aerial part of Cypress tree, *Cupressus sempervirens* L. (Cupressaceae) and 05 x 2008, 1 (♀), aerial part of Field Bindweed bushes, *Convolvulus arvensis* L. (Convolvulaceae).

Subfamily Typhlodrominae Wainstein**Tribe Typhlodromini Wainstein, 1962****Genus *Typhlodromus* Scheuten, 1857*****Typhlodromus (Anthoseius) bagdasarjani***

Typhlodromus bagdasarjani Wainstein & Arutunjan, 1967: 1765.

Amblydromella (Aphanoseia) bagdasarjani Denmark & Welbourn (2002)

World distribution: Armenia (First time on fruit tree), Azerbaijan, Turkmenistan (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007; Asali Fayaz *et al.*, 2011a) and new record for the collected regions of this survey.

Regional distribution: Palearctic.

Material examined: Faculty of Agriculture, 25 x 2008 and 10 xi 2009, 2 (♀♀), aerial part of Cypress tree, *Cupressus sempervirens* L. (Cupressaceae), infested by *Cenopalpus pulcher* (Canestrini and Fanzago, 1876) (Tenuipalpidae) and also 10 xi, 14 x 2009, 3 (♀♀), aerial part of Plane tree, *Platanus orientalis* L. (Platanaceae), and 24 ix 2009, 1 (♀), aerial part of apple infested by *Bryobia rubrioculus* (Scheuten) (Tetranychidae); Heydareh village, 09 ix 2009, 1 (♀), aerial part of apple; Ghorveh Dar Jazin region, [20–23 x 2009, 12 (♀♀), aerial part of grape, *Vitis vinifera* L.: Vitaceae, 23 x 2009, 1 (♀), aerial part of apple]; Malham Abad village, 19 iii 2011, 3 (♀♀), aerial part of plum infested by scale, *Tecaspis asiatica* Balachowsky (Diaspididae); Sanandaj vicinity, 09 x 2009, 2 (♀♀), aerial part of apple and 1 (♀), aerial part of sour cherry, *Prunus cerasus* L. (Rosaceae), and 1 (♀), aerial part of eggplant, *Solanum melongena* L. (Solanaceae), and 1 (♀), aerial part of Peach tree, *Prunus persica* (L.) Batsch (Rosaceae); Chenu region, 18 x 2009, 1 (♀), aerial part of Sweet cherry tree; 09 xi 2009, 2 (♀♀), aerial part of apple; Sanandaj vicinity, 14 xi 2009, 3 (♀♀), aerial part of apple.

***Typhlodromus (Anthoseius) khosrovensis* (Arutunjan, 1971)**

Amblydromella khosrovensis Moraes *et al.*, (1986)

Typhlodromus khosrovensis Arutunjan, 1971: 306.

World distribution: Armenia (First time on *Betula* sp.) (Moraes *et al.*, 2004), Iran (Ueckermann *et al.*, 2009; Jafari *et al.*, 2011; Asali Fayaz *et al.*, 2011a) and new record for the collected regions of this survey.

Regional distribution: Palearctic.

Material examined: Faculty of Agriculture, [08 x 2008, 1 (♀), aerial part of Plane tree, *Platanus orientalis* L. (Platanaceae), 28 x 2008, 1 (♀), aerial part of apple and 01 viii 2010, 12 (♀♀), aerial part of plum infested by Brown mite, *Bryobia rubrioculus* (Scheuten) (Tetranychidae)]; Heydareh village, 06 xi 2009, 2 (♀♀), aerial part of apple tree infested by TSSM and 1 (♀), aerial part of Peach tree, *Prunus persica* (L.) Batsch (Rosaceae).

***Typhlodromus* (*Anthoseius*) *rhenanus* (Oudemans, 1905)**

Amblydromella (*Aphanoseia*) *rhenana* (Denmark and Welbourn (2002)

Seiulus rhenanus Oudemans, 1905: 78.

World distribution: Algeria, Azerbaijan, Belgium, Byelorussia, Canada, Cyprus, Denmark, England, Finland, France, Germany (First time on rotting leaves), Hungary, India, Iran, Israel, Italy, Kazakhstan, Madeira Island, Moldova, Montenegro, Netherlands, Northern Ireland, Norway, Poland, Portugal, Russia, Sweden, Switzerland, Turkey, Ukraine, USA (Moraes *et al.*, 2004) and new record for the collected regions of this survey.

Region distribution: Holarctic, Oriental.

Material examined: Heyran region, 28 ix 2008, 2 (♀♀), aerial part of Lady Fern bushes, *Athyrium filix-femina* (L.) Roth (Dryopteridaceae.)

***Typhlodromus* (*Anthoseius*) *rodriguezi* (Denmark & Daneshvar, 1982)**

Amblydromella rodriguezi Denmark & Daneshvar, 1982: 11.

World distribution: Iran (First time on *Malus* sp.)

Region distribution: Palearctic.

Material examined: Faculty of Agriculture, 17 x 2008, 1 (♀), unknown host.

***Typhlodromus* (*Anthoseius*) *tamaricis* (Kolodochka, 1982)**

Anthoseius (*Amblydromellus*) *tamaricis* Kolodochka, 1982: 11.

Amblydromella (*Aphanoseia*) *tamaricis* Denmark & Welbourn, 2002

World distribution: Turkmenistan (First time on *Tamarix* sp.) (Moraes *et al.*, 2004), Turkey (Kasap and Çobanoğlu, 2009).

Regional distribution: Palearctic.

Material examined: Heyran region, 28 ix 2008, 2 (♀♀), aerial part of Salt cedar tree.

***Typhlodromus* Scheuten, 1857**

Typhlodromus (*Typhlodromus*) *athiasae* (Porath & Swirski, 1965)

Typhlodromus athiasae Porath & Swirski, 1965: 60.

World distribution: Azerbaijan, Egypt, Greece, Iran, Israel (First time on *Citrus* sp.), Jordan, Turkey (Moraes *et al.*, 2004) and new record for the collected regions of this survey.

Regional Distribution: Palearctic.

Material examined: Heyran region, 28 ix 2008, 6 (♀♀), aerial part of Cypress tree *Cupressus sempervirens* L. (Cupressaceae).

Typhlodromus* (*Typhlodromus*) *leptodactylus

Typhlodromus leptodactylus Wainstein, 1961: 153.

Typhlodromus (*Oudemanus*) *leptodactylus* (Denmark, 1992)

World distribution: Armenia, Azerbaijan, Georgia (First time on *Juniperus* sp.), Israel, Ukraine (Moraes *et al.*, 2004), Iran (Faraji *et al.*, 2007) and new record for the collected regions of this survey.

Regional distribution: Palearctic.

Material examined: Heyran region, 26 xii 2008, 2 (♀♀) aerial part of Cypress tree, *Cupressus sempervirens* L. (Cupressaceae).

***Typhlodromus* (*Typhlodromus*) *tubifer* (Wainstein, 1961)**

Typhlodromus tubifer Wainstein, 1961: 157.

World distribution: Armenia, Azerbaijan, Belgium, Caucasus Region, Georgia (unspecified substrate), Iran, Moldova, Turkey (Moraes *et al.*, 2004) and new record for the collected regions of this survey.

Regional distribution: Palearctic.

Material examined: Fandoghlu forest, 20 v 2009, 1 (♀), aerial part of Hazelnut tree, *Corylus avellanae* L. (Corylaceae).

Acknowledgment

This paper is a part of M. Sc. thesis of the senior author which was financially supported by Vice Research of Bu–Ali Sina University, Hamedan, Iran. Also, we wish to thank Prof. H. A. Denmark (Emeritus Chief of Entomology, Entomology Bureau, division of Plant Industry, Florida Department of Agriculture and Consumer services, P. O. Box 147100, Gainesville, Florida 32614-7100, U.S.A) for critical review of the manuscript and also we would like appreciate Prof. Edward A. Ueckermann (ARC-Plant Protection Research Institute, Private Bag X134, Queenswood, Pretoria, 0121 South Africa and School of Environmental Sciences and Development, North–West University, Potchefstroom Campus 2520, South Africa) for supplying some literatures.

References

- Amitai, S. and Grinberg, T. 1971. Description of a new phytoseiid genus and species (Acarina: Mesostigmata) from Israel. *Israel Journal of Entomology*, 6: 327-335.
- Arutunjan, E. S. 1971. New species of the genus *Typhlodromus* Scheuten, 1857 (Parasitiformes, Phytoseiidae) [in Russian]. *Doklady Akademii Nauk Armyanskoi SSR*, 52: 305-308.
- Asali Fayaz, B., Khanjani, M., Molavi, F. and Ueckermann, E. A. 2011a. Phytoseiid mites (Acari: Phytoseiidae) of apple and almond trees in regions of western and south-western Iran. *Acarologia*, 51: 371-379.
- Asali Fayaz, B., Khanjani, M. and Ueckermann, E. A. 2011b. Description of immature stages and re–description of female and male of *Neoseiulus bicaudus* (Wainstain), 1962 (Acari: Phytoseiidae) from west of Iran. *Acta Phytopathologica et Entomologica Hungarica*, 46: 329-338.
- Canestrini, G. and Fanzago, F. 1876. Nuovi acari italiani (Seconda Serie). *Atti Società Veneto-Trentina di Scienze Naturali*, 5: 130-142.
- Chant, D. A. and McMurtry, J. A. 2003. Review of the subfamily Amblyseiniinae Muma (Acari: Phytoseiidae): Part I. Neoseiulini new tribe. *International Journal of Acarology*, 29: 3–46.
- Chant, D. A. and McMurtry, J. A. 2007. Illustrated keys and diagnoses and subgenera of the Phytoseiidae of the world (Acari: Mesostigmata). Indira Publishing House, West Bloomfield.
- Chant, D. A. and Yoshida–Shaul, E. 1992. Adult idiosomal setal patterns in the family Phytoseiidae (Acari: Gamasidae). *International Journal of Acarology*, 18 (3): 177-193.
- Daneshvar, H. 1980. Some predator mites from northern and western Iran. *Entomologie et Phytopathologie Appliquees*, 48: 15–17 (In English), 87–96 (In Persian).
- Daneshvar, H. 1987. Some predatory mites from Iran, with descriptions of one new genus and six new species (Acari: Phytoseiidae, Ascidae). *Entomologie et Phytopathologie Appliquees*, 54 (1-2): 13-37 (In English), 55-73 (In Persian).
- Daneshvar, H. and Denmark, H. A. 1982. Phytoseiids of Iran (Acarina: Phytoseiidae). *International Journal of Acarology*, 8: 3-14.
- Denmark, H. A. 1992. A revision of the genus *Typhlodromus* Scheuten (Acari: Phytoseiidae). *Occasional Papers of the Florida State Collection of Arthropods*, 7: 1-43.
- Denmark, H. A. and Welbourn, W. C. 2002. Revision of the genera *Amblydromella* Muma and *Anthoseius* De Leon (Acari: Phytoseiidae). *International Journal of Acarology*, 28 (4): 291–316.
- Dosse, G. 1957. Morphologie und biologie von *Typhlodromus zwoelferi* n. sp. (Acari, Phytoseiidae). *Zeitschrift für Angewandte Entomologie*, 41 (2–3): 301–311.
- Dosse, G. 1958. Ubereinigenue Raubmilbenarten (Acari: Phytoseiidae). *Pflanzenschutz Berichte*, 21: 44–61.
- Ehara, S. and Amano, H. 2004. Checklist and Keys to Japanese Amblyseiniinae (Acari:

- Gamasina: Phytoseiidae). Journal of the Acarological Society of Japan, 13 (1): 1-30.
- Faraji, F., Hajizadeh, J., Ueckermann, E.A., Kamali, K. and McMurtry, J. A. 2007. Two new records for Iranian phytoseiid mites with synonymy and keys to the species of *Typhloseiulus* Chant and McMurtry and Phytoseiidae in Iran (Acari: Mesostigmata). International Journal of Acarology, 33 (3): 231–239.
- Faraji, F., Sakenin-Chelav, H., Kamali, K. and McMurtry, J. A. 2008. Four new species records of Phytoseiidae (Acari: Mesostigmata) for Iran, and description of variability in the spermatheca of *Typhlodromus Bakeri*. Systematic and Applied Acarology, 13: 123–132.
- Gerson, U., Smiley, R. L. and Ochoa, R. 2003. Mites (Acari) for pest control. Blackwell Publishing, Oxford.
- Hajizadeh, J. 2007. Phytoseiid mites fauna of Guilan province, part II: subfamilies Amblyseinae Muma and Phytoseiinae Berlese (Acari: Phytoseiidae). Agriculture Research, 7 (1): 7–25 [In Persian].
- Hajizadeh, J., Hosseini, R. and McMurtry, J. A. 2002. Phytoseiid mites (Acari: Phytoseiidae) associated with eriophyid mites (Acari: Eriophyidae) in Guilan province of Iran. International Journal of Acarology, 28 (4): 373–378.
- Hajizadeh, J., Faraji, F., Rafatifard, M. and Kamranfard, F. 2010. The genus *Paragigagnathus* Amitai and Grinberg (Acari: Phytoseiidae) in Iran, with a key to the known species. Systematic and Applied Acarology, 15: 222–227.
- Jafari, S., Fathipour, Y. and Faraji, F. 2011. Redescriptions of *Amblyseius meghriensis* Arutunjan and *Typhlodromus haiastanius* (Arutunjan) with discussion on using preanal pores as a character in the subgenus *Anthoseius* (Mesostigmata: Phytoseiidae). International Journal of Acarology, 37 (3): 244: 254.
- Karg, W. 1960. Zur Kenntnis der Typhlodromiden (Acarina: Parasitiformes) aus Ackerund Grünlandboden. Zeitschrift für Angewandte Entomologie, 47: 440–452.
- Kasap, I. and Çobanoğlu, S. 2009. Phytoseiid mites of Hakkâri province, with *Typhlodromus (Anthoseius) tamaricis* Kolodochka, 1982 (Acari: Phytoseiidae), a new record for the predatory mite fauna of Turkey. Turkish Journal of Zoology, 33 (3): 301–308.
- Khalil-Manesh, B. 1973. Phytophagous mite fauna of Iran. Applied Entomology and Phytopathology, 35: 30–38. [In Persian]
- Koch, C. L. 1839. Deutschlands Crustaceen, Myriapoden und Arachniden. Regensburg, Germany, 6: 1–13.
- Kolodochka, L. A. 1982. New phytoseiid mites (Parasitiformes: Phytoseiidae) from Turkmen. Vestnik Zoologii, 6: 7–13 [in Russian].
- Livshitz, I. Z. and Kuznetsov, N. N. 1972. Phytoseiid mites from Crimea (Parasitiformes: Phytoseiidae). Proceedings of The All-Union V. I. Lenin Academy of Agricultural Science, The State Nikita Botanical Gardens, Yalta, Ukraine, pp. 13–64 [in Russian].
- McMurtry, J. A. 1977. Description and biology of *Typhlodromus persianus*, n. sp., from Iran, with notes on *T. kettanehi* (Acarina: Mesostigmata: Phytoseiidae). Annals of the Entomological Society of America, 70: 563–568.
- Meyerdirk, D. E. and Coudriet, D.L. 1986. Evaluation of two biotypes of *Euseius scutalis* as predators of *Bemesia tabaci*. Journal of Economic Entomology, 79 (3): 659–663.
- Moraes, G. J. De., McMurtry, J. A., Denmark, H. A., and Campos C. B. 2004. A revised catalog of the mite family Phytoseiidae. Zootaxa, 434: 1–494.
- Muma, M. H. 1961. Subfamilies, genera, and species of Phytoseiidae (Acarina: Mesostigmata). Florida State Museum Bulletin, 5 (7): 267–302.
- Muma, M. H. 1971. Food habits of Phytoseiidae (Acarina: Mesostigmata) including common species of Florida citrus. The Florida Entomologist, 54: 21-34.
- Nadimi, A., Kamali, K., Arbabi M. and Abdoli, F. 2009. Selectivity of three miticides to spider mite predator, *Phytoseius plumifer* (Acari: Phytoseiidae) under laboratory conditions. Agricultural Sciences in China, 8 (3): 326–331.

- Nomikou, M., Janseen, A. and Sabelis, M. W. 2003. Phytoseiid predators of whiteflies feed and reproduce on non-pray food sources. *Experimental and Applied Acarology*, 31: 15–26.
- Nourbakhsh, S. H. and Kamali, K. 1995. Biology of Brown mite (*Petrobia latens* Muller) in eastern Chahar–Mahaland Bakhtiari province (Iran). *Journal of Entomological Society of Iran*, 15: 15–24 [In Persian].
- Oudemans, A. C. 1905. Verslag van de zestigsteezomervergadering der Nederlandsche Entomologische Vereeniging, gehouddemtedriebergen op zaterdag, 20 Mei 1905, des morgens ten 11 ure. *Tijdschr. Tijdschriftvoor Entomologie*, The Netherlands, 48: 77–81.
- Papadoulis, G. T., Emmanouel, N. G. and Kapaxidi E. V. 2009. Phytoseiidae of Greece and Cyprus: (Acari: Mesostigmata). Indira Publishing House, West Bloomfield.
- Porath, A. and Swirski, E. 1965. A survey of phytoseiid mites (Acarina: Phytoseiidae) on citrus, with a description of one new species. *Israel Journal of Agricultural Research*, 15: 87–100.
- Rowell, H. J., Chant, D. A. and Hansell, R. I. C. 1978. The determination of setal homologies and setal patterns on the dorsal shield in the family Phytoseiidae (Acarina: Mesostigmata). *The Canadian Entomologist*, 110: 859–876.
- Sabelis, M. W. 1996. Phytoseiidae. In: Lindquist, E. E., Sabelis, M. W. and Bruin, J. (Eds.) *Eriophyoid mites, their biology, natural enemies and control*. Elsevier Science Publishing, Amsterdam, The Netherlands, pp. 427–456.
- Shirdel, D., Kamali, K. and Faraji, F. 2008. Redescription of *Typhloseiulus carmonae* (Chant and Yoshida-Shaul) (Mesostigmata: Phytoseiidae) new species for Iran. *Acarina*, 16 (1): 51–56.
- Shirdel, D., Arbabi, M. and Faraji, F. 2009. *Amblyseius ampullosus* Wu and Lan (Acari: Phytoseiidae), a new species record for the Iranian fauna. *Systematic and Applied Acarology*, 14: 136–139.
- Tixier, M. S., Kreiter, S. and Moraes, G. J. De 2008. Biogeographic distribution of Phytoseiidae (Acari: Mesostigmata). *Biological Journal of the Linnean Society*, 93: 845–856.
- Ueckermann, E. A., Jalaieian, M., Saboori, A. and Seyedoleslami, H. 2009. Redescription of *Typhlodromus (Anthoseius) khosrovensis*, first record for Iran (Acari: Phytoseiidae). *Acarologia*, 49 (1–2): 23–27.
- Walter, D. E. and Krantz, G. W. 2009. Collection, rearing and preparing specimens In: Krantz G. W. and Walter D. E. [Eds.]. *A Manual of Acarology*. 3rd Ed. Texas Tech University Press, Lubbock.
- Wainstein, B.A. 1960. New species and subspecies of the genus *Typhlodromus* Scheuten (Parasitiformes, Phytoseiidae) of the USSR fauna. *Zoologicheskii Zhurnal*, 39: 683–690 [in Russian].
- Wainstein, B.A. 1961. New species of mites of the genus *Typhlodromus* (Parasitiformes: Phytoseiidae) in Georgia. *Trudy Instituta Zoologii Akademii Nauk Gruzinskoy SSR*, 18: 153–162 [in Russian].
- Wainstein, B. A. 1962. Some new predatory mites of the family Phytoseiidae (Parasitiformes) of the USSR fauna. *Entomologicheskoe Obozrenie*, Russia. 41: 230–240 [in Russian]; *Entomological Review*, 41: 139–146 [English translation].
- Wainstein, B. A. 1975. Predatory mites of the family Phytoseiidae (Parasitiformes) of Yaroslavl Province. *Entomologicheskoe Obozrenie*, Ruussia, 54 (4): 914–922 [In Russian].
- Wainstein, B. A. and Arutunjan, E. S. 1967. New species of predaceous mites of the genera *Typhlodromus* Scheuten and *Paraseiulus* Muma (Parasitiformes, Phytoseiidae) *Zoologicheskii Zhurnal*, 46: 1764–1770 [In Russian].
- Wainstein, B. A. and Abbasova, E. D. 1974. Two new species of the genus *Amblyseius* (Parasitiformes: Phytoseiidae) from Azerbaijan. *Zoologicheskii Zhurnal*, 3: 796–798 [In Russian].

کنه‌های فیتوزییده (Acari: Mesostigmata: Phytoseiidae) در برخی از مناطق غربی و شمال غربی ایران

بهمن عسلی فیاض و محمد خانجانی*

گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه بوعلی سینا، همدان، ایران.

* پست الکترونیکی مسئول مکاتبه: mkhanjani@gmail.com

چکیده: مطالعه کنه‌های فیتوزییده در برخی از مناطق غربی و شمال غربی ایران در طی سالهای ۱۳۸۷-۱۳۹۰ صورت گرفت. در این مطالعه، ۲۱ گونه در قالب شش جنس مرتبط با قسمت‌های هوایی، خاک زیر گیاهان زراعی و غیر زراعی و برخی از کنه‌ها و حشرات گیاهخوار جمع‌آوری و شناسایی شد.

واژگان کلیدی: کنه، شکارگر، گیاه‌خوار، فون، ایران.