

Lethal and sublethal effects of abamectin and deltamethrin on potato tuber moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae)

Hooshang Rafiee-Dastjerdi^{1*}, Zainab Mashhadi¹ and Aziz Sheikhi Garjan²

1. Department of Plant Protection, Faculty of Agricultural, University of Mohaghegh Ardabili, Ardabil, Iran.

2. Iranian Research Institute of Plant Protection P. O. Box: 19395-1454, Tehran, Iran.

Abstract: The potato tuber moth (PTM), *Phthorimaea operculella* (Zeller) is one of the important pests of potato in tropical regions. In this research, the toxicity of two insecticides, abamectin and deltamethrin on *P. operculella* was studied at 26 ± 1 °C, $65 \pm 5\%$ RH and photoperiod of 16:8 h (L: D). The concentrations were determined by preliminary dose setting experiments. Distilled water was used as control. LC_{50} values for egg, first instar larvae and adult stages in abamectin were 0.92, 0.014 and 0.46 mg ai/l and in deltamethrin were 0.09, 0.024 and 0.29 mg ai/l, respectively. The sublethal effects of abamectin and deltamethrin on population growth parameters of *P. operculella* were determined at mentioned conditions. Four hundred 1 day old eggs of the pest were dipped in insecticides solutions (LC_{30}). After egg hatching, the first instar larvae were transferred on potato tubers and placed in transparent plastic dishes. After adults' emergence, 20 females were used for each treatment in life table experiments. In control, abamectin and deltamethrin treatments intrinsic rates of increase were estimated to be 0.060, 0.042 and 0.141 day⁻¹, respectively. Statistical analyses showed that deltamethrin had the higher toxicity to egg and adult stages of PTM and abamectin had the higher toxicity to its first instar larvae. Since deltamethrin increased the stable population parameters of *P. operculella* specially its intrinsic rate of increase, thus it might not be recommended for control of PTM. However based on lethal and sublethal effects, abamectin could be suitable for management of this pest.

Keywords: *Phthorimaea operculella*, lethal, sublethal, insecticides, demography

Introduction

The potato tuber moth (PTM), *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae), is the major world wide pest of potato (*Solanum tuberosum* L.), tobacco (*Nicotiana tabacum* L.), tomato (*Lycopersicon esculentum* Mill.), eggplant and pepper in fields and stores, especially in warm temperate and subtropical

climates (Sporleder *et al.*, 2004). Pest management tactics including mass trapping by pheromones and cultural and chemical control methods have been applied to manage of PTM (Hanafi, 1999). There are several reports for resistance of tuber moth to some insecticides from many countries (Haines, 1977). Frequently use of synthetic insecticides causes problems such as resurgence. Resurgence may occur in response to physiological or ecological mechanisms (Ripper, 1956; Luckey, 1968). In order to introduce new insecticides for control of PTM, in present study, lethal and sublethal effects of two insecticides, abamectin and deltamethrin were investigated on

Handling Editor: Dr. Ahad Sahragard

*Corresponding author, e-mail: rafiee@uma.ac.ir
Received: 26 July 2012; Accepted: 29 July 2013

P. operculella. Abamectin produced by the soil micro organism, *Streptomyces avermilitis* acts as insecticide against agricultural pests (White *et al.*, 1997). Deltamethrin is a pyrethroid insecticide that is used to control a range of pests. Deltamethrin paralyzes the insect nervous system rapidly, giving a quick knockdown effect (Haug And Hoffman, 1990). Abamectin has a rapidly disabling effect on feeding behavior of insects (Hayes and Laws, 1990).

Lethal assays could not completely determine the effects of insecticides on organisms (Walthall and Stark 1996). Thus, evaluating sublethal effect of insecticides is very important (Stapel *et al.*, 2000; Stark and Banks 2003). Demographic toxicology is usually considered the best way to evaluate total effects of pesticides. Sublethal doses can make changes in development time (Schneider *et al.*, 2003), percent of hatched eggs and number of laid eggs per female (Wang *et al.*, 2009), fecundity (Haseeb and Amano, 2002, reproduction parameters (Lashkari *et al.*, 2007), and sex ratio (Delpuech and Meyet, 2003). Then, the parameter defined as the intrinsic rate of increase (r_m) has been recommended to evaluate total effects of pesticides, because it is based on both survivorship and fecundity parameters (Stark and Wennergren 1995). This study was carried out to assess the lethal and sublethal effect of the insecticides on *P. operculella* and to determine the possibility of the application of these insecticides at reduced dose for control of this pest.

Materials and Methods

Insecticides

The tested insecticides were abamectin (Agrimec[®], 1.8% EC, Gyah Company, www.gyah.ir) and Deltamethrin (Decis[®] 2.5% EC, Gyah Company).

Insect rearing

Initial population of *P. operculella* was obtained from the University of Mohaghegh Ardabili, Ardabil, Iran. Adults of PTM were released in transparent plastic container (12 × 9 cm) to lay

eggs. The bottom of containers was covered with a filter paper and a slice of potato (Agria cultivar). Insects were reared on potato tubers at 26 ± 1 °C and $65 \pm 5\%$ RH and photoperiod of 16:8 h (L: D) in a growth chamber.

Bioassays

Lethal

The concentrations were determined by preliminary dose setting experiments. Distilled water was used as control. One-day-old eggs of tuber moth on the paper disc were dipped in different concentrations of insecticides for 10 seconds and then they were transferred to Petri dishes up to hatch (Sauor, 2008). For adult bioassay, Petri dishes were impregnated with insecticides at different concentrations and then PTM adults were transferred to them (Dogramaci and Tingey, 2008). Also, the toxicity of insecticides was assessed on first larvae of PTM using leaf dipping bioassay method. Potato leaves were dipped in the insecticide solutions for 10 seconds and let dry at room temperature for 15 minutes, then the larvae were transferred on them into Petri dishes (Symington, 2003). The mortalities were recorded 24 h after treatment in the larvae and adult stages and after hatching in the egg stage. Each experiment was replicated four times.

Sublethal

Four hundred one-day-old eggs of the pest were dipped in sublethal concentration (LC_{30}) of each insecticide. Then, they were transferred into 90mm Petri dishes. Distilled water was used as control. After egg hatching, the first instar larvae were transferred on potato tubers and placed in transparent plastic containers (4 × 8 cm). After adults' emergence, 20 females from each of the insecticide treatment and the control were individually transferred into plastic vials (5 cm in diameter and 10 cm height). The number of eggs laid by females was recorded daily until death of all females. All experiments were conducted at 26 ± 1 °C, $65 \pm 5\%$ RH and photoperiod of 16: 8 h (L: D). Carey's procedure was used to estimate the population growth parameters (Carey, 1993).

Data analysis

The biological data were submitted to analysis of variance and the means were compared with Tukey's tests using SPSS 16.00 software program (SPSS, 2004). The demographic parameters and their corresponding standard errors were estimated by the jackknife technique (Meyer, 1986).

Results

The toxicity of insecticides, abamectin and deltamethrin to eggs, larvae and adults of *P. operculella* is shown in Table 1. The results indicated that both tested insecticides had toxic and acceptable effect on eggs of *P. operculella* with LC_{50} of 0.92 and 0.095 mg AI/L, respectively. Based on the results, deltamethrin had approximately ten times higher toxicity to eggs and two times higher toxicity to adults of the pest compared with abamectin. In contrast abamectin had approximately two times higher toxicity to larvae of the pest compared with deltamethrin. Insecticide concentrations as high

as 3000 mg AI/L did not cause considerable mortality on pupal stage in the dipping method. Hence, their LC values were not estimated in the experiments.

Effects of sublethal doses of abamectin and deltamethrin on life table parameters of *P. operculella* are shown in Table 2. Gross reproductive rate (GRR) in deltamethrin was significantly higher than that for abamectin and control treatments ($F = 6.68$; $df = 2, 57$; $P < 0.01$). Also, both the net reproductive rate (R_0) and intrinsic rate of increase (r_m) in deltamethrin were significantly higher than those in control and abamectin treatments ($F = 2.59$; $df = 2, 57$; $P < 0.01$ and $F = 1.54$; $df = 2, 57$; $P < 0.01$, respectively). There was no significant difference between GRR, estimated R_0 , r_m and λ for control and abamectin treatment. The value of generation time (T) and doubling time (DT) obtained for deltamethrin was significantly lower than those for the control and abamectin treatments ($F = 1.16$; $df = 2, 57$; $P < 0.05$ and $F = 3.7$; $df = 2, 57$; $P < 0.01$, respectively).

Table 1 Toxicity of abamectin and deltamethrin to developmental stages of *Phthorimaea operculella*

Stage	Insecticides	n	$LC_{50}(95\%CL)$ (mg ai/l)	Slope \pm SE	χ^2
Egg	abamectin	360	0.92 ^a (0.59-1.27)	1.03 \pm 0.13	1.87
	deltamethrin	360	0.09 ^b (0.047 – 0.14)	0.94 \pm 0.17	1.2
Larvae	abamectin	360	0.014 ^b (0.006-0.024)	0.74 \pm 0.11	1.81
	deltamethrin	360	0.024 ^a (0.004-0.054)	0.58 \pm 0.13	2.21
Adult	abamectin	360	0.46 ^a (0.29-0.63)	1.72 \pm 0.29	2.28
	deltamethrin	360	0.29 ^b (0.16 – 0.52)	0.86 \pm 0.15	1.55

In each stage, values followed by different letters are significantly different ($P < 0.05$).

Table 2 Life table parameters (mean \pm SE) of *Phthorimaea operculella* exposed to sublethal dose of abamectin and deltamethrin

Parameters	GRR	R_0	r_m	λ	T	DT
Control	21.66 \pm 3.6 ^b	4.41 \pm 1.33 ^b	0.060 \pm 0.014 ^b	1.049 \pm 0.015 ^b	27.3 \pm 0.98 ^a	12.95 \pm 4.84 ^{ab}
Abamectin	12.68 \pm 0.45 ^b	2.83 \pm 0.92 ^b	0.042 \pm 0.016 ^b	1.026 \pm 0.017 ^b	28.89 \pm 0.65 ^a	26.93 \pm 9.04 ^a
Deltamethrin	46.54 \pm 6.5 ^a	20.42 \pm 3.19 ^a	0.141 \pm 0.0066 ^a	1.15 \pm 0.007 ^a	21.74 \pm 0.29 ^b	4.87 \pm 0.23 ^b

Means in the same column, followed by different letters are significantly different ($P < 0.05$).

Discussion

In present study, it was shown that both tested insecticides were effective against PTM. Based on the results, deltamethrin had the highest toxicity to egg and adult stages of PTM and abamectin had the highest toxicity to its first instar larvae. Corbitt *et al.*, (1989) reported that abamectin had highest toxicity on first instar larvae of *Heliothis* Sp. Also Bueno and Freitas (2004) reported that abamectin had no effect on eggs of predator, *Chrysoperla externa*. Thus abamectin could be suitable insecticide in PTM control in IPM. But deltamethrin increased the stable population parameters of PTM specially its intrinsic rate of increase, thus it's not recommended for control of this pest.

Our results demonstrated that the deltamethrin significantly affected life table parameters of *P. operculella*, but abamectin had no significant effect on them. Our results are similar to Bayram *et al.*, (2010) who reported higher intrinsic rate of increase (r_m) in *Telenomus busseolae* after exposure to deltamethrin. Hui-Dong *et al.*, (2004) reported that emamectin caused decreasing parameters including r_m , R_0 , λ , and T of *Plotella xylostella*. Mahdavi *et al.*, (2011) studied sublethal effects of abamectin on the ectoparasitoid *H. hebetor*. Their results were based on extended laboratory studies. Based on these laboratory results it seems that abamectin is potentially more compatible with a chosen IPM approach. Hamed *et al.*, (2011) reported adverse effects of abamectin on population growth of predator *Phytoseius plumifer*, so the results of their study can be used to develop approximate guidelines for the use of abamectin in order to minimize their impact on *P. plumifer* and related natural enemies.

The intrinsic rate of increase (r_m) is the most important parameter for evaluation of total effects of insecticides on population development, because it is based on both survivorship and fecundity (Carey, 1993, Stark and Wennergren, 1995). Higher intrinsic rate of increase (r_m) in deltamethrin indicated that this insecticide has positive effect on population

growth rates in *P. operculella*. Therefore, it is not suitable to control the pest. Even though results obtained in small laboratory conditions may not be realized under natural conditions (Kareiva, 1990), these kinds of investigations under the laboratory conditions can be near to storage conditions and helpful in selecting insecticides for application along with pest and its natural enemies in the pest management programs.

Acknowledgment

We thank the Department of Plant Protection of Mohaghegh Ardabili University, for their financial support and Entomology Research Section of the Iranian Research Institute of Plant Protection for scientific help.

References

- Bayram, A., Salerno, G., Onofri, A. and Conti, E. 2010. Sub-lethal effects of two pyrethroids on biological parameters and behavioural responses to host cues in the egg parasitoid *Telenomus busseolae*. *Biological Control*, 53:153-160.
- Bueno, A. F. Freitas, S., 2004. Effect of the insecticides abamectin and lufenuron on eggs and larvae of *Chrysoperla externa* under laboratory conditions. *Journal of Biological Control*, 49: 277- 283.
- Carey, J. R. 1993. Insect biodemography. *Annual Review of Entomology*, 46: 79-110.
- Corbitt, T. S., StGreen, A. J., Wright, D. J. 1989. Relative potency of abamectin against larval stages of *Spodoptera littoralis* (Boisd), *Heliothis armigera* (Hubn) and *Heliothis virescens* (F) (Lep: Noctuidae). *Crop Protection*, 8: 127-132.
- Delpuech, J. M. and Meyet, J. 2003. Reduction in the sex ratio of the progeny of a parasitoid wasp (*Trichogramma brassicae*) surviving the insecticide chlorpyrifos. *Archives of Environmental Contamination and Toxicology*, 45: 203-208.
- Dogramaci, M., Tingey, W. M. 2008. Comparison of insecticide resistance in a

- north American field population and laboratory colony of potato tuber worm (Lepidoptera: Gelechiidae). *Journal of Pest Science*, 81: 17-22.
- Haines, C. P. and Tropical Products Institute (Great Britain). 1977. The potato tuber moth, *Phthorimaea operculella* (Zeller): a bibliography of recent literature and a review of its biology and control on potatoes in the field and in store. Tropical Products Institute, London.
- Hamed, N., Fathipour, Y. and Saber, M. 2011. Sublethal effects of abamectin on the biological performance of the predatory mite, *Phytoseius plumifer* (Canestrini & Fanzago) (Acari: Phytoseiidae). *Experimental and Applied Acarology*, 53: 29-40.
- Hanafi, A. 1999. Integrated pest management of potato tuber moth in field and storage. *Potato Research*, 42: 373-380.
- Haseeb, M. and Amano, H. 2002. Effects of contact, oral and persistent toxicity of selected pesticides on *Cotesia plutellae* (Hym., Braconidae), a potential parasitoid of *Plutella xylostella* (Lep., Plutellidae). *Journal of Applied Entomology*, 126: 8-13.
- Haug, G. and Hoffman, H. (Eds.). 1990. *Chemistry of Plant Protection 4: Synthetic Pyrethroid Insecticides: Structures and Properties*. Springer-Verlag. Berlin, Heidelberg, New York.
- Hayes, W. J. and Laws, E. R. 1990. *Handbook of Pesticide Toxicology, General Principles*, Vol. 1. Academic Press, Inc., NY.
- Hui-Dong, L., Fang-Qiang, Z. and Wan-Chun, L. 2004. Toxicity of emamectin to the diamondback moth, *Plutella xylostella* and the effects on survivors of parent generation treated with sublethal dosage. *Acta Entomologica Sinica*, 47: 193-197.
- Kareiva, P. 1989. The spatial dimension in pest-enemy interactions. In: Mackauer, Ehler and Roland (Ed.). *Critical Issues in Biological Control*. Intercept, Andover, Intercept Ltd., U. K., pp. 213-227.
- Lashkari, M. R., Sahragard, A. and Ghadamyari, M. 2007. Sublethal effects of imidacloprid and pymetrozin on population growth parameters of cabbage aphid, *Brevicoryne brassicae* on rape seed, *Brassica napus* L. *Insect Science*, 14: 207-212.
- Mahdavi, V., Saber, M., Rafiee-Dastjerdi, H. and Mehrvar, A. 2011. Comparative study of the population level effects of carbaryl and abamectin on larval ectoparasitoid *Habrobracon hebetor* Say (Hymenoptera: Braconidae). *BioControl*, 56: 823-830.
- Meyer, J. S., Ingersoll, C. G., McDonald, L. L. and Boyce, M. S. 1986. Estimating uncertainty in population growth rates: jack knife vs. bootstrap techniques. *Ecology*, 67: 1150-1166.
- Ripper, W. E. 1956. Effect of pesticides on balance of arthropod populations. *Annual Review of Entomology*, 1: 403-438.
- Saour, G. 2008. Effect of thiacloprid against the potato tuber moth *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae). *Journal of Pest Science*, 81: 3-8.
- Schneider, M. I., Smagghe, G., Gobbi, A. and Vinuela, E. 2003. Toxicity and pharmacokinetic of insect growth regulators and other novel insecticides on pupae of *Hyposoter didymator* (Hymenoptera: Ichneumonidae), a parasitoid of early larval instars of lepidopteran pests. *Journal of Economic Entomology*, 96: 1054-1065.
- Sporleder, M., Kroschel, J., Gutierrez-Quispe, M. and Lagnaoui, A. 2004. A temperature-based simulation model for the potato tuberworm, *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae). *Environmental Entomology*, 33: 477-486.
- SPSS. 2004. *SPSS Base 13.0 User's Guide*. SPSS Incorporation, Chicago, IL.
- Stapel, J. O., Cortesero, A. M. and Lewis, W. J. 2000. Disruptive sublethal effects of insecticides on biological control: altered foraging ability and life span of a parasitoid after feeding on extrafloral nectar of cotton treated with systemic insecticides. *Biological Control*, 17: 243-249.
- Stark, J. D. and Banks, E. 2003. Population level effects of pesticides and other toxicants on arthropods. *Annual Review of Entomology*, 48: 505-519.

- Stark, J. D. and Wennergren, U. 1995. Can population effects of pesticides be predicted from demographic toxicological studies? *Journal of Economic Entomology*, 88: 1089-1096.
- Symington, C. A. 2003. Lethal and sublethal effects of pesticides on the potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) and its parasitoid *Orgilus lepidus* Muesebeck (Hymenoptera: Braconidae). *Crop Protection*, 22: 513-519.
- Walthall, W. K. and Stark, J. D. 1996. A comparison of acute mortality and population growth rate as endpoints of toxicological effect. *Ecotoxicology and Environmental Safety*, 37: 45-52.
- Wang, D., Qiu, X. H., Ren, X. X., Zhang, W. Z. and Wang, K. W. 2009. Sublethal effects of spinosad on survival, growth and reproduction of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Pest Management Science*, 65: 223-227.
- White, S. M., Dunbar, D. M., Brown, R., Cartwright, B., Cox, D., Eckel, C., Janson, R. K., Moorkerjee, P. K., Norton, J. A., Peterson, R. F. and Starner, V. R. 1997. Emamectin benzoate: a novel avermectin derivative for control of lepidopterous pests in cotton. *Proceedings of Beltwide Cotton Conference*, PP. 1078-10

اثرات کشندگی و غیر کشندگی آبامکتین و دلتامترین روی بید سیب زمینی *Phthorimaea operculella*

هوشنگ رفیعی دستجردی^{۱*}، زینب مشهدی^۱ و عزیز شیخی گرجان^۲

۱- دانشگاه محقق اردبیلی - دانشکده علوم کشاورزی - گروه گیاه پزشکی

۲- تهران - مؤسسه تحقیقات گیاه پزشکی کشور

* پست الکترونیکی نویسنده مسئول مکاتبه: rafiee@uma.ac.ir

دریافت: ۵ مرداد ۱۳۹۱؛ پذیرش: ۷ مرداد ۱۳۹۲

چکیده: بید سیب زمینی با نام علمی *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) یکی از آفات مهم سیب زمینی در مناطق گرمسیر جهان می باشد. در این تحقیق، اثرات کشندگی و غیر کشندگی دو حشره کش آبامکتین و دلتامترین به بید سیب زمینی در شرایط دمایی 1 ± 26 درجه ی سانتی گراد، رطوبت نسبی 5 ± 65 درصد و دوره ی نوری ۱۶ ساعت روشنایی و ۸ ساعت تاریکی مورد مطالعه قرار گرفت. مقادیر غلظت ها براساس آزمایشات مقدماتی تعیین دز تهیه شدند. آب مقطر به عنوان شاهد مورد استفاده قرار گرفت. مقادیر LC_{50} برای مراحل تخم، لارو سن یک و حشره کامل بید سیب زمینی به ترتیب در آبامکتین معادل ۰/۲۹، ۰/۱۴ و ۰/۴۶ mg ai/l و در دلتامترین معادل ۰/۰۹، ۰/۲۴ و ۰/۲۹ mg ai/l تعیین شدند. تجزیه های آماری نشان داد که دلتامترین سمیت بیشتری به مراحل تخم و حشره کامل و آبامکتین سمیت بیشتری به لارو سن یک بید سیب زمینی داشت. همچنین اثرات دز زیرکشنده LC_{30} دو حشره کش آبامکتین و دلتامترین روی پارامترهای رشد جمعیت بید سیب زمینی در شرایط مذکور تعیین شدند. در شاهد، آبامکتین و دلتامترین نرخ ذاتی افزایش جمعیت به عنوان مهمترین پارامتر به ترتیب معادل ۰/۰۶۰، ۰/۰۴۲ و ۰/۱۴۱ ماده بر ماده بر روز تخمین زده شد. دلتامترین پارامترهای رشد جمعیت بید سیب زمینی به خصوص نرخ ذاتی افزایش جمعیت را افزایش داد، بنابراین برای کنترل بید سیب زمینی توصیه نمی شود. بر اساس نتایج کشندگی و غیر کشندگی، آبامکتین می تواند ترکیب مناسبی برای کنترل بید سیب زمینی باشد.

واژگان کلیدی: بید سیب زمینی، آبامکتین، دلتامترین، پارامترهای رشد جمعیت، اثرات کشندگی و غیر کشندگی